




Helping People To Live Well
“सर्वदा स्वस्थ भवः”



Company Profile

The journey of the 13 years.....

Care Formulation Labs Pvt. Ltd. is providing its services since 2010 fully entered with the aim of distribution and selling of medical products in various medical fields. We are based in Delhi.

Company has been accredited with ISO:2008:2015 and manufacture pharmaceutical formulations as per the guidelines of WHO.

Our success is the result of focus on our clients on top priorities such as quality products, on time delivery, competitive rates and in paralleled awareness.

Now, Care Formulation Labs Pvt. Ltd. is engaged in selling and distributing the pharmaceutical products in 28 states of India and 8 Union Territories and has entered International boundaries as well.

An integrated healthcare solution provider with pharmaceutical product basket of approx. 1000 products.

Marketing and Export of pharmaceutical formulations covering a wide range such as Tablets, Capsules, Injection, Syrups, Dry syrups, Suspensions, Ointments, Creams, Gel, Ear Drops, Eye Drops, Nasal Spray, Rotta Cap, Cosmetics, Protein Powder and Nutritional Supplement etc.

Our products are formulated under the strict supervision of experienced chemists, we are dedicated to maintain the best quality in the products.

We are also engaged in institutional business especially for government and private institutions across the country.

Through a strategy of continuous improvement and teamwork we maintain the highest quality standards, value services and technology.

We assure life long relationship to provide with best of the products.

NAVEEN KUMAR JINDAL
(DIRECTOR)



Accimpak-333

Composition :

Each enteric coated tablet contains :
Acamprosate Calcium 333mg.
Excipients q.s.
Colour : Titanium Dioxide IP

Indication : This medication is used along with counseling and support to help people who are alcohol dependent not drink alcohol. Acamprosate works by restoring the natural balance of chemicals in the brain(neurotransmitters).



10x6 Tablets (Blister)

M.R.P. 76/-Per 6 Tablets

Acethama[®]-200 SR

Composition :

Each uncoated sustained release tablet contains :
Acebrophylline 200mg.
Excipients q.s.
Colour : Titanium Dioxide IP

Indication : Acebrophylline Works By Blocking The Effects Of Leukotriene (substance Produced By The Body In Response To Certain Triggers) That Cause Narrowing And Swelling Of Airways In The Lungs. Acebrophylline Is Used In The Prevention And Treatment Of Asthma And Chronic Obstructive pulmonary disease



10x10 Tablets (Alu Alu)

M.R.P. 175/-Per 10 Tablets

Acethama[®]-DM

Composition :

Each film coated tablet contains :
Acebrophylline 200mg.
(In Sustained Release Form)
Montelukast Sodium IP
Eq. to Montelukast 10mg.
Desloratadine 5mg.
Excipients q.s.
Colour : Sunset Yellow

Indication : Acebrophylline+Desloratadine + Montelukast is used to prevent Asthma. Acebrophylline ,Desloratadine and Montelukast. Acebrophylline is a mucolytic which thins and loosens mucus (phlegm), making it easier to cough out.



10x10 Tablets (Alu Alu)

M.R.P. 280/-Per 10 Tablets

Acethama[®]-FM

Composition :

Each film coated tablet contains :
Acebrophylline 200mg.
(In Sustained Release Form)
Montelukast Sodium IP
Eq. to Montelukast 10mg.
Fexofenadine
Hydrochloride IP 120mg.
Excipients q.s.

Indication : This reduces inflammation in the airways and nose, and improves symptoms of allergies. Fexofenadine is an antiallergic which relieves symptoms like runny nose, watery eyes and sneezing. Acebrophylline is a mucolytic which thins and loosens mucus (phlegm), making it easier to cough out



10x10 Tablets (Alu Alu)

M.R.P. 215/-Per 10 Tablets

Acethama[®]-M

Composition :

Each film coated tablet contains :
Acebrophylline 200mg.
(In Sustained Release Form)
Montelukast Sodium IP
Eq. to Montelukast 10mg.
Excipients q.s.
Colour : Titanium Dioxide IP

Indication : Montelukast + acebrophylline is a combination of two drugs: montelukast (leukotriene receptor antagonist) and acebrophylline (mucolytic bronchodilator). Prevent symptoms of asthma and improves seasonal allergy symptoms.



10x10 Tablets (Alu Alu)

M.R.P. 171/-Per 10 Tablets

Acetynet-600

Composition :

Each film coated tablet contains :
N-Acetylcysteine BP 600mg.
Excipients q.s.
Colour : Titanium Dioxide IP

Indication : When inhaled by mouth, acetylcysteine is used to help thin and loosen mucus in the airways due to certain lung diseases (such as emphysema, bronchitis, cystic fibrosis, pneumonia).This effect helps you to clear the mucus from your lungs so that you can breathe easier.



10x10 Tablets (Alu Alu)

M.R.P. 220/-Per 10 Tablets

Acetynet-AC

Composition :

Each film coated tablet contains :
 N-Acetylcysteine BP 600mg.
 Acebrophylline 100mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : When inhaled by mouth, acetylcysteine is used to help thin and loosen mucus in the airways due to certain lung diseases (such as emphysema, bronchitis, cystic fibrosis, pneumonia). This effect helps you to clear the mucus from your lungs so that you can breathe easier.

10x10 Tablets (Alu Alu)

M.R.P. 210/-Per 10 Tablets



Acetynet-T

Composition :

Each film coated tablet contains :
 N-Acetylcysteine USP 150mg.
 Taurine USP 500mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : When inhaled by mouth, acetylcysteine is used to help thin and loosen mucus in the airways due to certain lung diseases (such as emphysema, bronchitis, cystic fibrosis, pneumonia). This effect helps you to clear the mucus from your lungs so that you can breathe easier.

10x10 Tablets (Alu Alu)

M.R.P. 116.05/-Per 10 Tablets



Aciclovir-400

Composition :

Each uncoated tablet contains :
 Aciclovir IP 400mg.
 Excipients q.s.

Indication : is used to treat infections caused by certain types of viruses. It treats cold sores around the mouth (caused by herpes simplex), shingles (caused by herpes zoster), and chickenpox. This medication is also used to treat outbreaks of genital

20x5 Tablets (Blister)

M.R.P. 68.20/- Per 5 Tablets



Aciclovir-800

Composition :

Each uncoated tablet contains :
 Aciclovir IP 800mg.
 Excipients q.s.

Indication : is used to treat infections caused by certain types of viruses. It treats cold sores around the mouth (caused by herpes simplex), shingles (caused by herpes zoster), and chickenpox. This medication is also used to treat outbreaks of genital

10x10 Tablets (Blister)

M.R.P. 376.20/- Per 10 Tablets



Acimday-250

Composition:

Each uncoated tablet contains :
 Acetazolamide IP 250mg.
 Excipients q.s.

Indication : Acetazolamide used to prevent and reduce the symptoms of altitude sickness. Acetazolamide is also used with other medications to treat a certain type of eye problem (open-angle glaucoma). Acetazolamide is available in generic form.

10x10 Tablets (Blister)

M.R.P. 40/-Per 10 Tablets



Acotry-100

Composition :

Each film coated tablet contains :
 Acotiamide Hydrochloride Hydrate 100mg.
 Excipients q.s.
 Colour : Yellow oxide of Iron & Titanium Dioxide IP

Indication : Acotiamide is used in the treatment of functional dyspepsia. It helps to relieve the symptoms like bloating after a meal, pain/discomfort in the upper abdomen and early satiety.

10x15 Tablets (Blister)

M.R.P. 324/-Per 15 Tablets





TABLETS

Alendate-70

Composition :

Each uncoated tablet contains :
Sodium Alendronate Trihydrate USP 70mg
Eq.to Alendronic Acid
Excipients q.s.

Indication : Alendronic acid, sold under the brand name Fosamax among others, is a bisphosphonate medication used to treat osteoporosis and Paget's disease of bone. It is taken by mouth. Use is often recommended together with vitamin D, calcium supplementation, and lifestyle changes




10x1x4 Tablets (Blister)
M.R.P. 165/-Per 4 Tablets

Alfuzosin-D

Composition :

Each film coated bilayered tablet contains :
Alfuzosin Hydrochloride IP 10mg.
(In extended release form)
Dutasteride IP 0.5mg.
Excipients q.s.
Colour: Yellow oxide of Iron

Indication : Alfuzosin only comes as an extended-release oral tablet. Alfuzosin is used to treat benign prostatic hyperplasia (BPH) in adult men. It helps relax the muscles in your prostate and bladder, which can reduce BPH symptoms and improve your ability to urinate.



10x10 Tablets (Alu Alu)
M.R.P. 239/-Per 10 Tablets

Amedron-100

Composition :

Each uncoated tablet contains
Amiodarone Hydrochloride IP 100mg.
Excipients q.s.

Indication : This medication is used to treat certaintypes of serious (possibly fatal) irregularheartbeat (such as persistent ventricular fibrillation/ tachycardia).



10x10 Tablets (Alu Alu)
M.R.P. 68.70/-Per 10 Tablets

Alfuzosin-10

Composition :

Each film coated prolonged release tablet contains :
Alfuzosin Hydrochloride IP 10mg.
Excipients q.s.
Colour: Titanium Dioxide IP

Indication : Alfuzosin only comes as an extended-release oral tablet. Alfuzosin is used to treat benign prostatic hyperplasia (BPH) in adult men. It helps relax the muscles in your prostate and bladder, which can reduce BPH symptoms and improve your ability to urinate.



10x10 Tablets (Alu Alu)
M.R.P. 165/-Per 10 Tablets

Alpha-Ketozox

COMPOSITION:

Each film coated tablet contains:
Calcium-3-methyl-2-oxovalerate
(α - Ketoanalogue of isoleucine, Calcium salt) 67 mg.
Calcium-4-methyl-2-oxovalerate
(α - Ketoanalogue of leucine, Calcium salt) 101 mg.
Calcium-2-oxo-3-phenylpropionate
(α - Ketoanalogue of phenylalanine, Calcium salt) 68 mg.
Calcium-3-methyl-2-oxobutyrate
(α - Ketoanalogue of valine, Calcium salt) 88mg.
Calcium-DL-2-hydroxy-4-(methylthio)-butyrate
(α - hydroxylanalogue of methionine, Calcium-salt) 59mg.
L-Lysine Acetate USP 105 mg.
L-Threonine USP 53 mg.
L-Tryptophan USP 23 mg.
L-Histidine USP 38 mg.
L-Tyrosine USP 30 mg.
Total Nitrogen content / tablet 35mg.
Calcium / tablet 1.25 mmol \approx 0.05 g.
Colour: Ponceau 4R



10x10 Tablets (Alu Alu)
M.R.P. 410/-Per 10 Tablets

Amedron-200

Composition :

Each uncoated tablet contains
Amiodarone Hydrochloride IP 200mg.
Excipients q.s.

Indication : This medication is used to treat certaintypes of serious (possibly fatal) irregularheartbeat (such as persistent ventricular fibrillation/ tachycardia).



10x10 Tablets (Alu Alu)
M.R.P.115.02/-Per 10 Tablets

Amiport-100

Composition:

Each uncoated tablet contains :
 Amisulpride IP 100mg.
 Excipients q.s.

Indication : Amisulpride, sold under the brand name Solian among others, is an antipsychotic medication used to treat schizophrenia. It is also used to treat dysthymia. It is usually classed with the atypical antipsychotics.



10x10 Tablets (Blister)
 M.R.P. 180/-Per 10 Tablets

Amiport-200

Composition:

Each uncoated tablet contains :
 Amisulpride IP 200mg.
 Excipients q.s.

Indication : Amisulpride, sold under the brand name Solian among others, is an antipsychotic medication used to treat schizophrenia. It is also used to treat dysthymia. It is usually classed with the atypical antipsychotics.





10x10 Tablets (Blister)
 M.R.P. 220/-Per 10 Tablets

Amitocare-10

Composition :

Each film coated tablet contains :
 Amitriptyline Hydrochloride I.P 10 mg.
 Excipients q.s.
 Colour :Aluminium Lake of Indigo

Indication : Amitriptyline is a medicine used for treating pain. You can take it: to treat nerve pain (neuralgia) and back pain. to help prevent migraine attacks.





4x5x10 Tablets (Blister)
 M.R.P. 23.85/-Per 10 Tablets

Amitocare-25

Composition :

Each film coated tablet contains :
 Amitriptyline Hydrochloride I.P 25 mg.
 Excipients q.s.
 Colour :Aluminium Lake of Indigo

Indication : Amitriptyline is a medicine used for treating pain. You can take it: to treat nerve pain (neuralgia) and back pain. to help prevent migraine attacks.



4x5x10 Tablets (Blister)
 M.R.P. 24.19/-Per 10 Tablets

Amlodose-2.5

Composition :

Each uncoated tablet contains:
 Amlodipine Besylate I.P.
 eq. to Amlodipine 2.5 mg.
 Excipients q.s.

Indication : Amlodipine is used alone or in combination with other medications to treat high blood pressure in adults and children 6 years and older. It is also used to treat certain types of angina (chest pain) and coronary artery disease (narrowing of the blood vessels that supply blood to the heart).



4x5x10 Tablets (Blister)
 M.R.P. 20.04/-Per 10 Tablets

Amlodose-5

Composition :

Each uncoated tablet contains:
 Amlodipine Besylate I.P.
 eq. to Amlodipine 5 mg.
 Excipients q.s.

Indication : Amlodipine is used alone or in combination with other medications to treat high blood pressure in adults and children 6 years and older. It is also used to treat certain types of angina (chest pain) and coronary artery disease (narrowing of the blood vessels that supply blood to the heart).




10x5x10 Tablets (Blister)
 M.R.P. 24.97/-Per 10 Tablets

Amlodose-10

Composition :

Each uncoated tablet contains:
Amlodipine Besylate I.P. 10 mg.
eq. to Amlodipine q.s.
Excipients q.s.

Indication : Amlodipine is used alone or in combination with other medications to treat high blood pressure in adults and children 6 years and older. It is also used to treat certain types of angina (chest pain) and coronary artery disease (narrowing of the blood vessels that supply blood to the heart).



4x5x10 Tablets (Blister)
M.R.P. 60.92/-Per 10 Tablets

Anazos-1

Composition :

Each film coated tablet contains :
Anastrozole USP 1mg.
Excipients q.s.
Colour : Titanium Dioxide USNF

Indication : Anastrozole is a type of hormone treatment. It works by lowering the levels of oestrogen hormones in your body. It is mainly prescribed for women who have been through the menopause and have a type of cancer called hormone-dependent breast cancer.



3x10 Tablets (Blister)
M.R.P. 670/-Per 10 Tablets

Apxban-5

Composition :

Each film coated tablet contains :
Apxaban 5mg.
Excipients q.s.
Colour : Titanium Dioxide IP & Red oxide of Iron

Indication : Apxaban is used help prevent strokes or blood clots in people who have atrial fibrillation (a condition in which the heart beats irregularly, increasing the chance of clots forming in the body and possibly causing strokes) that is not caused by heart valve disease.




10x1x10 Tablets (Blister)
M.R.P. 695/-Per 10 Tablets

Amlodose-AT

Composition :

Each uncoated tablet contains :
Amlodipine Besylate I.P. 5 mg.
eq. to Amlodipine 5 mg.
Atenolol I.P. 50 mg.
Excipients q.s.
Colour : Sunset Yellow FCF

Indication : Amlodipine + Atenolol is a combination of two medicines: Amlodipine and Atenolol, which lowers blood pressure effectively. Amlodipine is a calcium channel blocker which works by relaxing blood vessels while atenolol is a beta blocker which works specifically on the heart to slow down the heart rate.



20x14 Tablets (Aluminium)
M.R.P. 60.76/-Per 14 Tablets

Apxban-2.5

Composition :

Each film coated tablet contains :
Apxaban 2.5mg.
Excipients q.s.
Colour : Titanium Dioxide IP & Yellow oxide of Iron

Indication : Apxaban is used help prevent strokes or blood clots in people who have atrial fibrillation (a condition in which the heart beats irregularly, increasing the chance of clots forming in the body and possibly causing strokes) that is not caused by heart valve disease.



10x1x10 Tablets (Blister)
M.R.P. 349/-Per 10 Tablets

Aripiz-10

Composition :

Each uncoated tablets contains :
Aripiprazole IP 10mg.
Excipients q.s.

Indication : Aripiprazole is used to treat certain mental/mood disorders (such as bipolar disorder, schizophrenia, Tourette's syndrome, and irritability associated with autistic disorder). It may also be used in combination with other medication to treat depression. Aripiprazole is known as an antipsychotic drug (atypical type).



10x10 Tablets (Alu Alu)
M.R.P. 110/-Per 10 Tablets

Ausporate Chorono-500

Composition :
Each film coated extended release tablet contains :
Sodium Valproate IP 333mg
Valproic Acid IP 145mg
(Both together correspond Sodium Valproate 500mg)
Excipients q.s.
Colours : Red Oxide of Iron & Titanium Dioxide I.P.

Valproate (VPA) and its valproic acid, sodium valproate, and valproate semisodium forms are medications primarily used to treat epilepsy and bipolar disorder and prevent migraine headaches. They are useful for the prevention of seizures in those with absence seizures, partial seizures, and generalized seizures.

10x10 Tablets (Alu Alu)
M.R.P. 103/-Per 10 Tablets



Ausporate Chorono-300

Composition :
Each film coated extended release tablet contains :
Sodium Valproate IP 200mg
Valproic Acid IP 87mg
(Both together correspond Sodium Valproate 300mg)
Excipients q.s.
Colours : Red Oxide of Iron & Titanium Dioxide I.P.

Valproate (VPA) and its valproic acid, sodium valproate, and valproate semisodium forms are medications primarily used to treat epilepsy and bipolar disorder and prevent migraine headaches. They are useful for the prevention of seizures in those with absence seizures, partial seizures, and generalized seizures.

10x10 Tablets (Alu Alu)
M.R.P. 68/-Per 10 Tablets



Ausporate-200

Composition :
Each enteric coated tablet contains :
Sodium Valproate IP 200 mg.
Excipients q.s.
Colour : Sunset Yellow

Indication : Sodium valproate is used to treat epilepsy and bipolar disorder. It's occasionally used to prevent migraine headaches. This medicine is only available on prescription. It comes as capsules, tablets and a liquid that you swallow. seizures, and generalized seizures.

10x10 Tablets (Alu Alu)
M.R.P. 34.60/-Per 10 Tablets




Ausric-100

Composition :
Each uncoated tablet contains :
Allopurinol IP 100mg.
Excipients q.s.

Indication : Allopurinol is used to treat gout and certain types of kidney stones. It is also used to prevent increased uric acid levels in patients receiving cancer chemotherapy. These patients can have increased uric acid levels due to release of uric acid from the dying cancer cells.

10x10 Tablets (Blister)
M.R.P. 18.92/-Per 10 Tablets



Ausric-300

Composition :
Each uncoated tablet contains :
Allopurinol IP 300mg.
Excipients q.s.

Indication : Allopurinol is used to treat gout and certain types of kidney stones. It is also used to prevent increased uric acid levels in patients receiving cancer chemotherapy. These patients can have increased uric acid levels due to release of uric acid from the dying cancer cells.

10x10 Tablets (Blister)
M.R.P. 62.94/-Per 10 Tablets



Austavir 0.5

Composition :
Each film coated tablet contains:
Entecavir IP 0.5mg.
(As monohydrate)
Excipients q.s.
Colour : Titanium Dioxide IP

Indication : Entecavir is used to treat chronic (long-term) hepatitis B infection (swelling of the liver caused by a virus) in adults and children 2 years of age and older who have liver damage. Entecavir is in a class of medications called nucleoside analogs.

10x1x10 Tablets (Alu Alu)
M.R.P. 760/-Per 10 Tablets





Austavir-1

Composition :

Each film coated tablet contains:
Entecavir IP 1mg.
(As monohydrate)
Excipients q.s.
Colour : Titanium Dioxide IP

Indication : Entecavir is used to treat chronic (long-term) hepatitis B infection (swelling of the liver caused by a virus) in adults and children 2 years of age and older who have liver damage. Entecavir is in a class of medications called nucleoside analogs.



10x10 Tablets (Alu Alu)
M.R.P. 1367.30/-Per 10 Tablets

Azyihyp-40

Composition :

Each film coated tablet contains :
Azilsartan Kamedoxomil
Eqv. to Azilsartan Medoxomil 40 mg.
Excipients q.s.
Colours : Quinoline Yellow Lake
& Titanium Dioxide IP

Indication : Azilsartan is used to treat high blood pressure (hypertension). Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. Azilsartan belongs to a class of drugs called angiotensin receptor blockers (ARBs). It works by relaxing blood vessels so that blood can flow more easily.



10x1x10 Tablets (Alu Alu)
M.R.P. 135/-Per 10 Tablets

Baclonip-10

Composition :

Each uncoated tablet contains :
Baclofen IP 10 mg.
Excipients q.s.

Indication : Baclofen is used to treat muscle spasms caused by certain conditions (such as multiple sclerosis, spinal cord injury/disease). It works by helping to relax the muscles.



10x10 Tablets (Blister)
M.R.P. 124.84/-Per 10 Tablets

Aboyvir-200

Composition :

Each uncoated tablet contains :
Aciclovir IP 200mg.
Excipients q.s.

Indication : Acyclovir is used to decrease pain and speed the healing of sores or blisters in people who have varicella (chickenpox), herpes zoster (shingles; a rash that can occur in people who have had chickenpox in the past), and first-time or repeat outbreaks of genital herpes (a herpes virus infection that causes sores to form around the genitals and rectum from time to time).



10x10 Tablets (Blister)
M.R.P. 84.22/-Per 10 Tablets

Azathioprine-50

Composition :

Each uncoated tablet contains :
Azathioprine IP 50mg.
Excipients q.s.

Indication : Azathioprine belongs to the group of medications known as immunosuppressive agents. It is used to prevent the rejection of kidney transplants and to treat severe rheumatoid arthritis that is not responding to conventional treatment.



10x10 Tablets (Blister)
M.R.P. 105.40/-Per 10 Tablets

Bendicop-4

Composition :

Each film coated tablet contains :
Benidipine Hydrochloride IP 4 mg.
Excipients q.s.

Indication : Benidipine is a dihydropyridine calcium channel blocker that is used for the treatment of high blood pressure. This makes the tachycardia minimal.



10x10 Tablets (Alu Alu)
M.R.P. 80/-Per 10 Tablets

Bendicop-8

Composition :

Each film coated tablet contains :
 Bendipine Hydrochloride IP 8 mg.
 Excipients q.s.

Indication : Bendipine is a dihydropyridine calcium channel blocker that is used for the treatment of high blood pressure. This makes the tachycardia minimal.



10x10 Tablets (Alu Alu)

M.R.P. 145/-Per 10 Tablets

Bicogone-2.5

Composition :

Each uncoated tablet contains :
 Bisoprolol Fumarate USP 2.5mg.
 Excipients q.s.

Indication : Bisoprolol is used with or without other medications to treat high blood pressure (hypertension). Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. This medication belongs to a class of drugs known as beta blockers.



10x10 Tablets (Blister)

M.R.P. 62/-Per 10 Tablets

Blosia-4

Composition :

Each uncoated tablet contains :
 Blonanserin 4mg.
 Excipients q.s.

Colour : Erythrosine

Indication : Blonanserin is an anti-psychotic drug which is used to treat patients with mental disorders like schizophrenia and bipolar disorder. It works by altering the balance of certain chemicals in the brain, so as to relieve the symptoms of schizophrenia and bipolar disorder.



10x10 Tablets (Alu Alu)

M.R.P. 130/-Per 10 Tablets

Bethanax-25

Composition :

Each uncoated tablet contains :
 Bethanechol Chloride USP 25mg.
 Excipients q.s.

Indication : This medication is used to treat certain bladder problems such as the inability to urinate or empty the bladder completely due to certain causes (e.g., surgery, bladder muscle problems). It works by helping the bladder muscle to squeeze better, thereby improving your ability to urinate



10x10 Tablets (Aluminium Strip)

M.R.P. 145/-Per 10 Tablets

Bicogone-5

Composition :

Each uncoated tablet contains :
 Bisoprolol Fumarate USP 5mg.
 Excipients q.s.

Indication : Bisoprolol is used with or without other medications to treat high blood pressure (hypertension). Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. This medication belongs to a class of drugs known as beta blockers.



10x10 Tablets (Blister)

M.R.P. 66/-Per 10 Tablets

Blosia-8

Composition :

Each uncoated tablet contains :
 Blonanserin 8mg.
 Excipients q.s.

Indication : Blonanserin is an anti-psychotic drug which is used to treat patients with mental disorders like schizophrenia and bipolar disorder. It works by altering the balance of certain chemicals in the brain, so as to relieve the symptoms of schizophrenia and bipolar disorder.



10x10 Tablets (Alu Alu)

M.R.P. 195/-Per 10 Tablets

Bisalto-20

Composition :

Each uncoated tablet contains :
 Bilastine 20mg.
 Excipients q.s.



Indication : Bilastine, a non-sedating H1-antihistamine, is indicated to treat the symptoms of allergic disorders (e.g. rhinoconjunctivitis and urticaria) in adults and adolescents and, more recently, in children.

10x10 Tablets (Alu Alu)

M.R.P. 145/-Per 10 Tablets

Bisalto-M

Composition :

Each film coated tablet contains :
 Bilastine 20mg.
 Montelukast Sodium IP eq. to Montelukast 10mg.
 Excipients q.s.
 Colour : Red oxide of Iron, Yellow Oxide of Iron & Titanium Dioxide IP



Indication : Bilastine belongs to the class of medications called antihistamines. It may be used by people 12 year of age and older to treat symptoms of seasonal allergies, including sneezing, itchy and runny nose; itchy, red and watery eyes; and skin rash and irritations.

10x10 Tablets (Alu-Alu)

M.R.P. 245/-Per 10 Tablets

Bosentan-62.5

Composition

Each film coated tablet contains :
 Bosentan Monohydrate eq. to Bosentan 62.5mg.
 Excipients q.s.



Indication : Bosentan is used to treat pulmonary arterial hypertension, which is high blood pressure of the blood vessels between the heart and lungs. Bosentan works to block the effect of a natural compound called endothelin, which is something the body makes to tighten blood vessels.

10x1x10 Tablets (Blister)

M.R.P. 798.52/-Per 10 Tablets

Bisalto-40

Composition :

Each uncoated tablet contains :
 Bilastine 40mg.
 Excipients q.s.



Indication : Bilastine, a non-sedating H1-antihistamine, is indicated to treat the symptoms of allergic disorders (e.g. rhinoconjunctivitis and urticaria) in adults and adolescents and, more recently, in children.

10x10 Tablets (Alu Alu)

M.R.P. 265/-Per 10 Tablets

Bivogrid-50

Composition

Each film-coated tablet contains :
 Brivaracetam 50mg.
 Excipients q.s.
 Colour : Red Oxide of Iron Lake Yellow Oxide of Iron & Titanium



Brivaracetam is used along with other medications to control partial onset seizures (seizures that involve only one part of the brain) in adults and children 4 years of age and older. Brivaracetam is in a class of medications called anticonvulsants.

10x1x10 Tablets (Blister)

M.R.P. 242/-Per 10 Tablets

Bosentan-125

Composition

Each film coated tablet contains :
 Bosentan Monohydrate eq. to Bosentan 125mg.
 Excipients q.s.



Indication : Bosentan is used to treat pulmonary arterial hypertension, which is high blood pressure of the blood vessels between the heart and lungs. Bosentan works to block the effect of a natural compound called endothelin, which is something the body makes to tighten blood vessels.

10x1x10 Tablets (Blister)

M.R.P. 1590/-Per 10 Tablets

Bromcrep 2.5

Composition :

Each uncoated tablet contains :
 Bromocriptine Mesylate IP
 eq. to Bromocriptin 2.5mg.
 Excipients q.s.

Indication : The most recently FDA-approved indication for bromocriptine mesylate is a supplemental medication for adults with type 2 diabetes mellitus (type 2 DM) to help with glycemic control.



10x1x10 Tablets (Aluminium)
 M.R.P. 499/-Per 10 Tablets

Buskron-10

Composition :

Each uncoated tablet contains :
 Buspirone Hydrochloride IP
 Eq. to Buspirone 10mg.
 Excipients q.s.
 Colour : Brilliant Blue

Indication : Buspirone is primarily used to treat generalized anxiety disorder (GAD). It is an FDA-approved medicine for managing anxiety disorders or the short-term relief of anxiety symptoms. Off-labeled buspirone is used for the augmentation of unipolar depression.



10x10 Tablets (Blister)
 M.R.P. 75/-Per 10 Tablets

Cabergoline 0.5

Composition :

Each uncoated tablet contains:
 Cabergoline IP 0.5mg.
 Excipients q.s.

Indication : Cabergoline is a dopamine receptor agonist. It works by reducing the amount of prolactin (a hormone) that is released from the pituitary gland. Cabergoline is used to treat a hormone imbalance in which there is too much prolactin in the blood (also called hyperprolactinemia).



10x1x4 Tablets (Aluminium)
 M.R.P. 399/-Per 4 Tablets

Buskron-5

Composition :

Each uncoated tablet contains :
 Buspirone Hydrochloride IP
 Eq. to Buspirone 5mg.
 Excipients q.s.
 Colour : Brilliant Blue

Indication : Buspirone is primarily used to treat generalized anxiety disorder (GAD). It is an FDA-approved medicine for managing anxiety disorders or the short-term relief of anxiety symptoms. Off-labeled buspirone is used for the augmentation of unipolar depression.



10x10 Tablets (Blister)
 M.R.P. 55/-Per 10 Tablets

Byluta-50

Composition :

Each film coated tablet contains:
 Bicalutamide IP 50mg.
 Excipients q.s.
 Colours : Titanium Dioxide IP

Indication : Bicalutamide is used for the treatment of advanced prostate carcinoma. It is taken together with a drug known as a luteinising hormone-releasing hormone (LHRH) analogue which reduces the levels of androgens (male sex hormones) within the body, or with accompanying surgical removal of the testicles.



3x10 Tablets (Blister)
 M.R.P. 432.50/-Per 10 Tablets

Cabergoline-0.25

Composition :

Each uncoated tablet contains:
 Cabergoline IP 0.25mg.
 Excipients q.s.

Indication : Cabergoline is a dopamine receptor agonist. It works by reducing the amount of prolactin (a hormone) that is released from the pituitary gland. Cabergoline is used to treat a hormone imbalance in which there is too much prolactin in the blood (also called hyperprolactinemia).



10x10 Tablets (Aluminium)
 M.R.P. 249/-Per 4 Tablets



TABLETS

Cansovir-250

Composition :

Each film coated tablet contains:
Lopinavir IP 200mg.
Ritonavir IP 50mg.
Excipients q.s.

Indication : The combination of lopinavir and ritonavir is used with other medications to treat human immunodeficiency virus (HIV) infection. Lopinavir and ritonavir are in a class of medications called protease inhibitors. They work by decreasing the amount of HIV in the blood.



1x30 Tablets

M.R.P. 1510/-Per 30 Tablets

Cardifill-7.5

Composition :

Each film coated tablet contains :
Ivabradine Hydrochloride IP 7.5mg.
Excipients q.s.

Indications : This medication is used by people with a certain heart problem (chronic heart failure), to help prevent it from getting worse and needing treatment in a hospital. Heart failure is a condition where your heart does not pump blood as well as it should. Ivabradine works by making your heart beat more slowly.



10x1x10 Tablets (Alu Alu)


M.R.P. 266/-Per 10 Tablets

Carelix[®]-SR 1.5

Composition :

Each film coated sustained release tablet contains :
Indapamide IP 1.5mg.
Excipients q.s.

Indication : Indapamide is also used to reduce extra fluid in the body (edema) caused by heart failure. Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. Indapamide belongs to a class of drugs known as diuretics/"water pills." It works by causing you to make more urine.



10x10 Tablets (Alu Alu)

M.R.P. 110/-Per 10 Tablets

Cardifill-5

Each film coated tablet contains :

Ivabradine Hydrochloride IP 5mg.
Excipients q.s.

Indications : This medication is used by people with a certain heart problem (chronic heart failure), to help prevent it from getting worse and needing treatment in a hospital. Heart failure is a condition where your heart does not pump blood as well as it should. Ivabradine works by making your heart beat more slowly.



10x1x10 Tablets (Alu Alu)

M.R.P. 210/-Per 10 Tablets

Caldicare-Ace 667

Composition :

Each uncoated tablet contains :
Calcium Acetate USP 667mg.
Excipients q.s.

Indication : Calcium acetate is used to prevent high blood phosphate levels in patients who are on dialysis due to severe kidney disease. Dialysis removes some phosphate from your blood, but it is difficult to remove enough to keep your phosphate levels balanced.



10x10 Tablets (Blister)

M.R.P.120/-Per 10 Tablets

Carelix[®]

Composition :

Each uncoated tablet contains :
Sodium Picosulphate B.P. 10 mg.
Excipients q.s.

Indication : Sodium picosulfate/magnesium oxide/citric acid solution is used to clean out the intestines before surgery or certain procedures (such as colonoscopy, X-rays). It is a laxative that works by drawing large amounts of water into your intestines and stimulating the colon. This causes watery bowel movements (diarrhea).



4x5x10 Tablets (Blister)

M.R.P. 51/-Per 10 Tablets

Care Zinc

Composition :

Each uncoated dispersible tablet contains :
 Zinc Sulphate (Monohydrate) IP 20mg.
 Eq. to Elemental Zinc 20mg.
 Excipients q.s.

Indication : Zinc is used by some people to help with the common cold or lung infections, malaria, or asthma. It may help with wound healing, ulcers, acne, and skin infections. Some people will use zinc to help with age-related problems like macular degeneration or Alzheimer disease. Others will use it to help with loose stools.



4x5x10 Tablets (Blister)

M.R.P. 35/-Per 10 Tablets

Carkozom-5

Composition :

Each uncoated tablet contains :
 Carbimazole IP 5mg.
 Excipients q.s.

Indication : Carbimazole is used to reduce the formation of thyroid hormones. Treatment in more serious cases, for example, to restore the normal function of the thyroid before its partial removal by surgery. It is also used together with other treatments for overactive thyroid.



1x100 Tablets

M.R.P. 237.44/-Per 100 Tablets

Carkozom-20

Composition :

Each uncoated tablet contains :
 Carbimazole IP 20mg.
 Excipients q.s.

Indication : Carbimazole is used to reduce the formation of thyroid hormones. Treatment in more serious cases, for example, to restore the normal function of the thyroid before its partial removal by surgery. It is also used together with other treatments for overactive thyroid.



1x100 Tablets

M.R.P. 892/-Per 100 Tablets

Caregrain

Composition :

Each uncoated tablet contains :
 Caffeine (Monohydrate) IP 100mg.
 Ergotamine Tartrate IP 1mg.
 Paracetamol IP 250mg.
 Prochlorperazine Maleate IP 2.5mg.
 Excipients q.s.
 Colour : Brilliant Blue FCF



Indication : is used to relieve the symptoms at the first sign of a migraine attack and to prevent the worsening of the attack.

10x1x10 Tablets (Blister)

M.R.P. 150/-Per 10 Tablets

Carkozom-10

Composition :

Each uncoated tablet contains :
 Carbimazole IP 10mg.
 Excipients q.s.

Indication : Carbimazole is used to reduce the formation of thyroid hormones. Treatment in more serious cases, for example, to restore the normal function of the thyroid before its partial removal by surgery. It is also used together with other treatments for overactive thyroid.



1x100 Tablets

M.R.P. 477.1/-Per 100 Tablets

Carzibo-25

Composition :

Each uncoated tablet contains :
 Acarbose IP 25mg.
 Excipients q.s.



Indication : Acarbose is used (with diet only or diet and other medications) to treat type 2 diabetes (condition in which the body does not use insulin normally and therefore cannot control the amount of sugar in the blood).

10x10 Tablets (Aluminium Strip)

M.R.P. 70/-Per 10 Tablets




Carzibo-50

Composition :

Each uncoated tablet contains :
Acarbose IP 50mg
Excipients q.s.

Indication : Acarbose is used (with diet only or diet and other medications) to treat type 2 diabetes (condition in which the body does not use insulin normally and therefore cannot control the amount of sugar in the blood).



10x10 Tablets (Aluminium Strip)
M.R.P. 152/-Per 10 Tablets

Chlotado 12.5

Composition :

Each film coated tablet contains :
Chlorthalidone IP 12.5mg.
Excipients q.s.

Indication : Chlorthalidone is used to treat high blood pressure (hypertension). Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. It is also used to reduce extra salt and water in the body caused by conditions such as heart failure, liver disease, and kidney disease.




10x10 Tablets (Blister)
M.R.P. 59/-Per 10 Tablets

Chymury-Forte

Composition :

Each enteric coated tablet contains :
1,00,000 Armour units of enzymatic activity (supplied by a purified concentrate which has specific trypsin and chymotrypsin activity in a ratio of approximately six to one)
Colours : Ponceau 4R & Sunset Yellow FCF

Indication : People use chymotrypsin to make medicine. People take chymotrypsin by mouth or as a shot to reduce redness and swelling associated with pockets of infection (abscesses), ulcers, surgery, or traumatic injuries; and to help loosen phlegm in asthma, bronchitis, lung diseases, and sinus infections.



10x20 Tablets (Blister)
M.R.P. 499/-Per 20 Tablets

Chlotado 6.25

Composition :

Each film coated tablet contains :
Chlorthalidone IP 6.25mg.
Excipients q.s.

Indication : Chlorthalidone is used to treat high blood pressure (hypertension). Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. It is also used to reduce extra salt and water in the body caused by conditions such as heart failure, liver disease, and kidney disease.



10x10 Tablets (Blister)
M.R.P. 56/-Per 10 Tablets

Chymury

Composition :

Each enteric coated tablet contains :
Trypsin BP 48mg.
Bromelain 90mg.
Rutoside Trihydrate BP 100mg.
Diclofenac Sodium IP 50mg.
Colour : Sunset Yellow & Titanium Dioxide IP

Indication : a combination of Trypsin, Bromelain, Rutoside Trihydrate and Diclofenac is a nonsteroidal anti-inflammatory painkiller. Dicoheal is used to reduce pain, swelling and improve knee function in people with arthritis.



10x10 Tablets (Alu-Alu)
M.R.P. 199/-Per 10 Tablets

Capedoz-500

Composition :

Each film coated tablet contains :
Capecitabine IP 500mg.
Excipients q.s.
Colours: Red oxide of Iron & titanium Dioxide IP

Indication : Capecitabine is a type of chemotherapy. It is used as a treatment for different types of cancer, including bowel cancer. You might have capecitabine on its own or with other chemotherapy drugs. You might also have it with radiotherapy.



3x10 Tablets (Blister)
M.R.P. 1125/-Per 10 Tablets

Ciloshot-100

Composition :

Each uncoated tablet contains :
 Cilostazol IP 100 mg.
 Excipients q.s.

Indication : Cilostazol is used to reduce the symptoms of intermittent claudication (pain in the legs that worsens when walking and improves when resting that is caused by narrowing of the blood vessels that supply blood to the legs).



10x10 Tablets (Alu Alu)

M.R.P. 263/-Per 10 Tablets

Ciloshot-50

Composition :

Each uncoated tablet contains :
 Cilostazol IP 50 mg.
 Excipients q.s.

Indication : Cilostazol is used to reduce the symptoms of intermittent claudication (pain in the legs that worsens when walking and improves when resting that is caused by narrowing of the blood vessels that supply blood to the legs).



10x10 Tablets (Alu Alu)

M.R.P. 140/-Per 10 Tablets

Citirise[®]-Plus

Composition :

Each film coated tablet contains :
 Citicoline Sodium IP
 Eq. to Citicoline 500mg.
 Piracetam IP 800mg.
 Excipients q.s.
 Colours: Titanium Dioxide IP

Indication : Citicoline + Piracetam is generally prescribed for memory enhancement, neurological and cognitive disorders and conditions such as Stroke, Parkinsonism and Alzheimer diseases.



10x10 Tablets (Alu Alu)

M.R.P. 670/-Per 10 Tablets

Citirise[®]

Composition :

Each film coated tablet contains :
 Citicoline Sodium IP
 Eq. to Citicoline 500mg.
 Excipients q.s.
 Colours: Red Oxide of Iron & Titanium Dioxide IP

Indication : Citicoline is taken by mouth or given as an injection to help memory loss due to aging, improve vision in people with glaucoma, and help with recovery in stroke patients. It is also used for Alzheimer disease, Parkinson disease, bipolar disorder, lazy eye, and other conditions of the brain.



10x10 Tablets (Alu Alu)

M.R.P. 540/-Per 10 Tablets

Clindoz-5

Composition :

Each film coated tablet contains:
 Cilnidipine IP 5mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Cilnidipine due to its blocking action at N-type and L type calcium channel dilates both arterioles and venules as a result the pressure in the capillary bed is reduced.



10x10 Tablets (Alu Alu)

M.R.P. 50/-Per 10 Tablets

Clindoz-10

Composition :

Each film coated tablet contains:
 Cilnidipine IP 10mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Cilnidipine due to its blocking action at N-type and L type calcium channel dilates both arterioles and venules as a result the pressure in the capillary bed is reduced.



10x10 Tablets (Alu Alu)

M.R.P. 71/-Per 10 Tablets

Clindoz-20

Composition :

Each film coated tablet contains:
 Cilnidipine IP 20mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP



Indication : Cilnidipine due to its blocking action at N-type and L type calcium channel dilates both arterioles and venules as a result the pressure in the capillary bed is reduced.

10x10 Tablets (Alu Alu)

M.R.P. 120/-Per 10 Tablets

Clomish-50

Composition :

Each uncoated tablet contains
 Clomiphene Citrate IP 50mg.
 Excipients q.s.



Indication : Clomiphene is used to induce ovulation (egg production) in women who do not produce ova (eggs) but wish to become pregnant (infertility). Clomiphene is in a class of medications called ovulatory stimulants. It works similarly to estrogen, a female hormone that causes eggs to develop in the ovaries and be released.

10x10 Tablets (Blister)

M.R.P. 98.8/-Per 10 Tablets

Clopogril-75

Composition :

Each film coated tablet contains :
 Clopidogrel Bisulphate IP 75mg.
 Eq. to Clopidogrel q.s.
 Colour : Red Oxide of Iron & Titanium Dioxide IP



Indication : (clopidogrel bisulfate) is a thienopyridine class of drug that inhibits platelet aggregation and thus inhibits aspects of blood clotting used to treat patients with acute coronary syndrome, myocardial infarction (MI), peripheral vascular disease and some stroke (ischemic type) patients.

20x10 Tablets (Alu Alu)

M.R.P. 66.52/-Per 10 Tablets

Clomish-25

Composition :

Each uncoated tablet contains
 Clomiphene Citrate IP 25mg.
 Excipients q.s.



Indication : Clomiphene is used to induce ovulation (egg production) in women who do not produce ova (eggs) but wish to become pregnant (infertility). Clomiphene is in a class of medications called ovulatory stimulants. It works similarly to estrogen, a female hormone that causes eggs to develop in the ovaries and be released.

10x10 Tablets (Blister)

M.R.P. 63.25/-Per 10 Tablets

Clonidoc-100

Composition :

Each uncoated tablet contains:
 Clonidine Hydrochloride IP 100mcg
 Excipients q.s.



Indication : Clonidine treats high blood pressure by decreasing your heart rate and relaxing the blood vessels so that blood can flow more easily through the body. Clonidine extended-release tablets may treat ADHD by affecting the part of the brain that controls attention and impulsivity.

10x10 Tablets (Alu-Alu)

M.R.P. 30/-Per 10 Tablets

Clopogril-AS

Composition:

Each uncoated bilayered tablet contains:
 Clopidogrel Bisulphate IP
 eq. to Clopidogrel 75 mg
 Aspirin IP 75 mg
 Colour : Red Oxide of Iron



Clopidogrel is used alone or with aspirin to prevent serious or life-threatening problems with the heart and blood vessels in people who have had a stroke, heart attack, or severe chest pain.

10x10 Tablets (Aluminium)

M.R.P. 38.95/-Per 10 Tablets

Clopidogril-AS Forte

Composition:

Each uncoated bilayered tablet contains:
 Clopidogrel Bisulphate IP 75 mg
 eq. to Clopidogrel
 Aspirin IP 150 mg
 Colour : Red Oxide of Iron



Indication : Clopidogrel is used alone or with aspirin to prevent serious or life-threatening problems with the heart and blood vessels in people who have had a stroke, heart attack, or severe chest pain.

10x10 Tablets (Aluminium)
M.R.P. 39.75/-Per 10 Tablets

Cvdol-3.125

Composition :

Each uncoated tablet contains :
 Carvedilol IP 3.125mg
 Excipients q.s.
 Colour Red oxide of Iron



Indication : Carvedilol is used alone or together with other medicines to treat high blood pressure (hypertension). High blood pressure adds to the workload of the heart and arteries.

4x5x10 Tablets (Blister)
M.R.P. 35/-Per 10 Tablets

Cvdol-12.5

Composition :

Each uncoated tablet contains :
 Carvedilol IP 12.5mg
 Excipients q.s.
 Colour :Quinoline Yellow



Indication : Carvedilol is used alone or together with other medicines to treat high blood pressure (hypertension). High blood pressure adds to the workload of the heart and arteries.

4x5x10 Tablets (Blister)
M.R.P. 65/-Per 10 Tablets

Colchicine-0.5

Composition :

Each uncoated tablet contains :
 Colchicine IP 0.5mg.
 Excipients q.s.



Indication : Colchicine works by decreasing swelling and lessening the build up of uric acid crystals that cause pain in the affected joint(s). This medication is also used to prevent attacks of pain in the abdomen, chest, or joints caused by a certain inherited disease (familial Mediterranean fever)

10x10 Tablets (Alu Alu)
M.R.P. 36.50/-Per 10 Tablets

Cvdol-6.25

Composition :

Each uncoated tablet contains :
 Carvedilol IP 6.25mg
 Excipients q.s.
 Colour :Quinoline Yellow



Indication : Carvedilol is used alone or together with other medicines to treat high blood pressure (hypertension). High blood pressure adds to the workload of the heart and arteries.

4x5x10 Tablets (Blister)
M.R.P. 45/-Per 10 Tablets

Cvdol-CR 10

Composition :

Each film coated controlled release tablet contains :
 Carvedilol IP 10mg.
 Excipients q.s.
 Colour : Red Oxide of Iron



Indication : Carvedilol is used alone or together with other medicines to treat high blood pressure (hypertension). High blood pressure adds to the workload of the heart and arteries.

10x10 Tablets (Alu Alu)
M.R.P. 99/-Per 10 Tablets

Dapafyzin-10

Composition :

Each film coated tablet contains :
 Dapagliflozin Propanediol Monohydrate
 Eq. to Dapagliflozin 10mg.
 Excipients q.s.
 Colour : Oxide of Iron &



Indication : Dapagliflozin is a medicine is mainly used to treat type 2 diabetes. Sometimes it can be used together with insulin to treat type 1 diabetes. Dapagliflozin is usually prescribed for type 2 diabetes if you cannot take metformin.

10x10 Tablets (Alu Alu)
 M.R.P. 245/-Per 10 Tablets

Dapafyzin-5

Composition :

Each film coated tablet contains :
 Dapagliflozin Propanediol Monohydrate
 Eq. to Dapagliflozin 5mg.
 Excipients q.s.
 Colour : Oxide of Iron & Titanium Dioxide IP



Indication : Dapagliflozin is a medicine is mainly used to treat type 2 diabetes. Sometimes it can be used together with insulin to treat type 1 diabetes. Dapagliflozin is usually prescribed for type 2 diabetes if you cannot take metformin.

10x10 Tablets (Alu Alu)
 M.R.P. 170/-Per 10 Tablets

Dapafyzin-M10/1000

Composition :

Each film coated bilayered tablet contains :
 Dapagliflozin propanediol monohydrate
 eq. to Dapagliflozin 10 mg.
 Metformin Hydrochloride IP 1000 mg. (In extended release form)
 Excipients q.s.
 Colours : Red Oxide of Iron & Titanium Dioxide IP



Indication : Dapagliflozin and metformin combination is used together with proper diet and exercise to treat type 2 diabetes. It is also used to lower the risk of hospitalization for heart failure in patients with type 2 diabetes and cardiovascular (heart or blood vessel) disease or multiple cardiovascular risk factors.

10x7 Tablets (Alu Alu)
 M.R.P. 198/-Per 7 Tablets

Dapafyzin M5/1000

Composition :

Each film coated bilayered tablet contains :
 Dapagliflozin Propanediol monohydrate
 eq. to Dapagliflozin 5gm.
 Metformin Hydrochloride IP 1000mg. (In extended release form)
 Excipients q.s.
 Colour : Red Oxide of Iron



Indication : Dapagliflozin and metformin combination is used together with proper diet and exercise to treat type 2 diabetes. It is also used to lower the risk of hospitalization for heart failure in patients with type 2 diabetes and cardiovascular (heart or blood vessel) disease or multiple cardiovascular risk factors.

10x7 Tablets (Alu Alu)
 M.R.P. 147/-Per 7 Tablets

Dapafyzin-T 5/20

Composition :

Each film coated tablet contains :
 Dapagliflozin Propanediol Monohydrate
 Eq. to Dapagliflozin 5mg.
 Tenelegliptin Hydrobromide Hydrate IP
 eq. to tenelegliptin 20mg
 Excipients q.s.
 Colour : Titanium Dioxide IP & ferric oxide Yellow USP NF



Indication : Dapagliflozin+Tenelegliptin is used in the treatment of Type 2 diabetes mellitus.

10x10 Tablets (Alu Alu)
 M.R.P. 185/-Per 10 Tablets

Dapafyzin-M10/500

Composition :

Each film coated bilayered tablet contains :
 Dapagliflozin propanediol monohydrate
 eq. to Dapagliflozin 10 mg.
 Metformin Hydrochloride IP 500 mg. (In extended release form)
 Excipients q.s.
 Colours : Red Oxide of Iron & Titanium Dioxide IP



Indication : Dapagliflozin and metformin combination is used together with proper diet and exercise to treat type 2 diabetes. It is also used to lower the risk of hospitalization for heart failure in patients with type 2 diabetes and cardiovascular (heart or blood vessel) disease or multiple cardiovascular risk factors.

10x7 Tablets (Alu Alu)
 M.R.P. 168/-Per 7 Tablets

Defneed-1[®]

Composition :

Each uncoated tablet contains:
Deflazacort 1 mg.
Excipients q.s.

Indication : Deflazacort is used to treat Duchenne muscular dystrophy (DMD; a progressive disease in which the muscles do not function properly) in adults and children 2 years of age and older. Deflazacort is in a class of medications called corticosteroids.



10x10 Tablets (Alu Alu)
M.R.P. 80/-Per 10 Tablets

Defneed-6[®]

Composition :

Each uncoated tablet contains:
Deflazacort 6 mg.
Excipients q.s.

Indication : Deflazacort is used to treat Duchenne muscular dystrophy (DMD; a progressive disease in which the muscles do not function properly) in adults and children 2 years of age and older. Deflazacort is in a class of medications called corticosteroids.



10x10 Tablets (Alu Alu)
M.R.P. 190/-Per 10 Tablets

Defneed-30[®]

Composition :

Each uncoated tablet contains:
Deflazacort 30 mg.
Excipients q.s.

Indication : Deflazacort is used to treat Duchenne muscular dystrophy (DMD; a progressive disease in which the muscles do not function properly) in adults and children 2 years of age and older. Deflazacort is in a class of medications called corticosteroids.



10x10 Tablets (Alu Alu)
M.R.P. 350/-Per 10 Tablets

Derifac 7.5

Composition :

Each film coated prolonged release tablet contains:
Derifenacin Hydrochloride IP Eq. to Derifenacin 7.5mg.
Excipients q.s.

Indication : Darifenacin is used to treat an overactive bladder. By relaxing the muscles in the bladder, darifenacin improves your ability to control your urination. It helps to reduce leaking of urine, feelings of needing to urinate right away, and frequent trips to the bathroom.



10x1x10 Tablets (Alu Alu)
M.R.P. 262/-Per 10 Tablets

Dexatore-4

Composition :

Each uncoated tablet contains :
Dexamethasone IP 4mg.
Excipients q.s.

Indication : It relieves inflammation (swelling, heat, redness, and pain) and is used to treat certain forms of arthritis; skin, blood, kidney, eye, thyroid, and intestinal disorders (e.g., colitis); severe allergies; and asthma. Dexamethasone is also used to treat certain types of cancer.



10x10 Tablets (Blister)
M.R.P. 51.60/-Per 10 Tablets

Dexatore-8

Composition :

Each uncoated tablet contains :
Dexamethasone IP 8mg.
Excipients q.s.

Indication : It relieves inflammation (swelling, heat, redness, and pain) and is used to treat certain forms of arthritis; skin, blood, kidney, eye, thyroid, and intestinal disorders (e.g., colitis); severe allergies; and asthma. Dexamethasone is also used to treat certain types of cancer.



10x10 Tablets (Blister)
M.R.P. 95/-Per 10 Tablets



Desfaxin-50

Each uncoated sustained release tablet contains :
Desvenlafaxine Succinate
Eq. to Desvenlafaxine 50mg.
Excipients q.s.



Desvenlafaxine is used to treat depression. It may improve your mood, feelings of well-being, and energy level. Desvenlafaxine is known as a serotonin-norepinephrine reuptake inhibitor (SNRI). It works by helping to restore the balance of certain natural substances (serotonin and norepinephrine) in the brain.

10x10 Tablets (Blister)
M.R.P. 119/-Per 10 Tablets

Digoxin-0.25

Composition :
Each uncoated tablet contains:
Digoxin IP 0.25mg.
Excipients q.s.
Colour: Titanium Dioxide IP



Indication : Digoxin is derived from the leaves of a digitalis plant. Digoxin helps make the heart beat stronger and with a more regular rhythm. Digoxin is also used to treat atrial fibrillation, a heart rhythm disorder of the atria (the upper chambers of the heart that allow blood to flow into the heart).

10x10 Tablets (Blister)
M.R.P. 14.80/-Per 10 Tablets

Ditalok-30SR

Composition :
Each film coated sustained release Tablet contains :
Diltiazem Hydrochloride IP 30mg.
Excipients q.s.
Colour : Indigo Carmine & Titanium Dioxide IP



Indication : Diltiazem is used to treat high blood pressure and to control angina (chest pain). Diltiazem is in a class of medications called calcium-channel blockers. It works by relaxing the blood vessels so the heart does not have to pump as hard. It also increases the supply of blood and oxygen to the heart.

10x10 Tablets (Blister)
M.R.P. 26.85/-Per 10 Tablets

Ditalok-60

Composition :
Each film coated sustained release Tablet contains :
Diltiazem Hydrochloride IP 60mg.
Excipients q.s.
Colour : Indigo Carmine & Titanium Dioxide IP



Indication : Diltiazem is used to treat high blood pressure and to control angina (chest pain). Diltiazem is in a class of medications called calcium-channel blockers. It works by relaxing the blood vessels so the heart does not have to pump as hard. It also increases the supply of blood and oxygen to the heart.

10x10 Tablets (Blister)
M.R.P. 54.85/-Per 10 Tablets

Ditalok-90

Composition :
Each film coated sustained release Tablet contains :
Diltiazem Hydrochloride IP 90mg.
Excipients q.s.



Indication : Diltiazem is used to treat high blood pressure and to control angina (chest pain). Diltiazem is in a class of medications called calcium-channel blockers. It works by relaxing the blood vessels so the heart does not have to pump as hard. It also increases the supply of blood and oxygen to the heart.

10x10 Tablets (Blister)
M.R.P. 115.90/-Per 10 Tablets

Divaplay-250

Composition :
Each film coated extended release tablet contains :
Divalproex Sodium IP
Eq. to Valporic Acid 250mg.
Excipients q.s.
Colour : Sunset Yellow & Titanium Dioxide IP



Indication : This medication is used to treat seizure disorders, mental/mood conditions (such as manic phase of bipolar disorder), and to prevent migraine headaches. It works by restoring the balance of certain natural substances (neurotransmitters) in the brain.

10x10 Tablets (Alu Alu)
M.R.P. 82/-Per 10 Tablets

Glicaday-M 80

Composition :

Each uncoated tablet contains :
 Gliclazide IP 80 mg.
 Metformin Hydrochloride IP 500mg.
 Excipients q.s.
 Colour: Titanium Dioxide IP

Indication : Gliclazide and metformin is a combination of two oral diabetes medicines that help control blood sugar levels. Gliclazide and metformin is used together with diet and exercise to improve blood sugar control in adults with type 2 diabetes



10x10 Tablets (Blister)
 M.R.P. 64.60/-Per 10 Tablets

Glicaday-60 MR

Composition :

Each uncoated modified release tablet contains
 Gliclazide IP 60mg.
 Excipients q.s.

Indication : Gliclazide is one of a group of medicines called sulfonylureas that are used to lower the blood sugar level. Gliclazide tablets are used to treat maturity onset diabetes in patients that is not controlled by diet alone. They are not for use in children or in patients with diabetes requiring insulin.



10x10 Tablets (Blister)
 M.R.P. 143/-Per 10 Tablets

Glimkar-1

Composition :

Each uncoated tablet contains :
 Glimperide IP 1mg
 Excipients q.s.

Indication : Glimperide is a medicine used to treat type 2 diabetes. Type 2 diabetes is a condition where the body doesn't make enough insulin, or the insulin that it makes doesn't work properly. This can cause high blood sugar levels (hyperglycaemia).



4x5x10 Tablets (Blister)
 M.R.P. 36.60/-Per 10 Tablets

Glicaday-40

Composition :

Each uncoated tablet contains
 Gliclazide IP 40mg.
 Excipients q.s.

Indication : Gliclazide is one of a group of medicines called sulfonylureas that are used to lower the blood sugar level. Gliclazide tablets are used to treat maturity onset diabetes in patients that is not controlled by diet alone. They are not for use in children or in patients with diabetes requiring insulin.



10x10 Tablets (Blister)
 M.R.P. 95/-Per 10 Tablets

Glicaday-80

Composition :

Each uncoated tablet contains
 Gliclazide IP 80mg.
 Excipients q.s.

Indication: Gliclazide is one of a group of medicines called sulfonylureas that are used to lower the blood sugar level. Gliclazide tablets are used to treat maturity onset diabetes in patients that is not controlled by diet alone. They are not for use in children or in patients with diabetes requiring insulin.



10x10 Tablets (Blister)
 M.R.P. 82.20/-Per 10 Tablets

Glimkar-2

Composition :

Each uncoated tablet contains :
 Glimperide IP 2mg
 Excipients q.s.

Indication : Glimperide is a medicine used to treat type 2 diabetes. Type 2 diabetes is a condition where the body doesn't make enough insulin, or the insulin that it makes doesn't work properly. This can cause high blood sugar levels (hyperglycaemia).



4x5x10 Tablets (Blister)
 M.R.P. 57.20/-Per 10 Tablets

Dulixtyn-20 GR

Composition :

Each enteric coated tablet contains :
 Duloxetine Hydrochloride IP 20mg.
 Eq. to Duloxetine 20mg.
 Excipients q.s.

Indication : Duloxetine is an antidepressant medicine. It's used to treat depression and anxiety. It's also used to treat nerve pain, such as fibromyalgia, and can be used to treat stress urinary incontinence in women. Duloxetine comes as capsules and is only available on prescription.



10x10 Tablets (Alu Alu)

M.R.P. 79/-Per 10 Tablets

Dydrocare-10

Composition

Each film-coated tablet contains :
 Dydrogesterone IP 10mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Dydrogesterone helps to regulate the healthy growth and normal shedding of the uterus lining. Therefore, it may be useful in the treatment of menstrual disorders such as absent, irregular or painful menstrual periods, infertility, premenstrual syndrome and endometriosis.



10x1x10 Tablets (Blister)

M.R.P. 827/-Per 10 Tablets

Enacare-5

Composition :

Each uncoated tablet contains :
 Enalapril Maleate IP 5 mg.
 Excipients q.s.

Indication : Enalapril is a medicine used to reduce high blood pressure and to prevent or treat heart failure. If you have high blood pressure, taking enalapril will help prevent a future heart attack or stroke. This medicine is only available on prescription. It comes as tablets.



20x10 Tablets (Blister)

M.R.P. 41.44/-Per 10 Tablets

DutaStay®-0.5

Composition :

Each film coated tablet contains :
 Dutasteride IP 0.5mg
 Excipients q.s.
 Colour: Titanium Dioxide IP

Indication : Dutasteride is used alone or with another medication (tamsulosin [Flomax]) to treat benign prostatic hyperplasia (BPH; enlargement of the prostate gland). Dutasteride is used to treat symptoms of BPH and may reduce the chance of developing acute urinary retention (sudden inability to urinate).



10x10 Tablets (Alu Alu)

M.R.P. 140/-Per 10 Tablets

Enacare 2.5

Composition :

Each uncoated tablet contains :
 Enalapril Maleate IP 2.5 mg.
 Excipients q.s.

Indication : Enalapril is a medicine used to reduce high blood pressure and to prevent or treat heart failure. If you have high blood pressure, taking enalapril will help prevent a future heart attack or stroke. This medicine is only available on prescription. It comes as tablets.



20x10 Tablets (Blister)

M.R.P. 23.90/-Per 10 Tablets

Eplerout-25

Composition :

Each film coated tablet contains :
 Eplerenone IP 25 mg
 Excipients q.s.
 Colour : Yellow oxide of iron

Indication : It's used to treat heart failure and reduce the risk of you having other heart problems or a stroke. It also helps to stop heart failure getting worse. It can sometimes be used to treat a condition called hyperaldosteronism. This is when your body makes too much aldosterone, a hormone that controls your blood pressure.



10x1x10 Tablets (Alu Alu)

M.R.P. 436.08/-Per 10 Tablets

Esamlodose-2.5

Composition :

Each uncoated tablet contains :
S(-)Amlodipine Besylate IP
Equivalent to S(-)Amlodipine 2.5mg.
Excipients q.s.

Indication : S Amlodipine Tablet belongs to a class of medicines known as calcium channel blockers. It is used to treat high blood pressure (hypertension) and prevent angina (heart-related chest pain). It lowers blood pressure and reduces the heart's workload, which helps prevent heart attacks and strokes.



20x10 Tablets (Blister)

M.R.P. 51/-Per 10 Tablets

Efaearn-600

Composition :

Each uncoated tablet contains:
Efavirenz IP 600mg.
Excipients q.s.

Indication : Efavirenz is used along with other medications to treat human immunodeficiency virus (HIV) infection. Efavirenz is in a class of medications called non-nucleoside reverse transcriptase inhibitors (NNRTIs). It works by decreasing the amount of HIV in the blood.



1x30 Tablets

M.R.P. 2371.8/-Per 30 Tablets

Etobone-90

Composition :

Each film coated tablet contains :
Etoricoxib IP 90mg.
Excipients q.s.
Colour : Brillinat Blue Lake & Tartrazine

Indication : Etoricoxib eases pain and swelling (inflammation) in conditions like osteoarthritis, rheumatoid arthritis and ankylosing spondylitis, and it may also be used for short periods of time in gout



10x10 Tablets (Blister)


M.R.P. 120/-Per 10 Tablets

Esamlodose-5

Composition :

Each uncoated tablet contains :
S(-)Amlodipine Besylate IP
Equivalent to S(-)Amlodipine 5mg.
Excipients q.s.

Indication : S Amlodipine Tablet belongs to a class of medicines known as calcium channel blockers. It is used to treat high blood pressure (hypertension) and prevent angina (heart-related chest pain). It lowers blood pressure and reduces the heart's workload, which helps prevent heart attacks and strokes.



4x5x10 Tablets (Blister)

M.R.P. 52/-Per 10 Tablets

Etobone-60

Composition :

Each film coated tablet contains :
Etoricoxib IP 60mg.
Excipients q.s.
Colour : Sunset Yellow & brilliant blue

Indication : Etoricoxib eases pain and swelling (inflammation) in conditions like osteoarthritis, rheumatoid arthritis and ankylosing spondylitis, and it may also be used for short periods of time in gout



10x10 Tablets (Blister)


M.R.P. 90/-Per 10 Tablets

Etobone-120

Composition :

Each film coated tablet contains :
Etoricoxib IP 120mg.
Excipients q.s.
Colour : Brillinat Blue Lake & Tartrazine

Indication : Etoricoxib eases pain and swelling (inflammation) in conditions like osteoarthritis, rheumatoid arthritis and ankylosing spondylitis, and it may also be used for short periods of time in gout



10x10 Tablets (Blister)

M.R.P. 190/-Per 10 Tablets




Etobone-P

Composition :

Each film coated tablet contains :
Etoricoxib IP 60mg.
Paracetamol IP 325mg
Excipients q.s.
Colour : Quinoline Yellow Lake,
Brillinat Blue Lake, Oxide Black &
Titanium Dioxide IP

Etoricoxib+Paracetamol is a pain relieving medicine. It is used to reduce pain and inflammation in conditions like rheumatoid arthritis, ankylosing spondylitis, and osteoarthritis. It may also be used to relieve muscle pain, back pain, toothache, or pain in the ear and throat.



10x10 Tablets (Blister)
M.R.P. 170/-Per 10 Tablets

Ezzitry-10

Composition :

Each uncoated tablet contains:
Ezetimibe IP 10 mg.
Excipients q.s.

Indication : Ezetimibe is used together with lifestyle changes (diet, weight-loss, exercise) to reduce the amount of cholesterol (a fat-like substance) and other fatty substances in the blood. It may be used alone or in combination with an HMG-CoA reductase inhibitor (statin).



10x10 Tablets (Alu Alu)
M.R.P. 122/-Per 10 Tablets

Febaus-80

Composition :

Each film coated tablet contains :
Febuxostat IP 80mg
Excipients q.s.
Colour : Titanium Dioxide IP

Indication : Febuxostat is in a class of medications called xanthine oxidase inhibitors. It works by decreasing the amount of uric acid that is made in the body. Febuxostat is used to prevent gout attacks but not to treat them once they occur.




10x10 Tablets (Alu Alu)
M.R.P. 185/-Per 10 Tablets

Etobone-TH

Composition :

Each film coated tablet contains :
Etoricoxib IP 60mg.
Thiocolchicoside IP 4mg
Excipients q.s.
Colour : Tartrazine

Indication : Etoricoxib+Thiocolchicoside is used in the treatment of muscular pain. Etoricoxib + Thiocolchicoside is a combination of two medicines: Etoricoxib and Thiocolchicoside, which relieves pain and relaxes the muscles.



10x10 Tablets (Alu Alu)
M.R.P. 185/-Per 10 Tablets

Febaus-40

Composition :

Each film coated tablet contains :
Febuxostat IP 40mg
Excipients q.s.
Colour : Titanium Dioxide IP

Indication : Febuxostat is in a class of medications called xanthine oxidase inhibitors. It works by decreasing the amount of uric acid that is made in the body. Febuxostat is used to prevent gout attacks but not to treat them once they occur.



10x10 Tablets (Alu Alu)
M.R.P. 145/-Per 10 Tablets

Fenoday-145

Composition :

Each film coated tablet contains:
Fenofibrate IP 145mg.
Excipients q.s.
Colour : Red oxide of Iron &
Titanium Dioxide IP

Indication : Fenofibrate is used along with a proper diet to help lower "bad" cholesterol and fats (such as LDL, triglycerides) and raise "good" cholesterol (HDL) in the blood. It works by increasing the natural substance (enzyme) that breaks down fats in the blood.



10x10 Tablets (Blister)
M.R.P. 135/-Per 10 Tablets

Fenoday-160

Composition :

Each film coated tablet contains:
 Fenofibrate IP 160mg.
 Excipients q.s.
 Colour : Red oxide of Iron & Titanium Dioxide IP

Indication : Fenofibrate is used along with a proper diet to help lower "bad" cholesterol and fats (such as LDL, triglycerides) and raise "good" cholesterol (HDL) in the blood. It works by increasing the natural substance (enzyme) that breaks down fats in the blood.



10x10 Tablets (Blister)
 M.R.P. 155/-Per 10 Tablets

Fenpine-300 ER

Composition :

Each film coated extended release tablet contains :
 Faropenem Sodium eq. to Faropenem 300mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Faropenem is approved for use in the treatment of upper and lower respiratory tract infections, ear, nose and throat (ENT) infections, genito-urinary infections, skin and skin structure infections, and gynaecological infections.



10x1x6 Tablets (Alu-Alu)
 M.R.P. 1070/-Per 6 Tablets

Fenpine-200

Composition :

Each film coated tablet contains :
 Faropenem sodium eq. to faropenam 200 mg.
 excipients q.s.

Indication : Faropenem is approved for use in the treatment of upper and lower respiratory tract infections, ear, nose and throat (ENT) infections, genito-urinary infections, skin and skin structure infections, and gynaecological infections.



10x1x6 Tablets (Alu Alu)
 M.R.P. 870/-Per 6 Tablets

Fenoday-200

Composition :

Each film coated tablet contains:
 Fenofibrate IP 200mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Fenofibrate is used along with a proper diet to help lower "bad" cholesterol and fats (such as LDL, triglycerides) and raise "good" cholesterol (HDL) in the blood. It works by increasing the natural substance (enzyme) that breaks down fats in the blood.



10x10 Tablets (Blister)
 M.R.P. 185/-Per 10 Tablets

Fenpine-CV

Composition

Each film-coated tablet contains :
 Faropenem Sodium Hydrate JP eq. to Faropenem 200mg.
 Potassium Clavulanate Diluted IP eq. to clavulanic Acid 125mg.
 Excipients q.s.
 Colours : Red Oxide of Iron & Titanium Dioxide IP

Indication : Faropenem is approved for use in the treatment of upper and lower respiratory tract infections, ear, nose and throat (ENT) infections, genito-urinary infections, skin and skin structure infections, and ynaecological infections.



10x1x6 Tablets (Alu Alu)
 M.R.P. 1085/-Per 6 Tablets

Fexofine-120

Composition :

Each film coated tablet contains
 Fexofenadine HCl IP 120 mg.
 Excipients q.s.
 Colours : Yellow oxide of Iron, Red Oxide of Iron & Titanium Dioxide IP

Indication : Fexofenadine is an antihistamine used to relieve allergy symptoms such as watery eyes, runny nose, itching eyes/nose, sneezing, hives, and body itching. It works by blocking a certain natural substance (histamine) that your body makes during an allergic



10x10 Tablets (Blister)
 M.R.P. 190/-Per 10 Tablets

Fexofine-180

Composition :

Each film coated tablet contains
 Fexofenadine HCl IP 180 mg.
 Excipients q.s.
 Colours : Yellow oxide of Iron, Red
 Oxide of Iron & Titanium Dioxide IP

Indication : Fexofenadine is an antihistamine used to relieve allergy symptoms such as watery eyes, runny nose, itching eyes/nose, sneezing, hives, and body itching. It works by blocking a certain natural substance (histamine) that your body makes during an allergic



10x10 Tablets (Blister)

M.R.P. 225/-Per 10 Tablets

Finasteride-1

Composition :

Each film coated tablet contains :
 Finasteride IP 1mg
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Finasteride is used to treat men with an enlarged prostate (benign prostate enlargement). It can help ease your symptoms if: it's difficult to start peeing, you need to pee urgently or frequently more often.



10x10 Tablets (Alu-Alu)

M.R.P. 121/-Per 10 Tablets

Fluliq-M

Composition :

Each film coated tablet contains :
 Flupentixol Hydrochloride BP eq. to Flupentixol 0.5 mg.
 Melitracen Hydrochloride eq. to Melitracen 10 mg.
 Excipients q.s.
 Colours : Erythrosine & Indigo Carmine.

Flupentixol is a thiazolyl (thioanthracene) antipsychotic, and melitracen is a tricyclic antidepressant. Low dose Flupentixol (0.5mg-3mg) has antidepressant and anti-anxiety effects, while melitracen has antidepressant effect. The mixture of the two components is used to treat mild to moderate mental disorders.



10x10 Tablets (Blister)

M.R.P. 75/-Per 10 Tablets

Fexofine-M

Composition :

Each film coated tablet contains :
 Fexofenadine HCl IP 120mg,
 Montelukast Sodium IP 10 mg.
 Eq. to Montelukast 10 mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Montelukast+ Fexofenadine Is Used To Relieve Allergy Symptoms Such As Watery Eyes, Runny Nose, Itching Eyes/nose, And Sneezing.



10x1x10 Tablets (Alu Alu)

M.R.P. 193/-Per 10 Tablets

Finasteride-5

Composition :

Each film coated tablet contains:
 Finasteride IP 5mg.
 Excipients q.s.

Indication : Finasteride is used to treat men with an enlarged prostate (benign prostate enlargement). It can help ease your symptoms if: it's difficult to start peeing, you need to pee urgently or frequently more often.



10x10 Tablets (Blister)

M.R.P. 182/-Per 10 Tablets

Fluritime-5

Composition :

Each uncoated tablet contains:
 Flunarizine Dihydrochloride BP Eq. to Flunarizine 5mg.
 Excipients q.s.

Indication : Flunarizine is used to prevent migraine headaches with or without aura (warning signs that occur before the headache begins). This medication should not be used for treatment of acute migraine headaches (headaches that have already started)



20x10 Tablets (Blister)

M.R.P. 30.20/-Per 10 Tablets

Fluritime-10

Composition :

Each uncoated tablet contains:
 Flunarizine Dihydrochloride BP 10mg.
 Eq.to Flunarizine
 Excipients q.s.

Indication : Flunarizine is used to prevent migraine headaches with or without aura (warning signs that occur before the headache begins). This edication should not be used for treatment of acute migraine headaches (headaches that have already started)



10x10 Tablets (Blister)
 M.R.P. 48.16/-Per 10 Tablets

Fluppy-P

Composition :

Each uncoated bilayered tablet contains :
 Flupirtine Maleate 100mg.
 Paracetamol IP 325mg.
 Excipients q.s.
 Colour : Sunset Yellow

Indication : Flupirtine Maleate+paracetamol May Be Used To Treat Acute And Chronic Pain, Fever, Headache, Joint Pains, Dysmenorhea (period Pain), Postoperative Pain, Toothache After Dental Extraction, And Myalgia (muscle Pain). Pain Is A Term Used To Describe An Unpleasant Feeling or discomfort.



10x10 Tablets
 M.R.P. 158/-Per 10 Tablets

Fluvomax-50

Composition :

Each film coated tablet contains :
 Fluvoxamine Maleate IP 50mg.
 Excipients q.s.
 Colour : Yellow oxide of Iron & Titanium Dioxide IP

Indication : Fluvoxamine is used to treat obsessive-compulsive disorder (bothersome thoughts that won't go away and the need to perform certain actions over and over) and social anxiety disorder (extreme fear of interacting with others or performing in front of others that interferes with normal life).



10x10 Tablets (Aluminium)
 M.R.P. 198/-Per 10 Tablets

Folaus

Composition :

Each film coated tablet contains:
 Methylcobalamin IP 1500 mcg.
 L-5 Methyltetrahydrofolate Calcium eq.to L-Methyl Folate 1 mg.
 Pyridoxal-5 Phosphate 0.5 mg.
 Excipients q.s.
 Colour : Ponceau 4r Lake & Titanium Dioxide IP

L-Methylfolate+Methylcobalamin+Pyridoxal 5 Phosphate Is Used To Treat Nutritional Deficiencies. It Contains L-methylfolate, Pyridoxal-5-phosphate, And Methylcobalamin. L-methylfolate Helps The Formation Of Red Blood Cells And Prevents Dna Changes That May Lead To Cancer. It Prevents And Treats Folate Deficiency.



10x10 Tablets (Alu Alu)
 M.R.P. 150/-Per 10 Tablets

Folifresh[®]-5

Composition :

Each uncoated tablet contains :
 Folic Acid IP 5mg.
 Excipients q.s.

Indication : Folic acid is used to: treat or prevent folate deficiency anaemia. help your unborn baby's brain, skull and spinal cord develop properly to avoid development problems (called neural tube defects) such as spina bifida.



10x1x30 Tablets (Blister)
 M.R.P. 50.37/-Per 30 Tablets



Furosemide-40

Composition :

Each uncoated tablet contains:
 Furosemide IP 40mg.
 Excipients q.s.

Indication : Furosemide is a loop diuretic (water pill) that prevents your body from absorbing too much salt. This allows the salt to instead be passed in your urine. Furosemide is used to treat fluid retention (edema) in people with congestive heart failure, liver disease, or a kidney disorder such as nephrotic syndrome.



20x10 Tablets (Blister)

M.R.P. 10/-Per 10 Tablets

Gebcare-100

Composition :

Each film coated tablet contains :
 Gabapentin IP 100mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Gabapentin is used with other medications to prevent and control seizures. It is also used to relieve nerve pain following shingles (a painful rash due to herpes zoster infection) in adults. Gabapentin is known as an anticonvulsant or antiepileptic drug.



10x15 Tablets (Blister)

M.R.P. 132/-Per 15 Tablets

Gefidoz-250

Composition :

Each film coated tablet contains:
 Gefitinib IP 250mg.
 Excipients q.s.
 Colours : Red oxide of Iron
 Yellow oxide of Iron &
 Titanium Dioxide IP

Indication : Gefitinib is used to treat metastatic (cancer that has already spread) non-small cell lung cancer in patients who have certain types of abnormal epidermal growth factor (EGFR) genes, and did not receive cancer medicines in the past. Your doctor will perform a test before you take this medicine.



3x10 Tablets (Blister)

M.R.P. 1965/-Per 10 Tablets

Gebcare-NT 100/10

Composition :

Each film coated tablet contains :
 Gabapentin IP 100mg.
 Nortriptyline Hydrochloride IP
 eq. to Nortriptyline 10mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Gabapentin+nortriptyline Belongs To A Combination Medicine Of Anticonvulsant (gabapentin) And Antidepressant (nortriptyline) Used For The Management Of Neuropathic Pain. It Is A Chronic Progressive Nerve Disease That Causes Nerve Pain.



10x10 Tablets (Alu-Alu)

M.R.P. 155/-Per 10 Tablets

GEB CARE-NT

Composition :

Each film coated tablet contains :
 Gabapentin IP 400mg.
 Nortriptyline HCl IP 10 mg.
 Excipients q.s.
 Colour : Ponceau 4R

Indication : Gabapentin relieves pain arising due to nerve damage, compression in nerves. Nortriptyline. These two medicines work effectively in reducing neuropathic pain (pain due to nerve damage). You should take this medicine in a dose and duration prescribed by your doctor.



10x10 Tablets (Alu Alu)

M.R.P. 349/-Per 10 Tablets

Glicaday-30 MR

Composition :

Each uncoated modified release tablet contains
 Gliclazide IP 30mg.
 Excipients q.s.

Indication : Gliclazide is one of a group of medicines called sulfonylureas that are used to lower the blood sugar level. Gliclazide tablets are used to treat maturity onset diabetes in patients that is not controlled by diet alone. They are not for use in children or in patients



10x10 Tablets (Blister)

M.R.P. 83/-Per 10 Tablets

Glicaday-M 80

Composition :

Each uncoated tablet contains :
 Gliclazide IP 80 mg.
 Metformin Hydrochloride IP 500mg.
 Excipients q.s.
 Colour: Titanium Dioxide IP

Indication : Gliclazide and metformin is a combination of two oral diabetes medicines that help control blood sugar levels. Gliclazide and metformin is used together with diet and exercise to improve blood sugar control in adults with type 2 diabetes



10x10 Tablets (Blister)
 M.R.P. 64.60/-Per 10 Tablets

Glicaday-60 MR

Composition :

Each uncoated modified release tablet contains
 Gliclazide IP 60mg.
 Excipients q.s.

Indication : Gliclazide is one of a group of medicines called sulfonylureas that are used to lower the blood sugar level. Gliclazide tablets are used to treat maturity onset diabetes in patients that is not controlled by diet alone. They are not for use in children or in patients with diabetes requiring insulin.



10x10 Tablets (Blister)
 M.R.P. 143/-Per 10 Tablets

Glimkar-1

Composition :

Each uncoated tablet contains :
 Glimperide IP 1mg.
 Excipients q.s.

Indication : Glimperide is a medicine used to treat type 2 diabetes. Type 2 diabetes is a condition where the body doesn't make enough insulin, or the insulin that it makes doesn't work properly. This can cause high blood sugar levels (hyperglycaemia).



4x5x10 Tablets (Blister)
 M.R.P. 36.60/-Per 10 Tablets

Glicaday-40

Composition :

Each uncoated tablet contains
 Gliclazide IP 40mg.
 Excipients q.s.

Indication : Gliclazide is one of a group of medicines called sulfonylureas that are used to lower the blood sugar level. Gliclazide tablets are used to treat maturity onset diabetes in patients that is not controlled by diet alone. They are not for use in children or in patients with diabetes requiring insulin.



10x10 Tablets (Blister)
 M.R.P. 95/-Per 10 Tablets

Glicaday-80

Composition :

Each uncoated tablet contains
 Gliclazide IP 80mg.
 Excipients q.s.

Indication: Gliclazide is one of a group of medicines called sulfonylureas that are used to lower the blood sugar level. Gliclazide tablets are used to treat maturity onset diabetes in patients that is not controlled by diet alone. They are not for use in children or in patients with diabetes requiring insulin.



10x10 Tablets (Blister)
 M.R.P. 82.20/-Per 10 Tablets

Glimkar-2

Composition :

Each uncoated tablet contains :
 Glimperide IP 2mg.
 Excipients q.s.

Indication : Glimperide is a medicine used to treat type 2 diabetes. Type 2 diabetes is a condition where the body doesn't make enough insulin, or the insulin that it makes doesn't work properly. This can cause high blood sugar levels (hyperglycaemia).



4x5x10 Tablets (Blister)
 M.R.P. 57.02/-Per 10 Tablets



Glimkar-3

Composition :

Each uncoated tablet contains :
Glimepiride IP 3mg
Excipients q.s.



Indication : Glimepiride is a medicine used to treat type 2 diabetes. Type 2 diabetes is a condition where the body doesn't make enough insulin, or the insulin that it makes doesn't work properly. This can cause high blood sugar levels (hyperglycaemia).

10x10 Tablets (Blister)

M.R.P. 160/-Per 10 Tablets

Glimkar-M 0.5

Composition :

Each uncoated bilayered tablet contains :
Glimepiride IP 0.5mg.
Metformin HCl IP 500mg.
(In sustained Release form)
Excipients q.s.
Colour : Sunset Yellow



Indication : Glimepiride+Metformin is used in the treatment of type 2 diabetes mellitus. Glimepiride + Metformin is a combination of two antidiabetic medicines: Glimepiride and Metformin. Glimepiride is a sulfonylurea which works by increasing the amount of insulin released by the pancreas in order to lower the blood glucose.

10x10 Tablets (Blister)

M.R.P. 60/-Per 10 Tablets

Glimkar-M1 SR

Each uncoated bilayered tablet contains :
Glimepiride IP 1mg
Metformin Hydrochloride IP 500mg
(In Sustained Release Form)
Excipients q.s.
Colour : Sunset Yellow



Glimepiride+Metformin is used in the treatment of type 2 diabetes mellitus. Glimepiride + Metformin is a combination of two antidiabetic medicines: Glimepiride and Metformin. Glimepiride is a sulfonylurea which works by increasing the amount of insulin released by the pancreas in order to lower the blood glucose.

10x10 Tablets (Blister)

M.R.P. 62/-Per 10 Tablets

Glimkar-4

Composition :

Each uncoated tablet contains :
Glimepiride IP 4mg
Excipients q.s.



Indication : Glimepiride is a medicine used to treat type 2 diabetes. Type 2 diabetes is a condition where the body doesn't make enough insulin, or the insulin that it makes doesn't work properly. This can cause high blood sugar levels (hyperglycaemia).

10x10 Tablets (Blister)

M.R.P. 192/-Per 10 Tablets

Glimkar-M1

Composition :

Each uncoated tablet contains :
Metformin Hydrochloride IP 500mg.
Glimepiride IP 1mg.
Excipients q.s.
Colour : Titanium Dioxide IP



Indication : Glimepiride+Metformin is used in the treatment of type 2 diabetes mellitus. Glimepiride + Metformin is a combination of two antidiabetic medicines: Glimepiride and Metformin. Glimepiride is a sulfonylurea which works by increasing the amount of insulin released by the pancreas in order to lower the blood glucose.


10x10 Tablets (Blister)

M.R.P. 40.30/-Per 10 Tablets

Glimkar-M2

Composition

Each uncoated tablet contains :
Metformin Hydrochloride IP 1000mg.
Glimepiride IP 2mg.
Excipients q.s.
Colour : Titanium Dioxide IP



Indication : Glimepiride+Metformin is used for the treatment of type 2 diabetes mellitus, when diet, exercise and glimepiride or metformin alone is unable to control your blood sugar levels adequately.

10x10 Tablets (Blister)

M.R.P. 47.25/-Per 10 Tablets

Glimkar-M2 SR

Composition :

Each uncoated bilayered tablet contains :


Glimepiride IP	2mg
Metformin Hydrochloride IP (In Sustained Release Form)	500mg
Excipients	q.s.

Color : Quinoline Yellow

Indication : Glimepiride+Metformin is used in the treatment of type 2 diabetes mellitus. Glimepiride + Metformin is a combination of two antidiabetic medicines: Glimepiride and Metformin. Glimepiride is a sulfonylurea which works by increasing the amount of insulin released by the pancreas in order to lower the blood glucose.

10x10 Tablets (Blister)

M.R.P. 78/-Per 10 Tablets



Glimkar-M3 SR

Each uncoated bilayered tablet contains :

Glimepiride IP	3mg
Metformin Hydrochloride IP (In Sustained Release Form)	500mg
Excipients	q.s.

Colour : Sunset Yellow

Glimepiride+Metformin is used in the treatment of type 2 diabetes mellitus. Glimepiride + Metformin is a combination of two antidiabetic medicines: Glimepiride and Metformin. Glimepiride is a sulfonylurea which works by increasing the amount of insulin released by the pancreas in order to lower the blood glucose.

10x10 Tablets (Blister)

M.R.P. 89/-Per 10 Tablets

Glimkar-M1 Forte

Composition :

Each uncoated bilayered tablet contains :

Glimepiride IP	1mg.
Metformin Hydrochloride IP (In sustained Release form)	1000mg.
Excipients	q.s.

Colour : Sunset Yellow

Indication : Glimepiride+Metformin is used in the treatment of type 2 diabetes mellitus. Glimepiride + Metformin is a combination of two antidiabetic medicines: Glimepiride and Metformin. Glimepiride is a sulfonylurea which works by increasing the amount of insulin released by the pancreas in order to lower the blood glucose.

10x10 Tablets (Blister)

M.R.P. 72.45/-Per 10 Tablets



Glimkar-M2 Forte

Composition :

Each uncoated bilayered tablet contains :

Glimepiride IP	2mg.
Metformin Hydrochloride IP (In sustained Release form)	1000mg.
Excipients	q.s.

Colour : Quinoline Yellow

Indication : Glimepiride+Metformin is used in the treatment of type 2 diabetes mellitus. Glimepiride + Metformin is a combination of two antidiabetic medicines: Glimepiride and Metformin. Glimepiride is a sulfonylurea which works by increasing the amount of insulin released by the pancreas in order to lower the blood glucose.

10x10 Tablets (Blister)

M.R.P. 98/-Per 10 Tablets

Glimkar-M3-Forte

Composition :

Each uncoated bilayered tablet contains :

Glimepiride IP	3mg.
Metformin Hydrochloride IP (In sustained Release form)	1000mg.
Excipients	q.s.

Colour : Ponceau 4R

Indication : Glimepiride+Metformin is used in the treatment of type 2 diabetes mellitus. Glimepiride + Metformin is a combination of two antidiabetic medicines: Glimepiride and Metformin. Glimepiride is a sulfonylurea which works by increasing the amount of insulin released by the pancreas in order to lower the blood glucose.

10x10 Tablets (Blister)

M.R.P. 93/-Per 10 Tablets



Glimkar-M4-Forte

Composition :

Each uncoated bilayered tablet contains :

Glimepiride IP	4mg.
Metformin Hydrochloride IP (In sustained Release form)	1000mg.
Excipients	q.s.

Colour : Black Oxide of Iron

Indication : Glimepiride+Metformin is used in the treatment of type 2 diabetes mellitus. Glimepiride + Metformin is a combination of two antidiabetic medicines: Glimepiride and Metformin. Glimepiride is a sulfonylurea which works by increasing the amount of insulin released by the pancreas in order to lower the blood glucose.

10x10 Tablets (Blister)

M.R.P. 107/-Per 10 Tablets

Glimkar-PG1

Composition

Each uncoated bilayered tablet contains :

Glimepiride	IP	1 mg.
Pioglitazone Hydrochloride	IP	15 mg.
Eq. to Pioglitazone		15 mg.
Metformin Hydrochloride	IP 500 mg.	
(In extended release form)		
Excipients		q.s.
Colour : Titanium Dioxide IP & Sunset Yellow FCF		

Indication : Glimepiride is a medicine used to treat type 2 diabetes. Type 2 diabetes is a condition where the body doesn't make enough insulin, or the insulin that it makes doesn't work properly. This can cause high blood sugar levels (hyperglycaemia).

10x10 Tablets (Blister)

M.R.P. 53.56/-Per 10 Tablets



Glimkar-Pg2

Composition :

Each uncoated bilayered tablet contains :

Glimepiride	IP	2 mg.
Pioglitazone Hydrochloride	IP	15 mg.
Eq. to Pioglitazone		15 mg.
Metformin Hydrochloride	IP 500 mg.	
(In extended release form)		
Excipients		q.s.
Colour : Titanium Dioxide IP & Sunset Yellow FCF		

Composition : Glimepiride is a medicine used to treat type 2 diabetes. Type 2 diabetes is a condition where the body doesn't make enough insulin, or the insulin that it makes doesn't work properly. This can cause high blood sugar levels (hyperglycaemia).

10x10 Tablets (Blister)

M.R.P. 72.87/-Per 10 Tablets



HC-Cort-10

Composition :

Each uncoated tablet contains

Hydrocortisone IP	10mg.
Excipients	q.s.

Indication : Hydrocortisone is a steroid (corticosteroid) medicine. It works by calming down your body's immune response to reduce pain, itching and swelling (inflammation). It can also be used as hormone replacement for people who do not have enough of the natural stress hormone, cortisol.

10x1x10 Tablets (Blister)

M.R.P. 66/-Per 10 Tablets



HC-Cort-20

Composition :

Each uncoated tablet contains

Hydrocortisone IP	20mg.
Excipients	q.s.

Indication : Hydrocortisone is a steroid (corticosteroid) medicine. It works by calming down your body's immune response to reduce pain, itching and swelling (inflammation). It can also be used as hormone replacement for people who do not have enough of the natural stress hormone, cortisol.

10x1x10 Tablets (Blister)

M.R.P. 166.50/-Per 10 Tablets



Hydrochlorothiazide-12.5

Composition :

Each uncoated tablet contains:

Hydrochlorothiazide	IP 12.5mg.
Excipients	q.s.

Indication : Hydrochlorothiazide is used to treat edema (fluid retention; excess fluid held in body tissues) caused by various medical problems, including heart, kidney, and liver disease and to treat edema caused by using certain medications including estrogen and corticosteroids.

20x10 Tablets (Blister)

M.R.P. 11.07/-Per 10 Tablets



Hydrochlorothiazide-25

Composition :

Each uncoated tablet contains:

Hydrochlorothiazide	IP 25mg.
Excipients	q.s.

Indication : Hydrochlorothiazide is used to treat high blood pressure (hypertension). Hydrochlorothiazide is also used to treat fluid retention (edema) in people with congestive heart failure, cirrhosis of the liver, or kidney disorders, or edema caused by taking steroids or estrogen

20x10 Tablets (Blister)

M.R.P. 19.48/-Per 10 Tablets



Hyrocare[®]-10

Composition :

Each film coated tablet contains :
 Hydroxyzine
 Hydrochloride I.P 10 mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP



Indication : Hydroxyzine is used in adults and children to relieve itching caused by allergic skin reactions. It is also used alone or with other medications in adults and children to relieve anxiety and tension.

20x15 Tablets (Blister)

M.R.P. 35/-Per 15 Tablets

Hyrocare[®]-25

Composition :

Each film coated tablet contains :
 Hydroxyzine
 Hydrochloride I.P 25 mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP



Indication : Hydroxyzine is used in adults and children to relieve itching caused by allergic skin reactions. It is also used alone or with other medications in adults and children to relieve anxiety and tension.

20x15 Tablets (Blister)

M.R.P. 55/-Per 15 Tablets

Halpido-5

Composition :

Each uncoated tablet contains :
 Haloperidol IP 5mg.
 Excipients q.s.



Indication : Haloperidol is a medication that works in the brain to treat schizophrenia. It is also known as a first generation antipsychotic (FGA) or typical antipsychotic. Haloperidol rebalances dopamine to improve thinking, mood, and behavior.

10x10 Tablets (Blister)

M.R.P. 35.60/-Per 10 Tablets

Ibossis-150

Composition :

Each film coated tablet contains :
 Ibandronic Acid Monosodium
 Monohydrate
 Eq. to Ibandronic Acid 150mg
 Excipients q.s.
 Colour : Titanium Dioxide IP



Indication : Ibandronic acid belongs to a group of drugs called bisphosphonates. It can be used to treat: high levels of calcium in the blood caused by cancer that has spread to the bones (secondary bone cancer) bone weakness or pain caused by breast cancer that has spread to the bones.


10x1x1 Tablets (Blister)

M.R.P. 235/-Per 1 Tablets

Isoaus-10

Composition:

Each uncoated tablet contains :
 Diluted Isosorbide Mononitrate IP
 Eq. to Isosorbide Mononitrate 10mg
 Excipients q.s.



Indication : Isosorbide mononitrate is a nitrate that dilates (widens) blood vessels, making it easier for blood to flow through them and easier for the heart to pump. Isosorbide mononitrate is used to prevent angina attacks (chest pain). Isosorbide mononitrate will not treat an angina attack that has already begun.


4x5x10 Tablets (Blister)

M.R.P. 24.10/-Per 10 Tablets

Isoaus-20

Composition :

Each uncoated tablet contains :
 Diluted Isosorbide Mononitrate IP
 Eq. to Isosorbide Mononitrate 20mg
 Excipients q.s.



Indication : Isosorbide mononitrate is a nitrate that dilates (widens) blood vessels, making it easier for blood to flow through them and easier for the heart to pump. Isosorbide mononitrate is used to prevent angina attacks (chest pain). Isosorbide mononitrate will not treat an angina attack that has already begun.

4x5x10 Tablets (Blister)

M.R.P. 34/-Per 10 Tablets

Imaten-400

Composition

Each film coated tablet contains :
Imatinib Mesylate IP
equivalent to imatinib 400mg.
Excipients q.s.
Colour : Ferric oxide Red

Indication : Imatinib is used to treat certain types of leukemia (cancer that begins in the white blood cells) and other cancers and disorders of the blood cells.



3x10 Tablets

M.R.P. 1535/-Per 10 Tablets

Isoniazid-100

Composition :

Each uncoated tablet contains :
Isoniazid IP 100mg.
Excipients q.s.

Indication : Isoniazid is used with other medications to treat active tuberculosis (TB) infections. It is also used alone to prevent active TB infections in people who may be infected with the bacteria (people with positive TB skin test). Isoniazid is an antibiotic and works by stopping the growth



10x10 Tablets (Blister)

M.R.P. 6.80/-Per 10 Tablets

Ivertac-6

Composition :

Each uncoated tablet contains :
Ivermectin USP 6mg.
Excipients q.s.

Indication : Ivermectin is used to treat strongyloidiasis (threadworm; infection with a type of roundworm that enters the body through the skin, moves through the airways and lives in the intestines).



10x10 Tablets (Alu Alu)

M.R.P. 175/-Per 10 Tablets

IBA-Quin

Composition :

Each film coated tablet contains :
Hydroxychloroquine
Sulphate IP 200mg.
Colour : Titanium Dioxide IP

Indication : Hydroxychloroquine is a quinoline medicine used to treat or prevent malaria, a disease caused by parasites that enter the body through the bite of a mosquito. Malaria is common in areas such as Africa, South America, and Southern Asia. This medicine is not effective against all strains of malaria.



10x15 Tablets (Blister)

M.R.P. 106.60/-Per 15 Tablets

Isoniazid-300

Composition :

Each uncoated tablet contains :
Isoniazid IP 300mg.
Excipients q.s.

Indication : Isoniazid is used with other medications to treat active tuberculosis (TB) infections. It is also used alone to prevent active TB infections in people who may be infected with the bacteria (people with positive TB skin test). Isoniazid is an antibiotic and works by stopping the growth



10x10 Tablets (Blister)

M.R.P. 13.54/-Per 10 Tablets

Ivertac-12

Composition :

Each uncoated tablet contains :
Ivermectin USP 12mg.
Excipients q.s.

Indication : Ivermectin is used to treat strongyloidiasis (threadworm; infection with a type of roundworm that enters the body through the skin, moves through the airways and lives in the intestines).



10x10 Tablets (Alu Alu)

M.R.P. 325/-Per 10 Tablets

Ketozox-DT

Composition :

Each uncoated dispersible tablet contains:
Ketorolac Tromethamine I.P. 10 mg.
Excipients q.s.

Indication : Ketorolac is used for the short-term treatment of moderate to severe pain in adults. It is usually used before or after medical procedures or after surgery. Reducing pain helps you recover more comfortably so that you can return to your normal daily activities.



20x15 Tablets (Alu Alu)
M.R.P. 130/-Per 15 Tablets

Lacavir-ET

Composition:

Each film coated tablet contains :
Tenofovir Disoproxil Fumarate IP 300 mg.
Lamivudine IP 300 mg.
Efavirenz IP 600 mg.
Excipients q.s.
Colour : Lake Indigo Caramel

Indication : Efavirenz, lamivudine, and tenofovir combination is used alone to treat human immunodeficiency virus (HIV) infection. HIV is the virus that causes acquired immune deficiency syndrome (AIDS).



1x30 Tablets
M.R.P. 2900/-Per 30 Tablets

Lanscool-15

Composition

Each uncoated orally disintegrating tablet contains :
Lansoprazole IP 15mg.
Excipients q.s.

Indication : It's used for indigestion, heartburn, acid reflux and gastroesophageal-reflux-disease (GORD). Lansoprazole is also taken to prevent and treat stomach ulcers. Sometimes, lansoprazole is taken for a rare condition caused by a tumour in the pancreas or gut called Zollinger-Ellison syndrome.



10x15 Tablets (Alu Alu)
M.R.P. 162/-Per 15 Tablets

Ketoclean®

Composition :

Each uncoated tablet contains
Ketoconazole IP 200mg.
Excipients q.s.

Indication : Ketoconazole is used to treat fungal and yeast infections on your skin, hair, nails, and in your blood.



10x10 Tablets (Alu Alu)
M.R.P. 198/-Per 10 Tablets

Lacavir-150

Composition

Each film coated tablet contains :
Lamivudine IP 150mg.
Excipients q.s.
Colour : Yellow oxide of Iron

Indication : Lamivudine (Epivir-HBV) is used to treat hepatitis B infection. Lamivudine is in a class of medications called nucleoside reverse transcriptase inhibitors (NRTIs). It works by decreasing the amount of HIV and hepatitis B in the blood.



1x30 Tablets
M.R.P. 360/-Per 30 Tablets

Lefoux-500

Composition :

Each film coated tablet contains :
Levofloxacin Hemihydrate IP 500 mg.
eq. to Levofloxacin 500 mg.
Excipients q.s.
Colour : Red Oxide of Iron & Titanium Dioxide I.P.

Indication : Dydrogesterone helps to regulate the healthy growth and normal shedding of the uterus lining. Therefore, it may be useful in the treatment of menstrual disorders such as absent, irregular or painful menstrual periods, infertility, premenstrual syndrome and endometriosis.



20x5 Tablets (Blister)
M.R.P. 49.28/-Per 5 Tablets

Lacosamide-50

Composition :

Each film coated tablet contains :
 Lacosamide 50mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Lacosamide is used alone and in combination with other medications to control certain types of seizures in adults and children 4 years of age and older. Lacosamide is in a class of medications called anticonvulsants. It works by decreasing abnormal electrical activity in the brain.



10x10 Tablets (Blister)

M.R.P. 99/-Per 10 Tablets

Lacosamide 100mg.

Composition :

Each film coated tablet contains :
 Lacosamide 100mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Lacosamide is used alone and in combination with other medications to control certain types of seizures in adults and children 4 years of age and older. Lacosamide is in a class of medications called anticonvulsants. It works by decreasing abnormal electrical activity in the brain.



10x10 Tablets (Blister)

M.R.P. 132/-Per 10 Tablets

Lecarto-500

Composition :

Each film coated tablet contains :
 Levocarnitine USP 500mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Levocarnitine is used to prevent and treat a lack of carnitine. It is used to prevent and treat this condition in patients with kidney disease on dialysis. It is given to people whose body cannot properly use carnitine from their diet. Lack of carnitine can lead to liver, heart, and muscle problems.



10x10 Tablets (Alu Alu)

M.R.P. 293/-Per 10 Tablets

Lefomycs-10

Composition :

Each film coated tablet contains :
 Leflunomide IP 10mg.
 Excipients

Indication : This medication is used to treat rheumatoid arthritis, a condition in which the body's defense system (immune system) fails to recognize the body as itself and attacks the healthy tissues around the joints. Leflunomide helps to reduce the joint damage/pain/swelling and helps you to move better.



10x10 Tablets (Aluminum Strip)

M.R.P. 124.40/-Per 10 Tablets

Lefomycs-20

Composition :

Each film coated tablet contains :
 Leflunomide IP 20mg.
 Excipients

Indication : This medication is used to treat rheumatoid arthritis, a condition in which the body's defense system (immune system) fails to recognize the body as itself and attacks the healthy tissues around the joints. Leflunomide helps to reduce the joint damage/pain/swelling and helps you to move better.



10x10 Tablets (Aluminium Strip)

M.R.P. 214/-Per 10 Tablets

Lerozot-2.5

Composition :

Each film coated tablet contains :
 Letrozole USP 2.5mg.
 Excipients q.s.
 Colour : Yellow Oxide of Iron & Titanium Dioxide IP

Indication : This medication is used to treat certain types of breast cancer (such as hormone-receptor-positive breast cancer) in women after menopause. Letrozole is also used to help prevent the cancer from returning. Some breast cancers are made to grow faster by a natural hormone called estrogen.



10x1x5 Tablets (Blister)

M.R.P. 164.13/-Per 10 Tablets

Levosulpiride-25

Composition :

Each uncoated tablet contains:
 Levosulpiride 25 mg.
 Excipients q.s.

Indication : Levosulpiride belongs to the class of antipsychotic medicines. However, it is primarily used for its gut motility properties i.e to improve food movement and treats disorders of the stomach and intestines.



10x10 Tablets (Alu Alu)

M.R.P. 82/-Per 10 Tablets

Levosulpiride-75

Composition :

Each uncoated tablet contains:
 Levosulpiride 75 mg.
 Excipients q.s.

Indication : Levosulpiride belongs to the class of antipsychotic medicines. However, it is primarily used for its gut motility properties i.e to improve food movement and treats disorders of the stomach and intestines.



10x10 Tablets (Blister)

M.R.P. 145/-Per 10 Tablets

Linezolid-600

Composition :

Each film coated tablet contains :
 Linezolid IP 600mg.
 Excipients q.s.

Indication : Linezolid is used to treat infections, including pneumonia, and infections of the skin . Linezolid is in a class of antibacterials called oxazolidinones. It works by stopping the growth of bacteria. Antibiotics such as linezolid will not work for colds, flu, and other viral infections



10x1x10 Tablets (Alu Alu)

M.R.P. 356.20/-Per 10 Tablets

Lisign-2.5

Composition :

Each uncoated tablet contains :
 Lisinopril IP
 Eq. to Anhydrous Lisinopril 2.5mg
 Excipients q.s.

Indication : Lisinopril is a medicine to treat high blood pressure and heart failure. It's also prescribed after a heart attack and in diabetic kidney disease. Lisinopril helps prevent future strokes and heart attacks. It also improves your survival if you're taking it after a recent heart attack or for heart failure.



10x10 Tablets (Blister)

M.R.P. 23/-Per 10 Tablets

Lisign-5

Composition :

Each uncoated tablet contains :
 Lisinopril IP
 Eq. to Anhydrous Lisinopril 5mg
 Excipients q.s.

Indication : Lisinopril is a medicine to treat high blood pressure and heart failure. It's also prescribed after a heart attack and in diabetic kidney disease. Lisinopril helps prevent future strokes and heart attacks. It also improves your survival if you're taking it after a recent heart attack or for heart failure.



10x10 Tablets (Blister)

M.R.P. 42/-Per 10 Tablets

Lisign-10

Composition :

Each uncoated tablet contains :
 Lisinopril IP
 Eq. to Anhydrous Lisinopril 10mg
 Excipients q.s.

Indication : Lisinopril is a medicine to treat high blood pressure and heart failure. It's also prescribed after a heart attack and in diabetic kidney disease. Lisinopril helps prevent future strokes and heart attacks. It also improves your survival if you're taking it after a recent heart attack or for heart failure.



10x10 Tablets (Blister)

M.R.P. 76/-Per 10 Tablets

Livogrid-250

Composition :

Each film coated tablet contains :
 Levetiracetam IP 250 mg.
 Excipients q.s.
 Colour: Yellow Oxide of Iron &
 Titanium Dioxide IP

Indication : Levetiracetam is an anticonvulsant used in combination with other medications to treat seizure disorders (epilepsy). Levetiracetam has been shown to decrease the number of seizures in adults and children. It is not known how it works to prevent seizures.

10x10 Tablets (Blister)
 M.R.P. 62.30/-Per 10 Tablets



Livogrid-500

Composition :

Each film coated tablet contains :
 Levetiracetam IP 500 mg.
 Excipients q.s.
 Colour: Yellow Oxide of Iron &
 Titanium Dioxide IP

Indication : Levetiracetam is an anticonvulsant used in combination with other medications to treat seizure disorders (epilepsy). Levetiracetam has been shown to decrease the number of seizures in adults and children. It is not known how it works to prevent seizures.

10x10 Tablets (Blister)
 M.R.P. 145.30/-Per 10 Tablets



Livogrid-750

Composition :

Each film coated tablet contains :
 Levetiracetam IP 750 mg.
 Excipients q.s.
 Colour: Red Oxide Of Iron
 & Titanium Dioxide IP

Indication : Levetiracetam is an anticonvulsant used in combination with other medications to treat seizure disorders (epilepsy). Levetiracetam has been shown to decrease the number of seizures in adults and children. It is not known how it works to prevent seizures.

10x10 Tablets (Blister)
 M.R.P. 190/-Per 10 Tablets



Livogrid-1000

Composition :

Each film coated tablet contains :
 Levetiracetam IP 1000 mg.
 Excipients q.s.
 Colour: Yellow Oxide of Iron &
 Titanium Dioxide IP

Indication : Levetiracetam is an anticonvulsant used in combination with other medications to treat seizure disorders (epilepsy). Levetiracetam has been shown to decrease the number of seizures in adults and children. It is not known how it works to prevent seizures.

10x10 Tablets (Blister)
 M.R.P. 360/-Per 10 Tablets



Loriday

Composition :

Each uncoated tablet contains :
 Loratadine BP 10mg.
 Excipients q.s.
 Colour : Erythrosine

Indication : Loratadine is indicated for the symptomatic relief of allergy such as hay fever (allergic rhinitis), urticaria (hives), chronic idiopathic urticaria, and other skin allergies. For allergic rhinitis, loratadine is indicated for both nasal and eye symptoms - sneezing, runny nose, and itchy or burning eyes.

4x5x10 Tablets (Blister)
 M.R.P. 48/-Per 10 Tablets




Lozaart[®]-25

Composition :

Each film coated tablet contains :
 Losartan Potassium IP 25 mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Losartan is a medicine widely used to treat high blood pressure and heart failure, and to protect your kidneys if you have both kidney disease and diabetes. Losartan helps to prevent future strokes, heart attacks and kidney problems.

4x5x10 Tablets (Blister)
 M.R.P. 28.24/-Per 10 Tablets




Lozaart[®]-50

Composition :

Each film coated tablet contains :
 Losartan Potassium IP 50 mg.
 Excipients q.s
 Colour : Titanium Dioxide IP

Indication : Losartan is a medicine widely used to treat high blood pressure and heart failure, and to protect your kidneys if you have both kidney disease and diabetes. Losartan helps to prevent future strokes, heart attacks and kidney problems.



4x5x10 Tablets (Blister)
M.R.P. 47.98/-Per 10 Tablets

Lozaart[®]-AM

Composition :

Each film coated tablet contains :
 Losartan Potassium IP 50mg.
 Amlodipine Besylate IP eq. to Amlodipine 5 mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Losartan + Amlodipine is a combination of two medicines: Losartan and Amlodipine which lower blood pressure effectively. Losartan is an angiotensin receptor blocker (ARB) and Amlodipine is a calcium channel blocker (CCB).




4x5x10 Tablets (Blister)
M.R.P. 94.16/-Per 10 Tablets

Lozaart[®]-H

Composition :

Each film coated tablet contains :
 Losartan Potassium IP 50mg.
 Hydrochlorothiazide IP 12.5.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : This drug is used to treat high blood pressure. It is also used to lower the risk of strokes in patients with high blood pressure and an enlarged heart. Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems.



4x5x10 Tablets (Blister)
M.R.P. 81/-Per 10 Tablets

Linadia-5

Composition :

Each uncoated tablet contains :
 Linagliptin 5mg.
 Excipients q.s.
 Colour : Red Oxide of Iron & Titanium Dioxide IP

Indication : Linagliptin is a medicine used to treat type 2 diabetes. Type 2 diabetes is a condition where the body does not make enough insulin, or the insulin that it makes does not work properly. This can cause high blood sugar levels (hyperglycaemia).



10x10 Tablets (Blister)
M.R.P. 309/-Per 10 Tablets

Linadia-2.5/500

Composition :

Each film coated tablet contains :
 Linagliptin 2.5mg.
 Metformin Hydrochloride IP 500mg.
 Excipients q.s.
 Colour : Red oxide of iron & Titanium Dioxide IP

Indication : Linagliptin and metformin combination is used to treat high blood sugar levels caused by type 2 diabetes. Linagliptin helps to control blood sugar levels by increasing substances in the body that make the pancreas release more insulin.



10x10 Tablets (Blister)
M.R.P. 179/-Per 10 Tablets

Linadia-2.5/1000

Composition :

Each film coated tablet contains :
 Linagliptin 2.5mg.
 Metformin Hydrochloride IP 1000mg.
 Excipients q.s.
 Colour : Red oxide of iron & Titanium Dioxide IP

Indication : Linagliptin and metformin combination is used to treat high blood sugar levels caused by type 2 diabetes. Linagliptin helps to control blood sugar levels by increasing substances in the body that make the pancreas release more insulin.



10x10 Tablets (Blister)
M.R.P. 286/-Per 10 Tablets

Labetalok-100

Composition :

Each uncoated tablet contains:
 Labetalol Hydrochloride IP 100mg.
 Excipients q.s.



Indication : Labetalol is used with or without other medications to treat high blood pressure (hypertension). Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. This medication is both an alpha blocker and beta blocker.

10x10 Tablets (Alu Alu)
 M.R.P. 169.68/-Per 10 Tablets

Mbgron-50

Composition :

Each film coated extended release tablet contains :
 Mirabegron 50mg.
 Excipients q.s.
 Colour : Yellow oxide of Iron



Indication : Mirabegron is used in adults to treat overactive bladder with symptoms of frequent or urgent urination and urinary incontinence. Mirabegron is sometimes used together with another medicine called solifenacin (Vesicare).

10x1x10 Tablets (Alu-Alu)
 M.R.P. 360/-Per 10 Tablets

Medmycogest-10

Composition :

Each uncoated tablet contains :
 Medroxyprogesterone Acetate IP 10 mg.
 Excipients q.s.



Indication : Medroxyprogesterone is used to treat amenorrhea (unusual stopping of menstrual periods) and abnormal uterine bleeding. It is also used to prevent endometrial hyperplasia (thickening of the lining of the uterus or womb) in women who are taking conjugated estrogens.

10x10 Tablets (Alu Alu)
 M.R.P. 66.97/-Per 10 Tablets

Mbgron-25

Composition

Each film coated extended release tablet contains :
 Mirabegron 25 mg
 Excipients q.s.
 Colour : Red Oxide of Iron.



Indication : Mirabegron is used in adults to treat overactive bladder with symptoms of frequent or urgent urination and urinary incontinence. Mirabegron is sometimes used together with another medicine called solifenacin (Vesicare).

10x1x10 Tablets (Alu Alu)
 M.R.P. 195/-Per 10 Tablets

Mebaus-135

Composition :

Each film coated tablet contains :
 Mebeverine Hydrochloride IP 135mg.
 Excipients q.s.
 Colour : Yellow Oxide of Iron & Titanium Dioxide IP



Indication : Mebeverine hydrochloride. This belongs to a group of medicines called antispasmodics. This medicine is used to treat symptoms of irritable bowel syndrome (IBS) and similar problems such as chronic irritable colon, spastic constipation, mucous colitis and spastic colitis.

10x10 Tablets (Blister)
 M.R.P. 132/-Per 10 Tablets

Menata-5

Composition :

Each film coated tablet contains :
 Memantine Hydrochloride USP 5 mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP



Indication : Memantine is used to treat moderate to severe confusion (dementia) related to Alzheimer's disease. It does not cure Alzheimer's disease, but it may improve memory, awareness, and the ability to perform

10x10 Tablets (Blister)
 M.R.P. 85/-Per 10 Tablets

Menata-10

Composition :

Each film coated tablet contains :
 Memantine Hydrochloride USP 10 mg. q.s.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Memantine is used to treat moderate to severe confusion (dementia) related to Alzheimer's disease. It does not cure Alzheimer's disease, but it may improve memory, awareness, and the ability to perform



10x10 Tablets (Blister)
 M.R.P. 150/-Per 10 Tablets

Menata-D

Composition :

Each film coated tablet contains :
 Donepezil Hydrochloride IP 5mg
 Memantine hydrochloride USP 10 mg. q.s.
 Excipients q.s.
 Colour : Red oxide of Iron &

Indication : This combination medication is used to treat moderate to severe confusion (dementia) related to Alzheimer's disease. It does not cure Alzheimer's disease, but it may improve memory, awareness, and the ability to perform daily functions.



10x10 Tablets (Blister)
 M.R.P. 138.16/-Per 10 Tablets

Mesalov[®]-400

Composition :

Each delayed release tablet contains :
 Mesalamine IP 400 mg
 Excipients q.s.
 Colour : Red oxide of Iron

Indication : Mesalamine is used to treat ulcerative colitis (a condition which causes swelling and sores in the lining of the colon [large intestine] and rectum) and also to maintain improvement of ulcerative colitis symptoms. Mesalamine is in a class of medications called anti-inflammatory agents.



10x10 Tablets (Blister)
 M.R.P. 112/-Per 10 Tablets

Mesalov[®]-800

Composition :

Each film coated delayed release tablet contains :
 Mesalamine USP 800 mg
 Excipients q.s.

Indication : Mesalamine is used to treat ulcerative colitis (a condition which causes swelling and sores in the lining of the colon [large intestine] and rectum) and also to maintain improvement of ulcerative colitis symptoms. Mesalamine is in a class of medications called anti-inflammatory agents.



10x10 Tablets (Blister)
 M.R.P. 215/-Per 10 Tablets

Mesalov[®]-1200

Composition :

Each film coated delayed release tablet contains :
 Mesalamine USP 1200 mg
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Mesalamine is used to treat ulcerative colitis (a condition which causes swelling and sores in the lining of the colon [large intestine] and rectum) and also to maintain improvement of ulcerative colitis symptoms. Mesalamine is in a class of medications called anti-inflammatory agents.



10x10 Tablets (Blister)
 M.R.P. 250/-Per 10 Tablets

Metadoop-2.5

Composition

Each film-coated tablet contains :
 Midodrine Hydrochloride USP 2.5mg
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Midodrine is used to treat orthostatic hypotension (sudden fall in blood pressure that occurs when a person assumes a standing position). Midodrine is in a class of medications called alpha-adrenergic agonists. It works by causing blood vessels to tighten, which increases blood pressure.



10x10 Tablets (Alu Alu)
 M.R.P. 970/-Per 10 Tablets

Methodol-4

Composition :

Each uncoated tablet contains :
 Methylprednisolone IP 4 mg.
 Excipients q.s.

Indication : It relieves inflammation (swelling, heat, redness, and pain) and is used to treat certain forms of arthritis; skin, blood, kidney, eye, thyroid, and intestinal disorders (e.g., colitis); severe allergies; and asthma. Methylprednisolone is also used to treat certain types of cancer.



20x10 Tablets (Blister)
M.R.P. 65/-Per 10 Tablets

Methodol-16

Composition :

Each uncoated tablet contains :
 Methylprednisolone IP 16 mg.
 Excipients q.s.

Indication : It relieves inflammation (swelling, heat, redness, and pain) and is used to treat certain forms of arthritis; skin, blood, kidney, eye, thyroid, and intestinal disorders (e.g., colitis); severe allergies; and asthma. Methylprednisolone is also used to treat certain types of cancer.



20x10 Tablets (Alu Alu)
M.R.P. 99.56/-Per 10 Tablets

Methotie-5

Composition :

Each uncoated tablet contains:
 Methotrexate IP 5mg.
 Excipients q.s.

Indication : Methotrexate is used to treat certain types of cancer or to control severe psoriasis or rheumatoid arthritis that has not responded to other treatments. It may also be used to control juvenile rheumatoid arthritis. Methotrexate belongs to a class of drugs known as antimetabolites.



10x10 Tablets (Alu Alu)
M.R.P. 98.91/-Per 10 Tablets

Methodol-8

Composition :

Each uncoated tablet contains :
 Methylprednisolone IP 8 mg.
 Excipients q.s.

Indication : It relieves inflammation (swelling, heat, redness, and pain) and is used to treat certain forms of arthritis; skin, blood, kidney, eye, thyroid, and intestinal disorders (e.g., colitis); severe allergies; and asthma. Methylprednisolone is also used to treat certain types of cancer.



20x10 Tablets (Alu Alu)
M.R.P. 57/-Per 10 Tablets

Methotie-2.5

Composition :

Each uncoated tablet contains:
 Methotrexate IP 2.5mg.
 Excipients q.s.

Indication : Methotrexate is used to treat certain types of cancer or to control severe psoriasis or rheumatoid arthritis that has not responded to other treatments. It may also be used to control juvenile rheumatoid arthritis. Methotrexate belongs to a class of drugs known as antimetabolites.



10x10 Tablets (Alu Alu)
M.R.P. 55.80/-Per 10 Tablets

Methotie-7.5

Composition :

Each uncoated tablet contains:
 Methotrexate IP 7.5mg.
 Excipients q.s.

Indication : Methotrexate is used to treat certain types of cancer or to control severe psoriasis or rheumatoid arthritis that has not responded to other treatments. It may also be used to control juvenile rheumatoid arthritis. Methotrexate belongs to a class of drugs known as antimetabolites.



10X10 Tablets (Alu Alu)
M.R.P. 135.03/-Per 10 Tablets

Methotie-10

Composition :

Each uncoated tablet contains:
 Methotrexate IP 10mg.
 Excipients q.s.

Indication : Methotrexate is used to treat certain types of cancer or to control severe psoriasis or rheumatoid arthritis that has not responded to other treatments. It may also be used to control juvenile rheumatoid arthritis. Methotrexate belongs to a class of drugs known as antimetabolites.



10x10 Tablets (Alu Alu)
 M.R.P. 139.40/-Per 10 Tablets

Methyq-G

Composition :

Each film coated tablet contains
 Gabapentin IP 300mg.
 Methylcobalamin IP 500 mcg.
 Excipients q.s.
 Colour : Red oxide of Iron

Indication:

Gabapentin+Methylcobalamin is used in the treatment of neuropathic pain. Gabapentin + Methylcobalamin is a combination of two medicines: Gabapentin and Methylcobalamin. Gabapentin is an alpha 2 delta ligand which decreases pain by modulating calcium channel activity of the nerve



10x10 Tablets (Alu Alu)
 M.R.P. 180/-Per 10 Tablets

Metaus-600 ER

Composition

Each film-coated extended release tablet contains :
 Metronidazole IP 600mg.
 Excipients q.s.
 Colour : Red oxide of Iron, Yellow Oxide Iron & Titanium Dioxide IP

Indication : Metronidazole is an antibiotic. It's used to treat skin infections, rosacea and mouth infections, including infected gums and dental abscesses. It's also used to treat conditions such as bacterial vaginosis and pelvic inflammatory disease.



10x10 Tablets (Alu Alu)
 M.R.P. 85/-Per 10 Tablets

Metrate-25

Composition :

Each uncoated tablets contains :
 Metoprolol Tartrate IP 25mg.
 Excipients q.s.

Indication : Metoprolol is used with or without other medications to treat high blood pressure (hypertension). Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. This medication is also used to treat chest pain (angina) and to improve survival after a heart attack.



20x10 Tablets (Blister)
 M.R.P. 34/-Per 10 Tablets

Metrate-50

Composition :

Each uncoated tablets contains :
 Metoprolol Tartrate IP 50mg.
 Excipients q.s.

Indication : Metoprolol is used with or without other medications to treat high blood pressure (hypertension). Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. This medication is also used to treat chest pain (angina) and to improve survival after a heart attack.



20x10 Tablets (Blister)
 M.R.P. 58.60/-Per 10 Tablets

Metrate-AM 50

Composition :

Each film coated tablet contains :
 Metoprolol Succinate IP 50mg.
 (As prolonged-release form)
 Amlodipine Besylate IP
 eq. to Amlodipine 5mg
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Metoprolol is used with or without other medications to treat high blood pressure (hypertension). Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. This medication is also used to treat chest pain (angina) and to improve survival after a heart attack.



10X10 Tablets (Alu Alu)
 M.R.P. 66/-Per 10 Tablets

Metrate-XL 12.5

Each film coated extended release tablets contains :
 Metoprolol Succinate IP 11.875mg.
 Eq. to Metoprolol Tartrate 12.5mg.
 Excipients q.s.
 Colour: Titanium Dioxide IP

Metoprolol is used with or without other medications to treat high blood pressure (hypertension). Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. This medication is also used to treat chest pain (angina) and to improve survival after a heart attack.



10x10 Tablets (Alu Alu)
M.R.P. 34.50/-Per 10 Tablets

Metrate XL 50

Composition :
 Each film coated extended release tablets contains :
 Metoprolol Succinate IP 47.5mg.
 Eq. to Metoprolol Tartrate 50mg.
 Excipients q.s.
 Colour: Titanium Dioxide IP

Indication : Metoprolol is used with or without other medications to treat high blood pressure (hypertension). Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. This medication is also used to treat chest pain (angina) and to improve survival after a heart attack.



10X10 Tablets (Alu Alu)
M.R.P. 65/-Per 10 Tablets

Mipholate-S

Composition :
 Each enteric coated tablet contains :
 Mycophenolate Sodium USP
 Eq. to Mycophenolic Acid 360mg.
 Excipients q.s.
 Colour : Sunset Yellow

Indication : Mycophenolate is used in combination with other medications to keep your body from attacking and rejecting your transplanted kidney. It belongs to a class of medications called immunosuppressants



10X10 Tablets (Alu Alu)
M.R.P. 875/-Per 10 Tablets

Metrate XL 25

Composition :
 Each film coated extended release tablets contains :
 Metoprolol Succinate IP 23.75mg.
 Eq. to Metoprolol Tartrate 25mg.
 Excipients q.s.
 Colour: Titanium Dioxide IP

Indication : Metoprolol is used with or without other medications to treat high blood pressure (hypertension). Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. This medication is also used to treat chest pain (angina) and to improve survival after a heart attack.



10X10 Tablets (Alu Alu)
M.R.P. 47/-Per 10 Tablets

Metrate XL-100

Composition :
 Each film coated extended release tablets contains :
 Metoprolol Succinate IP 95mg.
 Eq. to Metoprolol Tartrate 100mg.
 Excipients q.s.
 Colour: Titanium Dioxide IP

Indication : Metoprolol is used with or without other medications to treat high blood pressure (hypertension). Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. This medication is also used to treat chest pain (angina) and to improve survival after a heart attack.



10X10 Tablets (Alu Alu)
M.R.P. 92.89/-Per 10 Tablets

Mipholate-M

Composition :
 Each film coated tablet contains :
 Mycophenolate Mofetil IP 500mg.
 Excipients q.s.
 Colour : Red Oxide of Iron

Indication : Mycophenolate is used in combination with other medications to keep your body from attacking and rejecting your transplanted organ (such as kidney, liver, heart). It belongs to a class of medications called immunosuppressants.



10x10 Tablets (Alu Alu)
M.R.P. 737.90/-Per 10 Tablets

Mitroja-7.5

Composition :

Each film coated tablet contains :
 Mirtazapine IP 7.5mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Mirtazapine is the generic form of the brand-name antidepressant Remeron, which is used to treat depression and other ailments. In some patients, depression may be associated with abnormal levels of neurotransmitters, the brain chemicals that nerve cells use to communicate with each other.



10X10 Tablets (Blister)
 M.R.P. 58/-Per 10 Tablets

Mitroja-15

Composition :

Each film coated tablet contains :
 Mirtazapine IP 15mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Mirtazapine is the generic form of the brand-name antidepressant Remeron, which is used to treat depression and other ailments. In some patients, depression may be associated with abnormal levels of neurotransmitters, the brain chemicals that nerve cells use to communicate with each other.



10X10 Tablets (Blister)
 M.R.P. 95/-Per 10 Tablets

MOXAUS 0.2

Composition :

Each film coated tablet contains :
 Moxonidine BP 0.2mg
 Excipients q.s.

Indication : In patients with mild to moderate hypertension, moxonidine reduces blood pressure (BP) as effectively as most first-line antihypertensives when used as monotherapy and is also an effective adjunctive therapy in combination with other antihypertensive agents.



10x10 Tablets (Alu Alu)
 M.R.P. 72.75/-Per 10 Tablets

MOXAUS 0.3

Composition :

Each film coated tablet contains :
 Moxonidine BP 0.3mg
 Excipients q.s.

Indication : In patients with mild to moderate hypertension, moxonidine reduces blood pressure (BP) as effectively as most first-line antihypertensives when used as monotherapy and is also an effective adjunctive therapy in combination with other antihypertensive agents.



10x10 Tablets (Alu Alu)
 M.R.P. 122/-Per 10 Tablets

Moxiyst-400

Composition :

Each film coated tablet contains :
 Moxifloxacin Hydrochloride BP 400mg.
 Excipients q.s.
 Colour : Red Oxide of Iron & Titanium Dioxide IP

Indication : Moxifloxacin is used to treat a variety of bacterial infections. This medication belongs to a class of drugs called quinolone antibiotics. It works by stopping the growth of bacteria. This antibiotic treats only bacterial infections.



20x5 Tablets (Blister)
 M.R.P. 127/-Per 5 Tablets

Mycogest[®]-SR 200

Each film coated sustained release tablet contains :

Progesterone IP 200mg.
 (Natural Micronized)
 Excipients q.s.

Progesterone is used to cause menstrual periods in women who have not yet reached menopause but are not having periods due to a lack of progesterone in the body. It is also used to prevent overgrowth in the lining of the uterus in postmenopausal women who are receiving estrogen hormone replacement therapy.



10x1x10 Tablets (Blister)
 M.R.P. 280/-Per 10 Tablets

Natigone-60

Composition

Each film-coated tablet contains :
 Nateglinide 60mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Nateglinide is used together with diet and exercise to control blood sugar levels in patients with type 2 diabetes. It works by causing your pancreas to release more insulin into the blood stream. This medicine does not help patients who have insulin-dependent or type 1 diabetes.



10x10 Tablets (Alu Alu)

M.R.P. 96/-Per 10 Tablets

Naproxen-250

Composition :

Each film coated tablet contains:
 Naproxen Sodium PH EUR 250mg.

Indication : Naproxen is used to relieve pain from various conditions such as headache, muscle aches, tendonitis, dental pain, and menstrual cramps. It also reduces pain, swelling, and joint stiffness caused by arthritis, bursitis, and gout attacks. This medication is known as a nonsteroidal anti-inflammatory drug (NSAID).



10x10 Tablets (Blister)

M.R.P. 74/-Per 10 Tablets

Naproxen-D

Composition :

Each film coated tablet contains :
 Naproxen Sodium PH EUR 500mg
 Domperidone IP 10mg
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Naproxen+Domperidone is used to prevent Migraine. Naproxen + Domperidone is a combination of two medicines: Naproxen and Domperidone. Naproxen is a non-steroidal anti-inflammatory drug (NSAID) which works by blocking the release of certain chemical messengers that cause pain due to migraine.



10x10 Tablets (Blister)

M.R.P. 191/-Per 10 Tablets

Natigone-M

Composition

Each film-coated tablet contains :
 Nateglinide 60mg.
 Metformin Hydrochloride IP 500mg.
 (as sustained release)
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Nateglinide alone and in combination with metformin improves glycemic control by reducing mealtime glucose levels in type 2 diabetes. Diabetes Care.



10x10 Tablets (Alu Alu)

M.R.P. 295/-Per 10 Tablets

Naproxen-500

Composition :

Each film coated tablet contains:
 Naproxen Sodium PH EUR 500mg.
 Excipients q.s.

Indication : Naproxen is used to relieve pain from various conditions such as headache, muscle aches, tendonitis, dental pain, and menstrual cramps. It also reduces pain, swelling, and joint stiffness caused by arthritis, bursitis, and gout attacks. This medication is known as a nonsteroidal anti-inflammatory drug (NSAID).



10X10 Tablets (Blister)

M.R.P. 178/-Per 10 Tablets

Navsum-400

Composition

Each enteric coated tablet contains :
 Ademetionine 1,4-Butane Disulfonate
 Eq. to Ademetionine Ion 400mg.
 Excipients q.s.
 Colour : Sunset Yellow

Indication : It is produced from methionine (amino acid) and is used to treat liver disease, depression, and osteoarthritis. ADEMATIONINE ensures sufficient levels of glutathione in the liver are maintained to minimize liver damage. It also prevents the increase of lipids associated with cirrhosis outside the liver.



10x1x10 Tablets (Alu Alu)

M.R.P. 1125/-Per 10 Tablets

N-Dril-5

Composition :

Each uncoated tablet contains :
 Nicorandil IP 5 mg
 Excipients q.s.



Indication : Nicorandil is a medicine used to prevent and treat chest pain caused by angina. Your doctor will usually prescribe nicorandil when other heart medicines have not worked or are not suitable for you.

10X10 Tablets (Alu Alu)
 M.R.P. 105/-Per 10 Tablets

N-Dril-10

Composition :

Each uncoated tablet contains :
 Nicorandil IP 10 mg
 Excipients q.s.



Indication : Nicorandil is a medicine used to prevent and treat chest pain caused by angina. Your doctor will usually prescribe nicorandil when other heart medicines have not worked or are not suitable for you.

10X10 Tablets (Alu Alu)
 M.R.P. 175/-Per 10 Tablets

Nebawol-2.5

Composition :

Each uncoated tablet contains :
 Nebivolol Hydrochloride IP
 Eq. to Nebivolol 2.5 mg.
 Excipients q.s.



Indication : Nebivolol is used to treat high blood pressure. Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. This medication belongs to a class of drugs known as beta blockers.

10X10 Tablets (Alu Alu)
 M.R.P. 70/-Per 10 Tablets

Nebawol-5

Composition :

Each uncoated tablet contains :
 Nebivolol Hydrochloride IP
 Eq. to Nebivolol 5mg.
 Excipients q.s.



Indication : Nebivolol is used to treat high blood pressure. Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. This medication belongs to a class of drugs known as beta blockers.

10X10 Tablets (Alu Alu)
 M.R.P. 69/-Per 10 Tablets

Nefysity-10

Composition:

Each film coated tablet contains:
 Nifedipine IP 10 mg
 Colours : Ferric Oxide USPNF
 Yellow & Titanium Dioxide IP



Indication : Nifedipine is a medicine used to treat high blood pressure. If you have high blood pressure, taking nifedipine helps to prevent future heart disease, heart attacks and strokes. Nifedipine is also used to prevent chest pain caused by angina.

20x10 Tablets (Blister)
 M.R.P. 14.38/-Per 10 Tablets

Nefysity-20

Composition :

Each film coated extended release tablet contains :
 Nifedipine IP 20mg.
 Excipients q.s.
 Colour : Sunset Yellow FCF




Indication : Nifedipine is a medicine used to treat high blood pressure. If you have high blood pressure, taking nifedipine helps to prevent future heart disease, heart attacks and strokes. Nifedipine is also used to prevent chest pain caused by angina.

20x10 Tablets (Blister)
 M.R.P. 23.50/-Per 10 Tablets

Nekszole[®]-40

Composition :

Each enteric coated tablet contains :
 Esomeprazole
 Magnesium Trihydrate IP 40mg.
 Eq. to Esomeprazole q.s.
 Excipients q.s.
 Colour : Red oxide of Iron & Titanium Dioxide IP



Indication : Esomeprazole reduces the amount of acid your stomach makes. It's used for heartburn, acid reflux and gastro-oesophageal reflux disease (GORD) - GORD is when you keep getting acid reflux. It's also taken to prevent and treat stomach ulcers.

10X10 Tablets (Alu Alu)
 M.R.P. 130/-Per 10 Tablets

Newcontin-6.4

Each uncoated controlled release tablet contains :
 Diluted Nitroglycerin IP 6.4
 Eq. to Nitroglycerin q.s.
 Excipients q.s.
 Colour : Erythrosine



is used to treat and prevent heart-related chest pain (angina). Angina occurs when the heart muscle is not getting enough blood. This medicine works by relaxing and widening blood vessels so blood can flow more easily to the heart.

1x30 Tablets
 M.R.P. 217/-Per 30 Tablets

Nicomode-2

Composition :

Each uncoated tablet contains :
 Nicoumalone IP 2 mg.
 Excipients q.s.



Indication : Nicoumalone is used to prevent and treat harmful blood clots, reducing the risk of a stroke or heart attack. This medication helps to keep blood flowing smoothly in your body by decreasing the amount of clotting proteins in the blood.

10X10 Tablets (Aluminium Strip)
 M.R.P. 82/-Per 10 Tablets

Newcontin-2.6

Composition :

Each uncoated controlled release tablet contains :
 Diluted Nitroglycerin IP 2.6
 Eq. to Nitroglycerin q.s.
 Excipients q.s.
 Colour : Erythrosine



Indication : is used to treat and prevent heart-related chest pain (angina). Angina occurs when the heart muscle is not getting enough blood. This medicine works by relaxing and widening blood vessels so blood can flow more easily to the heart.

1x30 Tablets
 M.R.P. 159/-Per 30 Tablets

Nicomode-1

Composition :

Each uncoated tablet contains :
 Nicoumalone IP 1 mg.
 Excipients q.s.



Indication : Nicoumalone is used to prevent and treat harmful blood clots, reducing the risk of a stroke or heart attack. This medication helps to keep blood flowing smoothly in your body by decreasing the amount of clotting proteins in the blood.

10X10 Tablets (Aluminium Strip)
 M.R.P. 60/-Per 10 Tablets

Nicomode-3

Composition :

Each uncoated tablet contains :
 Nicoumalone IP 3 mg.
 Excipients q.s.



Indication : Nicoumalone is used to prevent and treat harmful blood clots, reducing the risk of a stroke or heart attack. This medication helps to keep blood flowing smoothly in your body by decreasing the amount of clotting proteins in the blood.

10X10 Tablets (Aluminium Strip)
 M.R.P. 115/-Per 10 Tablets

Nicomode-4

Composition :

Each uncoated tablet contains :
 Nicoumalone IP 4 mg.
 Excipients q.s.



Indication : Nicoumalone is used to prevent and treat harmful blood clots, reducing the risk of a stroke or heart attack. This medication helps to keep blood flowing smoothly in your body by decreasing the amount of clotting proteins in the blood.

10X10 Tablets (Aluminium Strip)
 M.R.P. 148/-Per 10 Tablets

Nitrofurantoin-100 SR

Composition :

Each film coated sustained release tablet contains :
 Nitrofurantoin IP 100mg.
 Excipients q.s.
 Colour : Quinoline Yellow




Indication : This medication is used to treat or prevent certain urinary tract infections. This medication is an antibiotic that works by stopping the growth of bacteria.

10X10 Tablets (Blister)
 M.R.P. 89.90/-Per 10 Tablets

Northeal

Composition :

Each uncoated tablet contains :
 Norethisterone IP 5mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP



Indication : Norethisterone Tablets has many uses. You can take Norethisterone Tablets to treat or manage: · Heavy periods · Painful periods · Irregular periods or periods that are more frequent than normal · Premenstrual tension (PMT) · Endometriosis (where tissue from your womb is found outside your

6x5x10 Tablets (Blister)
 M.R.P. 63.50/-Per 10 Tablets

Nimotime-30

Composition :

Each film coated tablet contain :
 Nimodipine BP 30 mg.
 Excipients q.s.



Indication : Nimodipine is used to decrease brain damage that may be caused by a subarachnoid hemorrhage (bleeding in the space surrounding the brain that occurs when a weakened blood vessel in the brain bursts). Nimodipine is in a class of medications called calcium channel blockers.

10X10 Tablets (Alu Alu)
 M.R.P. 160/-Per 10 Tablets

Noraus-25

Composition :

Each film coated tablet contains :
 Nortriptyline Hydrochloride IP
 Eq. to Nortriptyline 25mg.
 Excipients q.s.




Indication : Nortriptyline is a tricyclic antidepressant. It affects chemicals in the brain that may be unbalanced in people with depression. Nortriptyline is used to treat symptoms of depression. Nortriptyline may also be used for purposes not listed in this medication guide

10X10 Tablets (Blister)
 M.R.P. 69/-Per 10 Tablets

Norfaux-400

Composition

Each film-coated tablet contains :
 Norfloxacin IP 400mg.
 Excipients q.s.
 Colour : Sunset Yellow FCF & Titanium Dioxide IP



Indication : Norfloxacin is used to treat certain bacterial infections in many different parts of the body. Norfloxacin may mask or delay the symptoms of syphilis. It is not effective against syphilis infections. Norfloxacin belongs to the class of medicines known as quinolone antibiotics.

6x5x10 Tablets (Blister)
 M.R.P. 85/-Per 10 Tablets



TABLETS

Obetech-5

Composition :

Each film coated tablet contains :
 Obeticholic Acid 5mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Obeticholic acid is used alone or in combination with ursodiol (Actigall, Urso) to treat primary biliary cholangitis (PBC; a type of liver disease that destroys bile ducts, which allows bile to stay in the liver and cause damage) in people who cannot take ursodiol or in people who were not treated successfully with ursodiol alone. Obeticholic acid is in a class of medications called farnesoid X receptor agonists. It works by decreasing the production of bile in the liver and increasing the removal of



10x10 Tablets (Alu Alu)
M.R.P. 195/-Per 10 Tablets

Obetech-10

Composition :

Each film coated tablet contains :
 Obeticholic Acid 10mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Obeticholic acid is used alone or in combination with ursodiol (Actigall, Urso) to treat primary biliary cholangitis (PBC; a type of liver disease that destroys bile ducts, which allows bile to stay in the liver and cause damage) in people who cannot take ursodiol or in people who were not treated successfully with ursodiol alone. Obeticholic acid is in a class of medications called farnesoid X receptor agonists. It works by decreasing the production of bile in the liver and increasing the removal of



10x10 Tablets (Alu Alu)
M.R.P. 380/-Per 10 Tablets

Olanzapine-5

Composition :

Each film coated tablet contains :
 Olanzapine IP 5mg
 Excipients q.s.
 Colour : Quinoline Yellow

Indication : Olanzapine is an antipsychotic medication that affects chemicals in the brain. Olanzapine is used to treat the symptoms of psychotic conditions such as schizophrenia and bipolar disorder (manic depression) in adults and children who are at least 13 years old.



10X10 Tablets (Blister)
M.R.P. 34.80/-Per 10 Tablets

Olanzapine-10

Composition :

Each film coated tablet contains :
 Olanzapine IP 10mg
 Excipients q.s.
 Colour : Quinoline Yellow

Indication : Olanzapine is an antipsychotic medication that affects chemicals in the brain. Olanzapine is used to treat the symptoms of psychotic conditions such as schizophrenia and bipolar disorder (manic depression) in adults and children who are at least 13 years old.



10X10 Tablets (Blister)
M.R.P. 69/-Per 10 Tablets

Olaphine-F

Composition

Each film-coated tablet contains :
 Olanzapine IP 10mg.
 Fluoxetine Hydrochloride IP eq. to Fluoxetine 20mg.
 Excipients q.s.

Indication : Olanzapine and fluoxetine combination is used to treat depression that is a part of bipolar disorder, and depression in patients who received other antidepressants that did not work well. This medicine should not be used to treat behavioral problems in older adult patients who have dementia or Alzheimer's disease.



10x10 Tablets (Blister)
M.R.P. 135/-Per 10 Tablets

Olmedose-10

Composition :

Each film coated tablet contains :
 Olmesartan Medoxomil IP 10 mg
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Olmesartan is used to treat high blood pressure (hypertension). Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. Olmesartan belongs to a class of drugs called angiotensin receptor blockers (ARBs). It works by relaxing blood vessels so that blood can flow more easily.



10X10 Tablets (Alu Alu)
M.R.P. 38.90/-Per 10 Tablets

Olmedose-20

Composition :

Each film coated tablet contains :
 Olmesartan Medoxomil IP 20 mg
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Olmesartan is used to treat high blood pressure (hypertension). Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. Olmesartan belongs to a class of drugs called angiotensin receptor blockers (ARBs). It works by relaxing blood vessels so that blood can flow more easily.



10X10 Tablets (Alu Alu)
 M.R.P. 110/-Per 10 Tablets

Olmedose-40 H

Composition :

Each film coated Tablet contains :
 Olmesartan Medoxomil IP 40 mg.
 Hydrochlorothiazide IP 12.5 mg.
 Excipients q.s.
 Colour: Red Oxide of Iron

Indication : This drug is used to treat high blood pressure (hypertension). Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. This product contains two medications: olmesartan and hydrochlorothiazide.



10X10 Tablets (Alu Alu)
 M.R.P. 195/-Per 10 Tablets

Olmedose-40 Trio

Composition :

Each film coated tablet contains :
 Olmesartan Medoxomil IP 40 mg
 Amlodipine Besylate IP 5 mg
 Eq. to Amlodipine 5 mg
 Hydrochlorothiazide IP 12.5 mg
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Olmesartan medoxomil, amlodipine, hydrochlorothiazide) Tablets is a combination of an angiotensin receptor blocker, a calcium channel blocker, and a diuretic used to treat high blood pressure



10X10 Tablets (Alu Alu)
 M.R.P. 210/-Per 10 Tablets

Olmedose-20 H

Composition :

Each film coated tablet contains :
 Olmesartan Medoxomil IP 20 mg
 Hydrochlorothiazide IP 12.5mg
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Olmesartan is used to treat high blood pressure (hypertension). Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. Olmesartan belongs to a class of drugs called angiotensin receptor blockers (ARBs). It works by relaxing blood vessels so that blood can flow more easily.



10X10 Tablets (Alu Alu)
 M.R.P. 130/-Per 10 Tablets

Olmedose-40

Composition :

Each film coated tablet contains :
 Olmesartan Medoxomil IP 40 mg
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Olmesartan is used to treat high blood pressure (hypertension). Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. Olmesartan belongs to a class of drugs called angiotensin receptor blockers (ARBs). It works by relaxing blood vessels so that blood can flow more easily.



10X10 Tablets (Alu Alu)
 M.R.P. 175/-Per 10 Tablets

Olmedose-AM

Composition :

Each film coated tablets contains:
 Olmesartan Medoxomil IP 40 mg
 Amlodipine Besylate IP 5 mg
 Eq. to Amlodipine 5 mg
 Excipients q.s.
 Colour : Titanium Dioxide IP

Olmesartan is used to treat high blood pressure (hypertension). Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. Olmesartan belongs to a class of drugs called angiotensin receptor blockers (ARBs). It works by relaxing blood vessels so that blood can flow more easily.



10X10 Tablets (Alu Alu)
 M.R.P. 130/-Per 10 Tablets

Olmedose-C

Composition :

Each film coated tablet contains :
 Olmesartan Medoxomil IP 20 mg
 Chlorthalidone IP 12.5 mg
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Olmesartan medoxomil+ Chlorthalidone is used in the treatment of Hypertension (high blood pressure). Olmesartan Medoxomil + Chlorthalidone is a combination of two medicines: Olmesartan and chlorthalidone which lower blood pressure effectively. Olmesartan is an angiotensin receptor blocker (ARB)



10X10 Tablets (Alu Alu)
 M.R.P. 155/-Per 10 Tablets

Opimon-50

Composition

Each film-coated tablet contains :
 Opipramol Dihydrochloride 50mg.
 Excipients q.s.
 Colour : Quinoline Yellow & Titanium Dioxide IP

Indication : Opipramol 2-HCl (4-[3-(5H-dibenz[b,f]-azepine-5-yl)-propyl]-1-piperazine-ethanol dihydrochloride, CAS 315-72-0) is an atypical anxiolytic and antidepressive drug. It is a psychotropic drug commonly used for therapy of anxious-depressive states, somatoform disorders and general anxiety disorders (1-4).



10x10 Tablets (Blister)
 M.R.P. 108/-Per 10 Tablets

Otinium-40

Composition

Each film-coated tablet contains :
 Otilonium Bromide 40mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Otilonium bromide (OB) is a drug with spasmolytic activity belonging to quaternary ammonium derivatives and extensively used to treat patients affected by the Irritable Bowel Syndrome (IBS). Thanks to its peculiar pharmacokinetic, OB concentrates in the large bowel wall and acts locally.




10x10 Tablets (Alu Alu)
 M.R.P. 210/-Per 10 Tablets

Orthofill

Nutritional Value per tablet (approximate Value) :		%RDA Value
Energy Value	0.075 kcal	**
Protein	0.00 g	**
Carbohydrate	0.0189 g	**
Sugar	0.00 g	**
Fat	0.00 g	**
Each Film coated tablet contain (approx) :		%RDA Value
A blend of Extracts of Scutellaria Baicalensis and acacia catechu	500 mg.	**
Excipients	q.s.	

Indication : It helps in relieving body pains such as knee pain, joint pain and back pain. It helps in strengthening the muscles, joints and bones.



10x1x10 Tablets (Alu Alu)
 M.R.P. 490/-Per 10 Tablets

Opimon-100

Composition

Each film-coated tablet contains :
 Opipramol Dihydrochloride 100mg.
 Excipients q.s.
 Colour : Quinoline Yellow & Titanium Dioxide IP

Indication : Opipramol 2-HCl (4-[3-(5H-dibenz[b,f]-azepine-5-yl)-propyl]-1-piperazine-ethanol dihydrochloride, CAS 315-72-0) is an atypical anxiolytic and antidepressive drug. It is a psychotropic drug commonly used for therapy of anxious-depressive states, somatoform disorders and general anxiety disorders (1-4).



10x10 Tablets (Blister)
 M.R.P. 195/-Per 10 Tablets

OXYCBZ-150

Composition :

Each film coated tablet contains :
 Oxcarbazepine IP 150mg.
 Excipients q.s.
 Colour : Yellow oxide of Iron & Titanium Dioxide IP

Indication : Oxcarbazepine is an anticonvulsant used to reduce the occurrence of epileptic episodes, and is not intended to cure epilepsy. Oxcarbazepine is used alone or in combination with other medications for the treatment of focal (partial) seizures in adults.



10X10 Tablets (Blister)
 M.R.P. 149/-Per 10 Tablets

OXYCBAZ 300

Each film coated tablet contains :
Oxcarbazepine IP 300mg.
Excipients q.s.
Colour : Yellow oxide of Iron &
Titanium Dioxide IP

Oxcarbazepine is an anticonvulsant used to reduce the occurrence of epileptic episodes, and is not intended to cure epilepsy. Oxcarbazepine is used alone or in combination with other medications for the treatment of focal (partial) seizures in adults.



10X10 Tablets (Blister)
M.R.P. 299/-Per 10 Tablets

OXYCBAZ 450

Composition :
Each film coated tablet contains :
Oxcarbazepine IP 450mg.
Excipients q.s.
Colour : Yellow oxide of Iron &
Titanium Dioxide IP

Indication : Oxcarbazepine is an anticonvulsant used to reduce the occurrence of epileptic episodes, and is not intended to cure epilepsy. Oxcarbazepine is used alone or in combination with other medications for the treatment of focal (partial) seizures in adults.



10x10 Tablets (Blister)
M.R.P. 399/-Per 10 Tablets

Parest®-1000

Composition :
Each uncoated bilayered tablet contains:
Paracetamol I.P. 300 mg
(As Immediate Release)
Paracetamol I.P. 700 mg
(As Sustained Release)
Excipients q.s.
Colour : Sunset Yellow FCF

Indication : Paracetamol is a common painkiller used to treat aches and pain. It can also be used to reduce a high temperature. It's available combined with other painkillers and anti-sickness medicines. It's also an ingredient in a wide range of cold and flu remedies.



10x10 Tablets (Blister)
M.R.P. 99/-Per 10 Tablets

Parocare-12.5 CR

Composition :
Each enteric coated controlled release tablet contains :
Paroxetine Hydrochloride IP 12.5 mg.
Excipients q.s.
Colours : Yellow Oxide of Iron

Indication : Paroxetine is used to treat depression, panic attacks, obsessive-compulsive disorder (OCD), anxiety disorders, and post-traumatic stress disorder. It works by helping to restore the balance of a certain natural substance (serotonin) in the brain. Paroxetine is known as a selective serotonin reuptake inhibitor (SSRI).



10x10 Tablets (Alu Alu)
M.R.P. 120/-Per 10 Tablets

Parocare-25

Composition :
Each film coated tablet contains :
Paroxetine Hydrochloride Hemihydrate IP
Eq. to Anhydrous Paroxetine 25 mg
Excipients q.s.

Indication : Paroxetine is used to treat depression, panic attacks, obsessive-compulsive disorder (OCD), anxiety disorders, and post-traumatic stress disorder. It works by helping to restore the balance of a certain natural substance (serotonin) in the brain. Paroxetine is known as a selective serotonin reuptake inhibitor (SSRI).



10x10 Tablets (Alu Alu)
M.R.P. 170/-Per 10 Tablets

PAZOTRY 2.5

Composition :
Each uncoated sustained release tablet contains :
Prazosin Hydrochloride IP
Eq. to Prazosin 2.5mg.
Excipients q.s.

Indication : Prazosin is used with or without other medications to treat high blood pressure. Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems.



10x1x10 Tablets (Blister)
M.R.P. 90/-Per 10 Tablets

PAZOTRY-5

Composition :

Each uncoated sustained release tablet contains :
 Prazosin Hydrochloride IP 5mg.
 Eq. to Prazosin q.s.
 Excipients q.s.

Indication : Prazosin is used with or without other medications to treat high blood pressure. Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems.



10x10 Tablets (Blister)
M.R.P. 130/-Per 10 Tablets

Perindopril-4

Composition :

Each uncoated tablet contains:
 Perindopril Erbumine BP 4mg.
 Excipients q.s.

Indication : Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. This medication is also used to prevent heart attacks in people with a certain type of heart disease (stable coronary artery disease). Perindopril belongs to a class of drugs known as ACEinhibitors.



10X10 Tablets (Blister)
M.R.P. 125/-Per 10 Tablets

Perindopril-8

Composition :

Each uncoated tablet contains:
 Perindopril Erbumine BP 8mg.
 Excipients q.s.

Indication : Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. This medication is also used to prevent heart attacks in people with a certain type of heart disease (stable coronary artery disease). Perindopril belongs to a class of drugs known as ACEinhibitors.



10X10 Tablets (Blister)
M.R.P. 146/-Per 10 Tablets

PGNERV-NT

Composition :

Each film coated tablet contains :
 Pregabalin IP 75mg.
 (As Sustained Release)
 Nortriptyline Hydrochloride IP 10mg.
 Excipients q.s.
 Colour : Ponceau 4R

Indication : Pregabalin+Nortriptyline is used in the treatment of neuropathic pain. Pregabalin + Nortriptyline is a combination of two medicines: Pregabalin and Nortriptyline. Pregabalin is an alpha 2 delta ligand which decreases the pain by modulating calcium channel activity of the nerve cells.



10x10 Tablets (Alu Alu)
M.R.P. 195/-Per 10 Tablets

Pgnerv-M Forte

Composition :

Each uncoated bilayered tablet contains :
 Pregabalin IP 75 mg
 (Sustained Release)
 Methylcobalamin 1500 mcg
 Excipients q.s.

Indication : Pregabalin+methylcobalamin Belongs To The Class Of Medications 'anticonvulsants' Used In The Treatment Of Neuropathic Pain. Neuropathic Pain Is A Chronic (long-term) Progressive Nerve Disease That Causes Nerve Pain Due To Nerve Damage Or The Nervous System's Malfunctioning.



10x10 Tablets (Alu Alu)
M.R.P. 245/-Per 10 Tablets

PGNERV-NM

Composition :

Each film coated tablet contains :
 Pregabalin IP 75 mg
 (As Sustained release form)
 Nortriptyline HCl IPeq. to
 Nortriptyline 10 mg.
 Methylcobalamin IP 1500mcg.
 Excipients q.s.

Indication : This medication is used to treat pain caused by nerve damage due to diabetes, shingles (herpes zoster) infection, or spinal cord injury. This medication is also used to treat pain in people with fibromyalgia. It is also used with other medications to treat certain types of seizures (focal seizures).



10X10 Tablets (Alu Alu)
M.R.P. 275/-Per 10 Tablets

Pheuton-100

Composition :

Each film coated tablet contains :
 Phenytoin Sodium IP 100mg
 Excipients q.s.
 Colour : Titanium Dioxide

Indication : Phenytoin is used to control certain type of seizures, and to treat and prevent seizures that may begin during or after surgery to the brain or nervous system. Phenytoin is in a class of medications called anticonvulsants.



1X100 Tablets

M.R.P. 147/-Per 100 Tablets

Piocare-7.5

Composition :

Each uncoated tablet contains :
 Pioglitazone Hydrochloride IP 7.5 mg
 Eq. to Pioglitazone 7.5 mg
 Excipients q.s.

Indication : Pioglitazone is used with a diet and exercise program and sometimes with other medications, to treat type 2 diabetes (condition in which the body does not use insulin normally and therefore cannot control the amount of sugar in the blood). Pioglitazone is in a class of medications called thiazolidinediones.



10X10 Tablets (Blister)

M.R.P. 59/-Per 10 Tablets

Piocare-30

Indication :

Each uncoated tablet contains :
 Pioglitazone Hydrochloride IP 30mg
 Eq. to Pioglitazone 30mg
 Excipients q.s.

Indication : Pioglitazone is used with a diet and exercise program and sometimes with other medications, to treat type 2 diabetes (condition in which the body does not use insulin normally and therefore cannot control the amount of sugar in the blood). Pioglitazone is in a class of medications called thiazolidinediones.



10X10 Tablets (Blister)

M.R.P. 95/-Per 10 Tablets

Picorite-MD

Composition

Each uncoated mouth dissolving tablet contains :
 Prochlorperazine Maleate IP 5mg.

Indication : Prochlorperazine suppositories and tablets are used to control severe nausea and vomiting. Prochlorperazine tablets are also used to treat the symptoms of schizophrenia (a mental illness that causes disturbed or unusual thinking, loss of interest in life, and strong or inappropriate emotions).



20x10 Tablets (Aluminium)

M.R.P. 95/-Per 10 Tablets

Piocare-15

Composition :

Each uncoated tablet contains :
 Pioglitazone Hydrochloride IP 15mg
 Eq. to Pioglitazone 15mg
 Excipients q.s.

Indication : Pioglitazone is used with a diet and exercise program and sometimes with other medications, to treat type 2 diabetes (condition in which the body does not use insulin normally and therefore cannot control the amount of sugar in the blood). Pioglitazone is in a class of medications called thiazolidinediones.



10X10 Tablets (Blister)

M.R.P. 54/-Per 10 Tablets

Piracetam-400

Composition :

Each film coated tablet contains:
 Piracetam IP 400 mg
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Piracetam is said to act as a nootropic, a class of drugs designed to enhance memory and boost cognitive function. In alternative medicine, it's thought that piracetam can help increase brain function by promoting communication between the left and right hemispheres of the brain.



10x10 Tablets (Blister)

M.R.P. 85/-Per 10 Tablets

Piracetam-800

Composition :

Each film coated tablet contains:
 Piracetam IP 800mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Piracetam is said to act as a nootropic, a class of drugs designed to enhance memory and boost cognitive function. In alternative medicine, it's thought that piracetam can help increase brain function by promoting communication between the left and right hemispheres of the brain.



10X10 Tablets (Blister)
 M.R.P. 140/-Per 10 Tablets

Piroxicam-20

Composition :

Each uncoated tablet contains :
 Piroxicam IP 20mg
 Excipients q.s.

Indication : Piroxicam is used to relieve pain, tenderness, swelling, and stiffness caused by osteoarthritis (arthritis caused by a breakdown of the lining of the joints) and rheumatoid arthritis (arthritis caused by swelling of the lining of the joints). Piroxicam is in a class of medications called NSAIDs.



20x10 Tablets (Blister)
 M.R.P. 51.45/-Per 10 Tablets

Prasugrel-5

Composition :

Each film coated tablet contains:
 Prasugrel Hydrochloride eq. to Prasugrel 5mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Prasugrel is used with aspirin by patients with heart disease (recent heart attack, unstable angina) who undergo a certain heart procedure (angioplasty). This medication helps to prevent other serious heart/blood vessel problems (such as heart attacks, strokes, blood clots in stents).



10X10 Tablets (Alu Alu)
 M.R.P. 155/-Per 10 Tablets

Pirfedo-200

Composition :

Each film coated tablet contains :
 Pirfenidone IP 200mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Pirfenidone is used to treat a certain lung disease called idiopathic pulmonary fibrosis (IPF). This disease causes the lungs to get scarred and become stiff, making it hard to breathe. Pirfenidone may help slow down the worsening of your IPF.




10x1x15 Tablets
 M.R.P. 304.50 /-Per 15 Tablets

Pramday-0.125

Composition :

Each uncoated tablets contains :
 Pramipexole Dihydrochloride Monohydrate BP 0.125mg
 Excipients q.s.
 Colour : Erythrosin

Indication : Pramipexole is used alone or with other medications to treat Parkinson's disease. ... This medication is also used to treat a certain medical condition (restless legs syndrome - RLS) that causes an unusual urge to move the legs.



10x10 Tablets (Aluminium Strip)
 M.R.P. 59.50/-Per 10 Tablets

Prasugrel-10

Composition :

Each film coated tablet contains:
 Prasugrel Hydrochloride 10 mg.
 Excipients q.s.

Indication : Prasugrel is used with aspirin by patients with heart disease (recent heart attack, unstable angina) who undergo a certain heart procedure (angioplasty). This medication helps to prevent other serious heart/blood vessel problems (such as heart attacks, strokes, blood clots in stents).



10X10 Tablets (Alu Alu)
 M.R.P. 245/-Per 10 Tablets

Propong-10

Composition :

Each uncoated tablet contains :
 Propranolol Hydrochloride IP 10mg.

Indication : Propranolol is used to treat tremors, angina (chest pain), hypertension (high blood pressure), heart rhythm disorders, and other heart or circulatory conditions. It is also used to treat or prevent heart attack, and to reduce the severity and frequency of migraine headaches.



4x5x10 Tablets (Blister)

M.R.P. 13/-Per 10 Tablets

Propong-40

Composition :

Each uncoated tablet contains :
 Propranolol Hydrochloride IP 40mg.
 Excipients q.s.

Indication : Propranolol is used to treat tremors, angina (chest pain), hypertension (high blood pressure), heart rhythm disorders, and other heart or circulatory conditions. It is also used to treat or prevent heart attack, and to reduce the severity and frequency of migraine headaches.



4x5x10 Tablets (Blister)

M.R.P. 32/-Per 10 Tablets

Prucapride-1

Composition :

Each film coated tablet contains :
 Prucalopride Succinate eq. to Prucalopride 1mg.
 Excipients q.s.
 Colour : Red Oxide of Iron & yellow oxide of Iron & titanium dioxide IP

Indication : Prucalopride is used to treat chronic idiopathic (unknown cause) constipation (CIC). It works by increasing peristalsis (contractions) in the colon, leading to an increase in the number of bowel movements. This medicine is available only with your doctor's prescription.



10X10 Tablets (Alu Alu)

M.R.P. 148/-Per 10 Tablets

Prucapride-2

Composition :

Each film coated tablet contains :
 Prucalopride Succinate eq. to Prucalopride 2mg.
 Excipients q.s.
 Colour : Red Oxide of Iron & yellow oxide of Iron & titanium dioxide IP

Indication : Prucalopride is used to treat chronic idiopathic (unknown cause) constipation (CIC). It works by increasing peristalsis (contractions) in the colon, leading to an increase in the number of bowel movements. This medicine is available only with your doctor's prescription.



10X10 Tablets (Alu Alu)

M.R.P. 290 /-Per 10 Tablets

PYTB-500

Composition

Each uncoated tablet contains :
 Pyrazinamide IP 500mg.
 Excipients q.s.

Indication : Pyrazinamide kills or stops the growth of certain bacteria that cause tuberculosis (TB). It is used with other drugs to treat tuberculosis. This medication is sometimes prescribed for other uses; ask your doctor or pharmacist for more information.



10x10 Tablets (Blister)

M.R.P. 51.24 /-Per 10 Tablets

PYTB-750

Composition

Each uncoated tablet contains :
 Pyrazinamide IP 750mg.
 Excipients q.s.

Indication : Pyrazinamide kills or stops the growth of certain bacteria that cause tuberculosis (TB). It is used with other drugs to treat tuberculosis. This medication is sometimes prescribed for other uses; ask your doctor or pharmacist for more information.



10x10 Tablets (Blister)

M.R.P. 75.60 /-Per 10 Tablets

Qticare-100 SR

Composition :

Each film coated sustained release tablet contains :
 Quetiapine Fumarate IP 100 mg
 Eq. to Quetiapine
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : This medication is used to treat certain mental/mood conditions (such as schizophrenia, bipolar disorder, sudden episodes of mania or depression associated with bipolar disorder). Quetiapine is known as an anti-psychotic drug (atypical type).

10X10 Tablets (Blister)

M.R.P. 110/-Per 10 Tablets



Qticare-200 SR

Composition :

Each film coated sustained release tablet contains :
 Quetiapine Fumarate IP 200 mg
 Eq. to Quetiapine
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : This medication is used to treat certain mental/mood conditions (such as schizophrenia, bipolar disorder, sudden episodes of mania or depression associated with bipolar disorder). Quetiapine is known as an anti-psychotic drug (atypical type).

10x10 Tablets (Blister)

M.R.P. 125/-Per 10 Tablets



Qticare-300 SR

Composition :

Each film coated sustained release tablet contains :
 Quetiapine Fumarate IP 300 mg
 Eq. to Quetiapine
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : This medication is used to treat certain mental/mood conditions (such as schizophrenia, bipolar disorder, sudden episodes of mania or depression associated with bipolar disorder). Quetiapine is known as an anti-psychotic drug (atypical type).

10X10 Tablets (Blister)

M.R.P. 160/-Per 10 Tablets



Qticare-25

Composition :

Each film coated tablet contains :
 Quetiapine Fumarate IP 25 mg
 Eq. to Quetiapine
 Excipients q.s.

Indication : This medication is used to treat certain mental/mood conditions (such as schizophrenia, bipolar disorder, sudden episodes of mania or depression associated with bipolar disorder). Quetiapine is known as an anti-psychotic drug (atypical type).

10x10 Tablets (Blister)

M.R.P. 38/-Per 10 Tablets



Qticare-50

Composition :

Each film coated tablet contains :
 Quetiapine Fumarate IP 50mg.
 Eq. to Quetiapine
 Excipients q.s.

Indication : This medication is used to treat certain mental/mood conditions (such as schizophrenia, bipolar disorder, sudden episodes of mania or depression associated with bipolar disorder). Quetiapine is known as an anti-psychotic drug (atypical type).

10x10 Tablets (Blister)

M.R.P. 54/-Per 10 Tablets



Raminice-2.5

Composition :

Each uncoated tablet contains :
 Ramipril IP 2.5mg.
 Excipients q.s.

Indication : Ramipril is a medicine widely used to treat high blood pressure and heart failure. It's also prescribed after a heart attack. Ramipril helps prevent future strokes, heart attacks and kidney problems. It also improves your survival if you're taking it for heart failure or after a heart

20x10 Tablets (Blister)

M.R.P. 58.3/-Per 10 Tablets



Raminice-5

Composition :

Each uncoated tablet contains :
 Ramipril IP 5mg.
 Excipients q.s.

Indication : Ramipril is a medicine widely used to treat high blood pressure and heart failure. It's also prescribed after a heart attack. Ramipril helps prevent future strokes, heart attacks and kidney problems. It also improves your survival if you're taking it for heart failure or after a heart



20x10 Tablets (Blister)
 M.R.P. 81.76/-Per 10 Tablets

Raminice-10

Composition :

Each uncoated tablet contains :
 Ramipril IP 10mg.
 Excipients q.s.

Indication : Ramipril is a medicine widely used to treat high blood pressure and heart failure. It's also prescribed after a heart attack. Ramipril helps prevent future strokes, heart attacks and kidney problems. It also improves your survival if you're taking it for heart failure or after a heart



20x10 Tablets (Blister)
 M.R.P. 164/-Per 10 Tablets

Raniday-150

Composition :

Each film coated tablet contains :
 Ranitidine HCL IP
 Eq. to Ranitidine 150 mg.
 Colour : Sunset Yellow Lake &
 Titanium Dioxide IP

Indication : Ranitidine reduces the amount of acid your stomach makes. It's used for indigestion and heartburn and acid reflux. It is also used for gastro-oesophageal reflux disease (GORD) - this is when you keep getting acid reflux. Ranitidine is also taken to prevent and treat stomach ulcers.



20x3x10 Tablets (Aluminium Strip)
 M.R.P. 13.66/-Per 10 Tablets

Raniday-300

Composition

Each film coated tablet contains :
 Ranitidine Hydrochloride IP
 eq. to Ranitidine 300mg.
 Colour : Tartrazine & Brilliant Blue

Indication : Ranitidine reduces the amount of acid your stomach makes. It's used for indigestion and heartburn and acid reflux. It is also used for gastro-oesophageal reflux disease (GORD) - this is when you keep getting acid reflux. Ranitidine is also taken to prevent and treat stomach ulcers.



20x10 Tablets (Aluminium Strip)
 M.R.P. 41/-Per 10 Tablets

Raniday-DM

Composition

Each film coated tablet contains :
 Ranitidine Hydrochloride IP
 Eq. to Ranitidine 150mg.
 Domperidone IP 10mg.
 Colour : Sunset Yellow

Indication : Domperidone+Ranitidine belongs to a group of medicines called gastrointestinal agents used to treat gastro-oesophageal reflux disease (GERD) and peptic ulcers. Gastroesophageal reflux disease (GERD) occurs when stomach acid frequently flows back into the food pipe (oesophagus).



20x10 Tablets
 M.R.P. 20/-Per 10 Tablets

Rasgocin-1

Composition

Each uncoated tablet contains :
 Rasagiline Masylate
 Eq. to Rasagiline 1mg.
 Excipients q.s.

Indication : Rasagiline is used alone or in combination with another medication to treat the symptoms of Parkinson's disease (a slowly progressing disease of the nervous system causing a fixed face without expression, tremor at rest, slowing of movements, walking with shuffling steps, stooped posture and muscle weakness).



10x10 Tablets
 M.R.P. /-Per 10 Tablets

Rasgocin-2

Composition

Each uncoated tablet contains :
 Rasagiline Masylate
 Eq. to Rasagiline 2mg.
 Excipients q.s.

Indication : Rasagiline is used alone or in combination with another medication to treat the symptoms of Parkinson's disease (a slowly progressing disease of the nervous system causing a fixed face without expression, tremor at rest, slowing of movements, walking with shuffling steps, stooped posture and muscle weakness).



10x10 Tablets

M.R.P. /-Per 10 Tablets

Risperdone-4

Composition :

Each film coated tablet contains:
 Risperidone BP 4mg.
 Excipients q.s.
 Colour : Brillinat Blue & Titanium Dioxide IP

Indication : Risperidone is used to treat certain mental/mood disorders (such as schizophrenia, bipolar disorder, irritability associated with autistic disorder). This medication can help you to think clearly and take part in everyday life. Risperidone belongs to a class of drugs called atypical antipsychotics



10X10 Tablets (Blister)

M.R.P. 112.20/-Per 10 Tablets

Rivaxab-15

Composition :

Each film coated tablet contains :
 Rivaroxaban Micronised 15 mg
 Excipients q.s.
 Colour : Ferric Oxide Red & Titanium Dioxide

Indication : Rivaroxaban is used to treat deep vein thrombosis (DVT; a blood clot, usually in the leg) and pulmonary embolism (PE; a blood clot in the lung). Rivaroxaban may be continued to prevent DVT and/or PE from happening again after initial treatment is completed.



10x2x14 Tablets (Blister)

M.R.P. 595/-Per 14 Tablets

Risperdone-2

Composition :

Each film coated tablet contains:
 Risperidone BP 2mg.
 Excipients q.s.
 Colour : Sunset Yellow FCF

Indication : Risperidone is used to treat certain mental/mood disorders (such as schizophrenia, bipolar disorder, irritability associated with autistic disorder). This medication can help you to think clearly and take part in everyday life. Risperidone belongs to a class of drugs called atypical antipsychotics



10X10 Tablets (Blister)

M.R.P. 50.01/-Per 10 Tablets

Rivaxab-10

Composition :

Each film coated tablet contains :
 Rivaroxaban Micronised 10 mg
 Excipients q.s.
 Colour : Ferric Oxide Red &

Indication : Rivaroxaban is used to treat deep vein thrombosis (DVT; a blood clot, usually in the leg) and pulmonary embolism (PE; a blood clot in the lung). Rivaroxaban may be continued to prevent DVT and/or PE from happening again after initial treatment is completed.



10x2x14 Tablets (Blister)

M.R.P. 520/-Per 14 Tablets

Rivaxab-20

Composition :

Each film coated tablet contains :
 Rivaroxaban Micronised 20 mg
 Excipients q.s.
 Colour : Ferric Oxide Red & Titanium Dioxide

Indication : Rivaroxaban is used to treat deep vein thrombosis (DVT; a blood clot, usually in the leg) and pulmonary embolism (PE; a blood clot in the lung). Rivaroxaban may be continued to prevent DVT and/or PE from happening again after initial treatment is completed.



10x2x14 Tablets (Blister)

M.R.P. 700/-Per 14 Tablets

Ronirol-0.5

Composition :

Each film coated tablet contains :
 Ropinirole Hydrochloride USP 0.5 mg.
 Eq. to Ropinirole
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Ropinirole is also used to treat restless legs syndrome (RLS). It may improve your sleep by decreasing the urge to move your legs and decreasing uncomfortable/ unpleasant feelings in the legs. This medication works by helping to restore the balance of a certain natural substance (dopamine) in the brain



10X10 Tablets (Aluminium)
 M.R.P. 74/-Per 10 Tablets

Rosudoz-20

Composition :

Each film coated tablet contains :
 Rosuvastatin Calcium IP 20 mg.
 Eq. to Rosuvastatin
 Excipients q.s.
 Colours : Red oxide of Iron & Titanium Dioxide I

Indication : Rosuvastatin is used along with a proper diet to help lower "bad" cholesterol and fats (such as LDL, triglycerides) and raise "good" cholesterol (HDL) in the blood. It belongs to a group of drugs known as "statins." It works by reducing the amount of cholesterol made by the liver.



10X10 Tablets (Alu Alu)
 M.R.P. 243.25/-Per 10 Tablets

Rosudoz-F

Composition :

Each film coated tablet contains :
 Rosuvastatin Calcium IP 10 mg.
 Eq. to Rosuvastatin
 Fenofibrate IP 160mg
 Excipients q.s.

Indication : Fenofibrate and Rosuvastatin. Fenofibrate works by lowering the levels of triglycerides in your blood, while Rosuvastatin prevents your body from making "bad" cholesterol (LDL). Both raise the levels of "good" cholesterol (HDL).



10X10 Tablets (Alu Alu)
 M.R.P.190/-Per 10 Tablets

Rosudoz-10

Indication :

Each film coated tablet contains :
 Rosuvastatin Calcium IP 10 mg.
 Eq. to Rosuvastatin
 Excipients q.s.
 Colours : Red oxide of Iron & Titanium Dioxide IP

Indication : Rosuvastatin is used along with a proper diet to help lower "bad" cholesterol and fats (such as LDL, triglycerides) and raise "good" cholesterol (HDL) in the blood. It belongs to a group of drugs known as "statins." It works by reducing the amount of cholesterol made by the liver.



10X10 Tablets (Alu Alu)
 M.R.P. 133/-Per 10 Tablets

Rosudoz-40

Composition :

Each film coated tablet contains :
 Rosuvastatin Calcium IP 40 mg.
 Eq. to Rosuvastatin
 Excipients q.s.
 Colours : Red oxide of Iron & Titanium Dioxide IP

Indication : Rosuvastatin is used along with a proper diet to help lower "bad" cholesterol and fats (such as LDL, triglycerides) and raise "good" cholesterol (HDL) in the blood. It belongs to a group of drugs known as "statins." It works by reducing the amount of cholesterol made by the liver.



10X10 Tablets (Alu Alu)
 M.R.P. 362/-Per 10 Tablets

Runflow-500

Composition :

Each film coated tablet extended release tablet contains :
 Ranolazine 500mg.
 Excipients q.s.

Indication : Ranolazine is used alone or with other medications to treat chronic angina (ongoing chest pain or pressure that is felt when the heart does not get enough oxygen). Ranolazine is in a class of medications



10X10 Tablets (Alu Alu)
 M.R.P. 150/-Per 10 Tablets

Ryfxacare-200

Composition :

Each film coated tablet contains :
Rifaximin PH EUR B.P. 200 mg.
Excipients q.s.

Indication : Rifaximin, sold under the brand name Xifaxan among others, is an antibiotic medication used to treat travelers' diarrhea, irritable bowel syndrome, and hepatic encephalopathy. It has poor absorption when taken by mouth. It is based on rifamycin.



10X10 Tablets (Alu Alu)
M.R.P. 299/-Per 10 Tablets

Ryfxacare-550

Composition :

Each film coated tablet contains :
Rifaximin BP.....550 mg.
Excipients.....q.s
Colours : Lake of Sunset Yellow & Titanium Dioxide IP

Indication : Rifaximin, sold under the brand name Xifaxan among others, is an antibiotic medication used to treat travelers' diarrhea, irritable bowel syndrome, and hepatic encephalopathy. It has poor absorption when taken by mouth. It is based on rifamycin.



10X10 Tablets (Alu Alu)
M.R.P. 599/-Per 10 Tablets

Salfaus-500

Composition :

Each delayed release tablet contains :
Sulfasalazine USP 500mg.
Excipients q.s.
Colour : Ferric oxide Yellow

Indication : Sulfasalazine is used to treat bowel inflammation, diarrhea (stool frequency), rectal bleeding, and abdominal pain in patients with ulcerative colitis, a condition in which the bowel is inflamed.



10x15 Tablets (Blister)
M.R.P. 70.72/-Per 15 Tablets

Ryfxacare-400

Composition :

Each film coated tablet contains :
Rifaximin PH EUR B.P. 400 mg.
Excipients q.s.

Indication : Rifaximin, sold under the brand name Xifaxan among others, is an antibiotic medication used to treat travelers' diarrhea, irritable bowel syndrome, and hepatic encephalopathy. It has poor absorption when taken by mouth. It is based on rifamycin.



10X1x10 Tablets (Alu Alu)
M.R.P. 499/-Per 10 Tablets

Re-Caution CD3

Composition :

Each film coated tablet contains :
Methycobalamin IP 1500 mcg
Alpha Lipoic Acid IP 100 mg
Benfotiamine 150 mg
Inositol IP 100 mg
Chromium Picolinate IP 200 mcg
Pyridoxine Hydrochloride IP 3 mg
Folic Acid IP 1.5 mg
Vitamin D3 IP 1000 IU
Calcium Carbonate IP 500 mg
Excipients q.s.
Appropriate overages added to compensate the loss on Storage.



10x1x10 Tablets (Alu Alu)
M.R.P. 320/-Per 10 Tablets

Salfaus-1000

Composition :

Each delayed release tablet contains :
Sulfasalazine USP 1000mg.
Excipients q.s.
Colour : Ferric oxide Yellow

Indication : Sulfasalazine is used to treat bowel inflammation, diarrhea (stool frequency), rectal bleeding, and abdominal pain in patients with ulcerative colitis, a condition in which the bowel is inflamed.



10X10 Tablets (Blister)
M.R.P. 120/-Per 10 Tablets

Sertraline-50

Composition :

Each film coated tablet contains:
Sertraline Hydrochloride IP 50mg.
Eq. to Sertraline 50mg.
Excipients q.s.



Indication : Sertraline is used to treat depression, panic attacks, obsessive compulsive disorder, post-traumatic stress disorder, social anxiety disorder (social phobia), and a severe form of premenstrual syndrome (premenstrual dysphoric disorder).

10X10 Tablets (Blister)

M.R.P. 75.65/-Per 10 Tablets

Sevelamer-400

Composition :

Each film coated tablet contains :
Sevelamer Carbonate 400mg.
Excipients q.s.
Colour : Titanium Dioxide IP



Indication : Sevelamer is used to lower high blood phosphorus (phosphate) levels in patients who are on dialysis due to severe kidney disease. Dialysis removes some phosphate from your blood, but it is difficult to remove enough to keep your phosphate levels balanced.

10X10 Tablets (Alu Alu)

M.R.P. 299/-Per 10 Tablets

Sevelamer-800

Composition :

Each film coated tablet contains :
Sevelamer Carbonate 800mg.
Excipients q.s.
Colour : Titanium Dioxide IP



Indication : Sevelamer is used to lower high blood phosphorus (phosphate) levels in patients who are on dialysis due to severe kidney disease. Dialysis removes some phosphate from your blood, but it is difficult to remove enough to keep your phosphate levels balanced.

10X10 Tablets (Alu Alu)

M.R.P. 410/-Per 10 Tablets

Simnol-25

Composition :

Each uncoated tablet contains:
Atenolol IP 25 mg.
Excipients q.s.



Indication : Atenolol belongs to a group of medicines called beta blockers. It's used to treat high blood pressure and irregular heartbeats (arrhythmia). It can also be used to prevent chest pain caused by angina. If you have high blood pressure, taking atenolol helps prevent future heart disease, heart attacks and strokes.

20X14 Tablets (Blister)

M.R.P. 23.40/-Per 14 Tablets

Simnol-50

Composition :

Each uncoated tablet contains:
Atenolol IP 50 mg.
Excipients q.s.



Indication : Atenolol belongs to a group of medicines called beta blockers. It's used to treat high blood pressure and irregular heartbeats (arrhythmia). It can also be used to prevent chest pain caused by angina. If you have high blood pressure, taking atenolol helps prevent future heart disease, heart attacks and strokes.

20x14 Tablets (Blister)

M.R.P. 30.98/-Per 14 Tablets

Sinzin-25

Composition :

Each film coated tablet contains :
Cinnarizine IP 25mg.
Excipients q.s.
Colour : Titanium Dioxide IP



Indication : It's used for: travel sickness (also called motion sickness) dizziness and sickness caused by inner ear problems (such as vertigo, tinnitus and Ménière's disease)

4X5X10 Tablets (Blister)

M.R.P. 76/-Per 10 Tablets

Sitadose-25

Composition :

Each film coated tablet contains:
 Sitagliptin Phosphate Monohydrate IP
 Eq. to Sitagliptin 25 mg
 Colours : Red Oxide of Iron, Yellow
 Oxide of Iron & Titanium Dioxide IP



Indication : Sitagliptin is used along with diet and exercise and sometimes with other medications to lower blood sugar levels in adults with type 2 diabetes (condition in which blood sugar is too high because the body does not produce or use insulin normally).

10X7 Tablets (Blister)

M.R.P. 255/-Per 7 Tablets

Sitadose-100

Composition :

Each film coated tablet contains:
 Sitagliptin Phosphate Monohydrate IP
 Eq. to Sitagliptin 100 mg
 Colours : Red Oxide of Iron, Yellow
 Oxide of Iron & Titanium Dioxide IP



Indication : Sitagliptin is used along with diet and exercise and sometimes with other medications to lower blood sugar levels in adults with type 2 diabetes (condition in which blood sugar is too high because the body does not produce or use insulin normally).

10X7 Tablets (Blister)

M.R.P. 315/-Per 7 Tablets

Sitadose-M 1000

Composition :

Each film coated tablet contains:
 Sitagliptin Phosphate
 Monohydrate IP
 Eq. to Sitagliptin 50 mg
 Metformin
 Hydrochloride IP 1000 mg
 Excipients q.s.
 Colour : Red Oxide of Iron, Black
 Oxide of Iron & Titanium Dioxide IP



Indication : Sitagliptin is used along with diet and exercise and sometimes with other medications to lower blood sugar levels in adults with type 2 diabetes (condition in which blood sugar is too high because the body does not produce or use insulin normally).

10x15 Tablets (Blister)

M.R.P. 336/-Per 15 Tablets

Sitadose-50

Composition :

Each film coated tablet contains:
 Sitagliptin Phosphate Monohydrate IP
 Eq. to Sitagliptin 50 mg
 Colours : Red Oxide of Iron, Yellow
 Oxide of Iron & Titanium Dioxide IP



Indication : Sitagliptin is used along with diet and exercise and sometimes with other medications to lower blood sugar levels in adults with type 2 diabetes (condition in which blood sugar is too high because the body does not produce or use insulin normally).

10x7 Tablets (Blister)

M.R.P. 290/-Per 7 Tablets

Sitadose-M 500

Composition :

Each film coated tablet contains:
 Sitagliptin Phosphate
 Monohydrate IP
 Eq. to Sitagliptin 50 mg
 Metformin
 Hydrochloride IP 500 mg
 Excipients q.s.
 Colour : Red Oxide of Iron, Black
 Oxide of Iron & Titanium Dioxide IP



Indication : Sitagliptin is used along with diet and exercise and sometimes with other medications to lower blood sugar levels in adults with type 2 diabetes (condition in which blood sugar is too high because the body does not produce or use insulin normally).

10x15 Tablets (Blister)

M.R.P. 308/-Per 15 Tablets

Sitadose-D 100/10

Composition

Each film coated tablet contains :
 Sitagliptin Phosphate Monohydrate IP
 Eqv. to Sitagliptin 100mg.
 Dapagliflozin Propanediol Monohydrate
 Eqv. to Dapagliflozin 10mg.
 Excipients q.s.
 Colour : Yellow oxide of Iron &
 Titanium Dioxide IP



Indication : Sitagliptin is used along with diet and exercise and sometimes with other medications to lower blood sugar levels in adults with type 2 diabetes (condition in which blood sugar is too high because the body does not produce or use insulin normally).

10x10 Tablets (Alu Alu)

M.R.P. 310/-Per 10 Tablets

Sitadose-DM

Composition

Each film coated bilayered tablet contains :
 Dapagliflozin Propanediol Monohydrate eq. to Dapagliflozin 10mg
 Sitagliptin Phosphate Monohydrate IP eq. to Sitagliptin 100mg.
 Metformin Hydrochloride IP 500mg. (As extended release)
 Excipients q.s.
 Colour : Yellow oxide of Iron USP-NF (in Dapagliflozin & sitagliptin Layer)

Indication : Sitagliptin is used along with diet and exercise and sometimes with other medications to lower blood sugar levels in adults with type 2 diabetes (condition in which blood sugar is too high because the body does not produce or use insulin normally).

10x10 Tablets (Alu Alu)
 M.R.P. 330/-Per 10 Tablets



Sodicare-500

Composition :

Each film coated tablet contains:
 Sodium Bicarbonate IP 500mg.
 Excipients q.s.
 Titanium Dioxide IP

Indication : Sodium bicarbonate , also known as baking soda, is used to relieve heartburn, sour stomach, or acid indigestion by neutralizing excess stomach acid. When used for this purpose, it is said to belong to the group of medicines called antacids. It may be used to treat the symptoms of stomach or duodenal ulcers.

10X10 Tablets (Blister)
 M.R.P. 32.90/-Per 10 Tablets



Sodicare-1000

Composition :

Each film coated tablet contains:
 Sodium Bicarbonate IP 1000mg.
 Excipients q.s.
 Titanium Dioxide IP

Indication : Sodium bicarbonate , also known as baking soda, is used to relieve heartburn, sour stomach, or acid indigestion by neutralizing excess stomach acid. When used for this purpose, it is said to belong to the group of medicines called antacids. It may be used to treat the symptoms of stomach or duodenal ulcers.

10X10 Tablets (Blister)
 M.R.P. 85/-Per 10 Tablets



Solifenacin-5

Composition :

Each film coated tablet contains:
 Solifenacin Succinate 5mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Solifenacin is used to treat an overactive bladder. By relaxing the muscles in the bladder, solifenacin improves your ability to control your urination. It helps to reduce leaking of urine, feelings of needing to urinate right away, and frequent trips to the bathroom..

10X10 Tablets (Blister)
 M.R.P. 294/-Per 10 Tablets



Solifenacin-10

Composition :

Each film coated tablet contains:
 Solifenacin Succinate 10mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Solifenacin is used to treat an overactive bladder. By relaxing the muscles in the bladder, solifenacin improves your ability to control your urination. It helps to reduce leaking of urine, feelings of needing to urinate right away, and frequent trips to the bathroom..

10X10 Tablets (Blister)
 M.R.P. 370/-Per 10 Tablets



Spiroton-25

Composition :

Each uncoated tablet contains :
 Spironolactone IP 25mg.
 Excipients q.s.

Indication : Spironolactone is used to treat certain patients with hyperaldosteronism (the body produces too much aldosterone, a naturally occurring hormone); low potassium levels; heart failure; and in patients with edema (fluid retention) caused by various conditions, including liver, or kidney disease.

10x10 Tablets (Blister)
 M.R.P. 23.80/-Per 10 Tablets



Spironot-50

Composition :

Each uncoated tablet contains :
 Spironolactone IP 50mg.
 Excipients q.s

Indication : Spironolactone is used to treat certain patients with hyperaldosteronism (the body produces too much aldosterone, a naturally occurring hormone); low potassium levels; heart failure; and in patients with edema (fluid retention) caused by various conditions,



10x10 Tablets (Blister)
 M.R.P. 43.60/-Per 10 Tablets

Spironot-F

Composition :

Each film coated tablet contains :
 Furosemide IP 20mg.
 Spironolactone IP 50mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : furosemide and spironolactone are used to reduce edema (fluid accumulation) caused by heart, liver or kidney problems, or high blood pressure (hypertension). Lasix may also be used with antihypertensive drugs to control high blood pressure (hypertension).



10X10 Tablets (Blister)
 M.R.P. 199/-Per 10 Tablets

Sugarmet-500

Composition :

Each film coated tablet contains
 Metformin Hydrochloride IP 500 mg.
 Excipients q.s.
 Colour: Titanium Dioxide IP

Indication : Metformin helps to control the amount of glucose (sugar) in your blood. It decreases the amount of glucose you absorb from your food and the amount of glucose made by your liver. Metformin also increases your body's response to insulin, a natural substance that controls the amount of glucose in the blood, including liver, or kidney disease.



20x20 Tablets (Blister)
 M.R.P. 39.60/-Per 20 Tablets

Sugarmet-SR 500

Composition :

Each uncoated Sustained Release tablet contains :
 Metformin Hydrochloride IP 500mg
 Excipients q.s.

Indication : Metformin is used with a proper diet and exercise program and possibly with other medications to control high blood sugar. It is used in patients with type 2 diabetes. Controlling high blood sugar helps prevent kidney damage, blindness, nerve problems, loss of limbs, and sexual function problems.



20x10 Tablets (Blister)
 M.R.P. 23.80/-Per 10 Tablets

Sugarmet-850

Composition :

Each uncoated tablet contains :
 Metformin Hydrochloride IP 850mg.
 Excipients q.s.

Indication : Metformin is used with a proper diet and exercise program and possibly with other medications to control high blood sugar. It is used in patients with type 2 diabetes. Controlling high blood sugar helps prevent kidney damage, blindness, nerve problems, loss of limbs, and sexual function problems.



10x10 Tablets (Blister)
 M.R.P. 23/-Per 10 Tablets

Sugarmet-SR 1000

Composition :

Each film coated sustained release tablet contains :
 Metformin Hydrochloride IP 1000mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Metformin is used with a proper diet and exercise program and possibly with other medications to control high blood sugar. It is used in patients with type 2 diabetes. Controlling high blood sugar helps prevent kidney damage, blindness, nerve problems, loss of limbs, and sexual function problems.



10x10 Tablets (Blister)
 M.R.P. 45.24/-Per 10 Tablets

Sumeday-50

Composition :

Each film coated tablet contains :
Sumatriptan Succinate IP
Eq. to Sumatriptan 50mg.
Excipients q.s.

Indication : Sumatriptan is a medicine to treat migraines and cluster headaches. It does not prevent these conditions. It belongs to a group of medicines called triptans, or serotonin (5-HT1) agonists. You take sumatriptan once a migraine or cluster headache has started.



10x1x10 Tablets (Alu Alu)
M.R.P. 623.60/-Per 10 Tablets

Sultaday-375

Composition :

Each film coated tablet contains :
Sultamicillin Tosilate Dihydrate B.P. 375mg.
(Equivalent to Sultabactam 147mg.
and Ampicillin 220mg)
Excipients q.s.
Colour : Titanium Dioxide IP

Indication : It is a 'broad-spectrum antibiotic' that treats various bacterial infections of the middle ear and sinus infections, throat or lung respiratory tract infections, urinary tract infections, skin infections, soft tissue infections, joint and bone infections, and post-operative infections.



10x1x10 Tablets (Alu Alu)
M.R.P. 432/-Per 10 Tablets

Sirlofast-1

Composition :

Each film coated tablet contains :
Sirolimus 1mg
Excipients q.s.
Colour : Titanium Dioxide IP

Indication : Sirolimus is used in combination with other medications to prevent rejection of kidney transplants. Sirolimus is in a class of medications called immunosuppressants. It works by suppressing the body's immune system.




10x1x10 Tablets (Alu Alu)
M.R.P. 1490/-Per 10 Tablets

Tammislin

Composition :

Each film coated tablet contains :
Tamsulosin HCl IP 0.4mg.
(As Modified Release)
Excipients q.s.
Colour : Titanium Dioxide IP
Red Oxide of Iron

Indication : Tamsulosin belongs to a group of medicines called alpha-blockers. It's used to treat men with symptoms of an enlarged prostate (benign prostate enlargement). It's also occasionally taken to treat kidney stones.



20x10 Tablets (Alu Alu)
M.R.P. 102/-Per 10 Tablets

Tammislin-D

Composition :

Each film coated tablet contains :
Tamsulosin HCl IP 0.4mg.
Dutasteride IP 0.5mg.
Excipients q.s.
Colour : Titanium Dioxide IP

Indication : This product is used in men to treat the symptoms of an enlarged prostate (benign prostatic hyperplasia-BPH). Dutasteride works by reducing the size of the enlarged prostate. Tamsulosin is known as an alpha-blocker and works by relaxing muscles in the bladder and prostate.



10x10 Tablets (Alu Alu)
M.R.P. 195/-Per 10 Tablets

Tamslin-F

Composition :

Each Film coated tablet contains :
Tamsulosin Hydrochloride IP 0.4mg.
Finasteride IP 5 mg.
Excipients q.s.
Colour : Titanium Dioxide IP

Indication : Tamsulosin belongs to a group of medicines called alpha-blockers. It's used to treat men with symptoms of an enlarged prostate (benign prostate enlargement). It's also occasionally taken to treat kidney stones.



10x10 Tablets (Blister)
M.R.P. 180/-Per 10 Tablets

Telmidoz[®]-CH 40/6.25

Composition :

Each film coated tablet contains:
 Telmisartan IP 40 mg.
 Chlorthalidone IP 6.25mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Telmisartan+Chlorthalidone is used in the treatment of Hypertension (high blood pressure). Telmisartan + Chlorthalidone is a combination of two medicines: Telmisartan and chlorthalidone, which lowers blood pressure effectively. Telmisartan is an angiotensin receptor blocker (ARB).



10X10 Tablets (Alu Alu)
 M.R.P. 70/-Per 10 Tablets

Telmidoz[®]-CH 40/12.5

Composition :

Each film coated tablet contains:
 Telmisartan IP 40 mg.
 Chlorthalidone IP 12.5mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication: Telmisartan+Chlorthalidone is used in the treatment of Hypertension (high blood pressure). Telmisartan + Chlorthalidone is a combination of two medicines: Telmisartan and chlorthalidone, which lowers blood pressure effectively. Telmisartan is an angiotensin receptor blocker (ARB).



10X10 Tablets (Alu Alu)
 M.R.P. 95/-Per 10 Tablets

Telmidoz[®]-20

Composition :

Each film coated Tablet contains
 Telmisartan IP 20 mg.
 Excipients q.s.
 Colour: Titanum Dioxide IP

Indication : Telmisartan is used alone or in combination with other medications to treat high blood pressure. Telmisartan is also used to decrease the chance of heart attack, stroke, or death in people 55 years of age or older who are at high risk for cardiovascular disease.



10X10 Tablets (Alu Alu)
 M.R.P.43.34/-Per 10 Tablets

Telmidoz[®]-40

Composition :

Each film coated Tablet contains
 Telmisartan IP 40 mg.
 Excipients q.s.
 Colour: Titanum Dioxide IP

Indication : Telmisartan is used alone or in combination with other medications to treat high blood pressure. Telmisartan is also used to decrease the chance of heart attack, stroke, or death in people 55 years of age or older who are at high risk for cardiovascular disease.



10x10 Tablets (Alu Alu)
 M.R.P.75.70/-Per 10 Tablets

Telmidoz[®]-80

Composition :

Each film coated Tablet contains
 Telmisartan IP 80 mg.
 Excipients q.s.
 Colour: Titanum Dioxide IP

Indication : Telmisartan is used alone or in combination with other medications to treat high blood pressure. Telmisartan is also used to decrease the chance of heart attack, stroke, or death in people 55 years of age or older who are at high risk for cardiovascular disease.



10X10 Tablets (Alu Alu)
 M.R.P.116.48/-Per 10 Tablets

Telmidoz[®]-AH

Composition :

Each film coated tablet contains :
 Telmisartan IP 40mg.
 Hydrochlorothiazide IP 12.5mg.
 Amlodipine Besylate IP
 eq. to Amlodipine 5mg.
 Excipients q.s.

Indication: Telmisartan/Hydrochlorothiazide is a combination of an angiotensin II receptor antagonist, telmisartan, and a thiazide diuretic, hydrochlorothiazide. The combination of these ingredients has an additive antihypertensive effect, reducing blood pressure to a greater degree than either component alone.



10X10 Tablets (Alu Alu)
 M.R.P. 79/-Per 10 Tablets

TABLETS

Telmidoz[®]-AZ 40/8

Composition :

Each uncoated tablet contains :

Azelnidipine	IP	8mg.
Telmisartan	IP	40mg.
Excipients		q.s.

Indication : Azelnidipine+Telmisartan Belongs To The Class Of Anti-hypertensive Drugs Primarily Used To Treat Hypertension (high Blood Pressure). Hypertension (high Blood Pressure) Is A Medical Condition In Which The Blood Pressure Is Elevated Persistently In The Arteries.

10X10 Tablets (Alu Alu)

M.R.P. 127/-Per 10 Tablets



Telmidoz[®]-AM

Composition

Each film coated Tablet contains

Telmisartan	IP	40mg
Amlodipine Besylate	IP	5mg
Excipients		q.s.
Colour:		Titanium Dioxide IP

Telmisartan and amlodipine is a combination of medicines that may be used alone or with other medicines to treat high blood pressure (hypertension). High blood pressure adds to the workload of the heart and arteries. If it continues for a long time, the heart and arteries may not function properly.

10X10 Tablets (Alu Alu)

M.R.P. 80.80/-Per 10 Tablets



Telmidoz[®]-CC

Composition :

Each film coated Tablet contains

Telmisartan	IP	40 mg.
Chlorthalidone	IP	12.5 mg.
Cilnidipine	IP	10 mg.
Excipients		q.s.

Indication : Telmisartan is an angiotensin receptor blocker (ARB) and Cilnidipine is a calcium channel blocker (CCB). They work by widening and relaxing the blood vessels, thereby lowering blood pressure. Chlorthalidone is a diuretic which removes extra water and certain electrolytes from the body through

10X10 Tablets (Alu Alu)

M.R.P. 150/-Per 10 Tablets



Telmidoz[®]-C

Composition :

Each film coated Tablet contains

Telmisartan	IP	40mg
Cilnidipine	IP	10mg
Excipients		q.s.
Colour:		Red oxide of Iron

Indication : Cilnidipine + Telmisartan is a combination of two medicines: Cilnidipine and Telmisartan, which lowers blood pressure effectively. Cilnidipine is a calcium channel blocker (CCB) and Telmisartan is an angiotensin receptor blocker (ARB).

10X10 Tablets (Alu Alu)

M.R.P. 120/-Per 10 Tablets



Telmidoz[®]-H

Composition :

Each uncoated bilayered tablet contains

Telmisartan	IP	40 mg.
Hydrochlorothiazide	IP	12.5 mg.
Excipients		q.s.
Colour :		Red Oxide of Iron

Indication Telmisartan/Hydrochlorothiazide is a combination of an angiotensin II receptor antagonist, telmisartan, and a thiazide diuretic, hydrochlorothiazide. The combination of these ingredients has an additive antihypertensive effect, reducing blood pressure to a greater degree than either component alone.

10X10 Tablets (Alu Alu)

M.R.P. 84/-Per 10 Tablets



Telmidoz[®]-M 25

Composition :

Each uncoated tablet contains :

Telmisartan IP	40mg.
(In Immediate Release Form)	
Metoprolol Succinate IP	23.75 mg.
Eq. to Metoprolol Tartrate	25mg.
(In Extended Release Form)	
Excipients	q.s.
Colour :	Sunset Yellow

Indication : Telmisartan is an angiotensin receptor blocker (ARB). It works by blocking the chemical angiotensin, thereby relaxing blood vessels. Metoprolol Succinate is a beta blocker which works specifically on the heart to reduce heart rate and lower blood pressure.

10X10 Tablets (Alu Alu)

M.R.P. 130/-Per 10 Tablets



Telmidoz[®]-M 50

Composition :

Each uncoated tablet contains :
 Telmisartan IP 40mg.
 (in Immediate Release Form)
 Metoprolol Succinate IP 47.5 mg.
 Eq. to Metoprolol Tartrate 50mg.
 (In Extended Release Form)
 Excipients q.s.
 Colour : Sunset Yellow

Indication : Telmisartan is an angiotensin receptor blocker (ARB). It works by blocking the chemical angiotensin, thereby relaxing blood vessels. Metoprolol Succinate is a beta blocker which works specifically on the heart to reduce heart rate and lower blood pressure.

10X10 Tablets (Alu Alu)
 M.R.P. 160.96/-Per 10 Tablets



Tenedose[®]-20

Composition :

Each film coated tablet contains
 Teneligliptin
 Hydrobromide Hydrate IP
 Equivalent to Teneligliptin 20 mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : teneligliptin has been commercially sold in Japan and has been used for the treatment of type 2 diabetes mellitus when patients do not show sufficient improvement after diet control and exercise or a combination of diet control, exercise, and sulfonylurea- or thiazolidine-class drugs.

10X10 Tablets (Alu Alu)
 M.R.P. 105/-Per 10 Tablets



Tenedose[®]-M

Composition :

Each uncoated bilayered tablet contains :
 Teneligliptin
 Hydrobromide Hydrate IP
 eq. to Teneligliptin 20 mg.
 Metformin Hydrochloride IP 500 mg.
 (In extended release form)
 Excipients q.s.
 Colour : Yellow Oxide of Iron.

Indication : By increasing and prolonging active incretin levels, teneligliptin increases insulin release and decreases glucagon levels in the circulation in a glucose-dependent manner. Metformin improves glucose tolerance in patients with type-2 diabetes (NIDDM), lowering both basal and postprandial plasma glucose.

10X10 Tablets (Alu Alu)
 M.R.P. 107.70/-Per 10 Tablets



Tenedose[®]-M Forte

Composition :

Each uncoated bilayered tablet contains :
 Teneligliptin
 Hydrobromide Hydrate IP
 eq. to Teneligliptin 20 mg.
 Metformin Hydrochloride IP 1000 mg.
 (In extended release form)
 Excipients q.s.
 Colour : Ferric Oxide Yellow USP NF

Indication : By increasing and prolonging active incretin levels, teneligliptin increases insulin release and decreases glucagon levels in the circulation in a glucose-dependent manner. Metformin improves glucose tolerance in patients with type-2 diabetes (NIDDM), lowering both basal and postprandial plasma glucose.

10X10 Tablets (Alu Alu)
 M.R.P. 122.30/-Per 10 Tablets



Terbicare

Composition :

Each uncoated tablet contains :
 Terbinafine Hydrochloride IP
 eq. to Terbinafine 250 mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Terbinafine is an antifungal medicine. It is used to treat skin infections caused by a fungus (yeast), including: athlete's foot, fungal nail infections.

20x1x7 Tablets (Blister)
 M.R.P. 210/-Per 7 Tablets



Terbicare-DS

Composition :

Each uncoated tablet contains :
 Terbinafine Hydrochloride IP
 eq. to Terbinafine 500 mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Terbinafine is an antifungal medicine. It is used to treat skin infections caused by a fungus (yeast), including: athlete's foot, fungal nail infections.

10x1x7 Tablets (Blister)
 M.R.P. 369/-Per 7 Tablets





TABLETS

Timeovir-300

Composition :

Each film coated tablet contains :
 Tenofovir Disoproxil Fumarate IP 300mg.
 Excipients q.s.

Indication : Tenofovir is used with other HIV medications to help control HIV infection. It helps to decrease the amount of HIV in your body so your immune system can work better. This lowers your chance of getting HIV complications (such as new infections, cancer) and improves your quality of life.



1x30 Tablets
M.R.P. 1282/-Per 30 Tablets

Tofacare-5

Composition :

Each film coated tablet contains :
 Tofacitinib Citrate eq. to Tofacitinib 5mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Tofacitinib is used alone or with other medications to treat rheumatoid arthritis (condition in which the body attacks its own joints causing pain, swelling, and loss of function) in adults who are unable to take or did not respond to one or more tumor necrosis factor (TNF) inhibitor medication(s).



6x10 Tablets (Alu Alu)
M.R.P. 401/-Per 10 Tablets

Tofpam-50

Composition

Each uncoated tablet contains :
 Tofisopam JP 50mg.
 Excipients q.s.

Indication : Tofisopam contains an 'anti-depression medication', Tofisopam, primarily used to treat depression and associated anxiety with depression. It works by increasing the action of a chemical messenger known as GABA (gamma-amino-butyric-acid), which suppresses the excessive and abnormal activity of nerve cells in the brain.



10x10 Tablets (Alu Alu)
M.R.P. 210/-Per 10 Tablets

Tikaglor-90

Composition :

Each film coated tablet contains :
 Ticagrelor IP 90mg.
 Excipients q.s.
 Colour Yellow oxide of Iron & Iron Titanium Dioxide IP

Indication : Ticagrelor is used to prevent a serious or life-threatening heart attack or stroke, or death in people who have had a heart attack or who have acute coronary syndrome (ACS; blockage of blood flow to the




10x14 Tablets (Blister)
M.R.P. 499/-Per 14 Tablets

Tofacare-11

Composition :

Each film coated tablet contains :
 Tofacitinib Citrate eq. to Tofacitinib 11mg.
 Excipients q.s.
 Colour : Red oxide of Iron & Black Oxide of Iron & Titanium Dioxide IP

Indication : Tofacitinib is used alone or with other medications to treat rheumatoid arthritis (condition in which the body attacks its own joints causing pain, swelling, and loss of function) in adults who are unable to take or did not respond to one or more tumor necrosis factor (TNF) inhibitor medication(s).



6x10 Tablets (Alu Alu)
M.R.P. 990/-Per 10 Tablets

Tofpam-100

Composition

Each uncoated tablet contains :
 Tofisopam JP 100mg.
 Excipients q.s.

Indication : Tofisopam contains an 'anti-depression medication', Tofisopam, primarily used to treat depression and associated anxiety with depression. It works by increasing the action of a chemical messenger known as GABA (gamma-amino-butyric-acid), which suppresses the excessive and abnormal activity of nerve cells in the brain.



10x10 Tablets (Alu Alu)
M.R.P. 360/-Per 10 Tablets

Tolvaptan

Composition :
Each uncoated tablet contains:
Tolvaptan IP 15mg.
Excipients q.s.

Indication : This medication is used to treat low levels of salt (sodium) in the blood, which can result from conditions such as heart failure and certain hormone imbalances. Tolvaptan belongs to a class of drugs known as vasopressin receptor antagonists.



10x1x4 Tablets (Alu Alu)
M.R.P. 574/-Per 4 Tablets

Topirmat-25

Composition :
Each film coated Tablet contains :
Topiramate IP 25mg.
Excipients q.s.
Colour : Indigo Carmine & Titanium Dioxide IP

Indication : Topiramate is a medicine used to treat epilepsy. These are also called anti-epileptic medicines. It can also be taken to prevent migraine.



10X10 Tablets (Blister)
M.R.P. 120/-Per 10 Tablets

Topirmat-100

Composition :
Each film coated Tablet contains :
Topiramate IP 100mg.
Excipients q.s.
Colour : Indigo Carmine & Titanium Dioxide IP

Indication : Topiramate is a medicine used to treat epilepsy. These are also called anti-epileptic medicines. It can also be taken to prevent migraine.



10X10 Tablets (Blister)
M.R.P. 155/-Per 10 Tablets

Tolvaptan-30

Composition :
Each uncoated tablet contains:
Tolvaptan IP 30mg.
Excipients q.s.

Indication : This medication is used to treat low levels of salt (sodium) in the blood, which can result from conditions such as heart failure and certain hormone imbalances. Tolvaptan belongs to a class of drugs known as vasopressin receptor antagonists.



10x1x4 Tablets (Alu Alu)
M.R.P. 760/-Per 4 Tablets

Topirmat-50

Composition :
Each film coated Tablet contains :
Topiramate IP 50mg.
Excipients q.s.
Colour : Indigo Carmine & Titanium Dioxide IP

Indication : Topiramate is a medicine used to treat epilepsy. These are also called anti-epileptic medicines. It can also be taken to prevent migraine.



10X10 Tablets (Blister)
M.R.P. 140/-Per 10 Tablets

Tormay-5

Composition :
Each uncoated tablet contains :
Torsemide IP 5mg.
Excipients q.s.

Indication : Torsemide is used alone or in combination with other medications to treat high blood pressure. Torsemide is used to treat edema (fluid retention; excess fluid held in body tissues) caused by various medical problems, including heart, kidney, or liver disease.



4x5x10 Tablets (Blister)
M.R.P. 36/-Per 10 Tablets



TABLETS

Tormay-10

Composition :

Each uncoated tablet contains :
Torsemide IP 10mg.
Excipients q.s.

Indication : Torsemide is used alone or in combination with other medications to treat high blood pressure. Torsemide is used to treat edema (fluid retention; excess fluid held in body tissues) caused by various medical problems, including heart, kidney, or liver disease.



4x5x10 Tablets (Blister)

M.R.P. 69/-Per 10 Tablets

Tormay-20

Composition :

Each uncoated tablet contains :
Torsemide IP 20mg.
Excipients q.s.

Indication : Torsemide is used alone or in combination with other medications to treat high blood pressure. Torsemide is used to treat edema (fluid retention; excess fluid held in body tissues) caused by various medical problems, including heart, kidney, or liver disease.



4x5x10 Tablets (Blister)

M.R.P. 89/-Per 10 Tablets

Tormay-40

Composition :

Each uncoated tablet contains :
Torsemide IP 40mg.
Excipients q.s.

Indication : Torsemide is used alone or in combination with other medications to treat high blood pressure. Torsemide is used to treat edema (fluid retention; excess fluid held in body tissues) caused by various medical problems, including heart, kidney, or liver disease.



10X10 Tablets (Alu Alu)

M.R.P. 140/-Per 10 Tablets

Tormay-100

Composition :

Each uncoated tablet contains :
Torsemide IP 100mg.
Excipients q.s.

Indication : Torsemide is used alone or in combination with other medications to treat high blood pressure. Torsemide is used to treat edema (fluid retention; excess fluid held in body tissues) caused by various medical problems, including heart, kidney, or liver disease.



10X10 Tablets (Blister)

M.R.P. 338/-Per 10 Tablets

Tormay-S

Composition :

Each uncoated tablet contains :
Torsemide IP 10mg.
Spironolactone IP 50mg.
Excipients q.s.

Indication : is used to reduce extra fluid in the body (edema) caused by conditions such as heart failure, liver disease, and kidney disease. This can lessen symptoms such as shortness of breath and swelling in your arms, legs, and abdomen. This drug is also used to treat high blood pressure.



10X10 Tablets (Blister)

M.R.P. 185/-Per 10 Tablets

Traxpain

Composition :

Each film coated tablet contains :
Tranexamic Acid IP 500 mg.
Excipients q.s.
Colour : Yellow oxide of Iron

Indication : Tranexamic acid (sometimes shortened to txa) is a medicine that controls bleeding. It helps your blood to clot and is used for nosebleeds and heavy periods. If you're having a tooth taken out, using tranexamic acid mouthwash can help stop bleeding.



10x10 Tablets (Alu Alu)

M.R.P. 194.60/-Per 10 Tablets

Thyrocut

Support Therapy
For Thyroid



Composition :
Each uncoated tablet contains :
Thyroxine Sodium I.P.
eq. to Anhydrous
Thyroxine Sodium 12.5mcg
Excipients q.s.


1x100 Tablets (Bottle)
M.R.P. 69.75/-Per 100 Tablets

Composition :
Each uncoated tablet contains :
Thyroxine Sodium I.P.
eq. to Anhydrous
Thyroxine Sodium 25mcg
Excipients q.s.




1x100 Tablets (Bottle)
M.R.P. 125/-Per 100 Tablets

Composition :
Each uncoated tablet contains :
Thyroxine Sodium I.P.
eq. to Anhydrous
Thyroxine Sodium 50mcg
Excipients q.s.

1x100 Tablets (Bottle)
M.R.P. 103.04/-Per 10 Tablets



Hypothyroidism | Normalizes Elevated Tsh Levels
Increased Metabolism | Growth And Development



Composition :
Each uncoated tablet contains :
Thyroxine Sodium I.P.
eq. to Anhydrous
Thyroxine Sodium 75mcg
Excipients q.s.

1x100 Tablets (Bottle)
M.R.P. 105.95/-Per 100 Tablets

Composition :
Each uncoated tablet contains :
Thyroxine Sodium I.P.
eq. to Anhydrous
Thyroxine Sodium 100mcg
Excipients q.s.

1x100 Tablets (Bottle)
M.R.P. 131/-Per 100 Tablets

Composition :
Each uncoated tablet contains :
Thyroxine Sodium I.P.
eq. to Anhydrous
Thyroxine Sodium 125mcg
Excipients q.s.

1x100 Tablets (Bottle)
M.R.P. 172.48/-Per 100 Tablets

Composition :
Each uncoated tablet contains :
Thyroxine Sodium I.P.
eq. to Anhydrous
Thyroxine Sodium 150mcg
Excipients q.s.

1x100 Tablets (Bottle)
M.R.P. 156/-Per 100 Tablets

Traxpain-MF

Composition :

Each film coated tablet contains :
 Tranexamic Acid I.P. 500 mg.
 Mefenamic Acid I.P. 250 mg.
 Excipients q.s.
 Colour : Titanium Dioxide I.P.

Indication : Tranexamic Acid+ Mefenamic Acid is used in the treatment of heavy menstrual bleeding and menstrual pain. Tranexamic Acid + Mefenamic Acid is a combination of two medicines: Tranexamic Acid and Mefenamic Acid. Tranexamic Acid is an anti-fibrinolytic.



10X10 Tablets (Alu Alu)
 M.R.P. 210/-Per 10 Tablets

Trendcare-Forte

Each uncoated bilayered tablet contains (approx):

		% RDA*
L-Arginine as sustained release form	500 mg	**
Chondroitin Sulfate Sodium	200 mg	**
Collagen Peptide Type 1	40 mg	**
Vitamin C	30 mg	88
Sodium Hyaluronate	30 mg	**

Excipients q.s.
 *Sodium Based on ICMR guidelines **RDA not established

Nutritional Information per serving / tablet (approx):	
Energy	3.40 kcal
Protein	0.51 g
Carbohydrate	0.31 g
Sugar	0 g
Fat	0 g

Figures based on calculated values.



10x1x15 Tablets (Aluminium)
 M.R.P. 460/-Per 15 Tablets

Trazodone-25

Composition

Each uncoated tablet contains :
 Trazodone Hydrochloride IP 25mg.
 Excipients q.s.
 Colour : Ponceau 4r

Indication : It's used to treat depression, anxiety, or a combination of depression and anxiety. Trazodone works by increasing your levels of serotonin and noradrenaline so you feel better. It can help if you're having problems like low mood, not sleeping (insomnia) and poor concentration.



10x10 Tablets (Blister)
 M.R.P. 42/-Per 10 Tablets



Composition

Each uncoated tablet contains :
 Trazodone Hydrochloride IP 50mg.
 Excipients q.s.
 Colour : Ponceau 4r

Indication : It's used to treat depression, anxiety, or a combination of depression and anxiety. Trazodone works by increasing your levels of serotonin and noradrenaline so you feel better. It can help if you're having problems like low mood, not sleeping (insomnia) and poor concentration.

10x10 Tablets (Blister)
 M.R.P. 87/-Per 10 Tablets

Trimzed-35

Composition :

Each film coated modified release tablet contains:
 Trimetazidine Dihydrochloride IP 35mg.
 Excipients q.s.
 Colour : Red oxide of Iron & Titanium Dioxide IP

Indication : Trimetazidine is an anti-ischemic (anti-anginal) metabolic agent, which improves myocardial glucose utilization through inhibition of fatty acid metabolism, also known as fatty acid oxidation inhibitor.



10X10 Tablets (Alu Alu)
 M.R.P. 165/-Per 10 Tablets



Composition :

Each film coated modified release tablet contains:
 Trimetazidine Dihydrochloride IP 60mg.
 Excipients q.s.
 Colour : Red oxide of Iron & Titanium Dioxide IP

Indication : Trimetazidine is an anti-ischemic (anti-anginal) metabolic agent, which improves myocardial glucose utilization through inhibition of fatty acid metabolism, also known as fatty acid oxidation inhibitor.

10X10 Tablets (Alu Alu)
 M.R.P. 215/-Per 10 Tablets

Tripaazine

Composition

Each film coated tablet contains :
 Trifluoperazine Hydrochloride IP 5mg.
 Excipients q.s.
 Colour : Titanium Dioxide of Iron

Indication : Trifluoperazine is used to treat the symptoms of schizophrenia (a mental illness that causes disturbed or unusual thinking, loss of interest in life, and strong or inappropriate emotions). Trifluoperazine is also used on a short-term basis to treat anxiety in people who have not been helped by other medications.



10X10 Tablets (Blister)

M.R.P. 26/-Per 10 Tablets

Trixcare-R2

Composition

Each uncoated tablet contains :
 Risperidone IP 2mg.
 Trihexyphenidyl Hydrochloride 2mg.
 Excipients q.s.
 Colour : Sunset Yellow

Indication : Risperidone +trihexyphenidyl Is Used To Treat Schizophrenia (a Mental Illness In Which A Person Is Confused). Risperidone+trihexyphenidyl Is A Fixed-dose Combination Composed Of Two Different Medicines, Namely, Risperidone And, Trihexyphenidyl.



20x10 Tablets (Blister)

M.R.P. 77/-Per 10 Tablets

Udicare-150 Tablets

Composition :

Each film coated tablet contains :
 Ursodeoxycholic Acid IP 150mg.
 Excipients q.s.
 Colour : Red oxide of Iron
 yellow oxide of Iron

Indication : Ursodeoxycholic acid is indicated in the treatment of primary biliary cirrhosis (PBC) and for the dissolution of radiolucent gallstones in patients with a functioning gall bladder.



5x2x10 Tablets (Blister)

M.R.P. 235/-Per 10 Tablets

Trixcare-2

Composition :

Each uncoated tablet contains
 Trihexyphenidyl Hydrochloride IP 2 mg.
 Excipients q.s.

Indication : Trihexyphenidyl is used for the symptomatic treatment of Parkinson's disease in mono and combination therapy. It is active in postencephalitic, arteriosclerotic, and idiopathic forms. The drug is also commonly used to treat extrapyramidal side effects occurring during antipsychotic treatment.



10X10 Tablets (Blister)

M.R.P. 14.10/-Per 10 Tablets

Trixcare-T

Composition

Each film coated tablet contains :
 Trifluoperazine Hydrochloride IP eqv. to Trifluoperazine 5mg.
 Trihexyphenidyl Hydrochloride IP 2mg.
 Colour : Indigo Carmine

Indication : Risperidone and aripiprazole are the only two medications approved by the Federal Drug Administration (FDA) for the treatment of irritability (including aggression, self-injurious behavior, temper tantrums, and mood swings) in school-age children and adolescents with autistic disorder.



10x10 Tablets (Blister)

M.R.P. 24/-Per 10 Tablets

Udicare-300 Tablets

Composition :

Each film coated tablet contains :
 Ursodeoxycholic Acid IP 300mg.
 Excipients q.s.
 Colour : Red oxide of Iron
 yellow oxide of Iron

Indication : Ursodeoxycholic acid is indicated in the treatment of primary biliary cirrhosis (PBC) and for the dissolution of radiolucent gallstones in patients with a functioning gall bladder.



5x2x10 Tablets (Blister)

M.R.P. 484/-Per 10 Tablets

Udicare-450 Tablets

Composition :

Each film coated tablet contains :
Ursodeoxycholic Acid IP 450mg.
(in sustained release form)
Excipients q.s.
Colour : Titanium Dioxide IP

Indication : Ursodeoxycholic acid is indicated in the treatment of primary biliary cirrhosis (PBC) and for the dissolution of radiolucent gallstones in patients with a functioning gall bladder.



5x2x10 Tablets (Blister)
M.R.P. 490/-Per 10 Tablets

URIPAIN[®]

Composition :

Each film coated tablet contains :
Flavoxate HCl IP 200 mg.
Excipients q.s.
Colour : Titanium Dioxide IP

Indication : Flavoxate is used to treat overactive bladder (a condition in which the bladder muscles contract uncontrollably and cause frequent urination, urgent need to urinate, and inability to control urination) relieve painful, frequent, or nighttime urination and urgency that may occur with infections of the prostate, bladder,



4x5x10 Tablets (Blister)
M.R.P. 185/-Per 10 Tablets

Valasivir-500

Composition :

Each film coated tablets contains :
Valacyclovir Hydrochloride USP
Eq. to Valacyclovir 500mg.
Excipients q.s.
Colour : Titanium Dioxide IP &

Indication : valacyclovir is an antiviral medication used to treat outbreaks of herpes simplex or herpes zoster. It is also used to prevent cytomegalovirus following a kidney transplant in high risk cases. It is taken by mouth. Common side effects include headache and vomiting.



10x1x3 Tablets (Blister)
M.R.P. 139/-Per 3 Tablets

Udicare-600 SR

Composition :

Each uncoated sustained release tablet contains :
Ursodeoxycholic Acid IP 600mg.
Excipients q.s.
Colour: Sunset Yellow FCF

Indication : Ursodeoxycholic acid is indicated in the treatment of primary biliary cirrhosis (PBC) and for the dissolution of radiolucent gallstones in patients with a functioning gall bladder.



10x1x10 Tablets (Blister)
M.R.P. 780/-Per 10 Tablets

URIPAIN[®]

Composition :

Each film coated tablet contains :
Flavoxate HCl IP 200 mg.
Excipients q.s.
Colour : Titanium Dioxide IP

Indication : Flavoxate is used to treat overactive bladder (a condition in which the bladder muscles contract uncontrollably and cause frequent urination, urgent need to urinate, and inability to control urination) relieve painful, frequent, or nighttime urination and urgency that may occur with infections of the prostate, bladder,



10x1x15 Tablets (Blister)
M.R.P. 275/-Per 15 Tablets

Valsartan-40

Composition :

Each film coated tablet contains:
Valsartan IP 40mg.
Excipients q.s.
Colour : Yellow oxide of iron & Titanium Dioxide IP

Indication : Valsartan belongs to a class of drugs called angiotensin receptor blockers (ARBs). It works by relaxing blood vessels so that blood can flow more easily. Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems.



10x1x10 Tablets (Alu Alu)
M.R.P. 85/-Per 10 Tablets

Valsartan-80

Composition :

Each film coated tablet contains:
 Valsartan IP 80mg.
 Excipients q.s.
 Colour : Yellow oxide of iron & Titanium Dioxide IP

Indication : Valsartan belongs to a class of drugs called angiotensin receptor blockers (ARBs). It works by relaxing blood vessels so that blood can flow more easily. Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems.



10x10 Tablets (Alu Alu)

M.R.P. 194/-Per 10 Tablets

Vasafe-20

Composition :

Each film coated tablet contains :
 Atorvastatin Calcium IP
 eq. to Atorvastatin 20mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Atorvastatin is used along with a proper diet to help lower "bad" cholesterol and fats (such as LDL, triglycerides) and raise "good" cholesterol (HDL) in the blood. It belongs to a group of drugs known as "statins." It works by reducing the amount of cholesterol made by the liver.



20x10 Tablets (Blister)

M.R.P. 138.24/-Per 10 Tablets

Vasafe-80

Composition :

Each film coated tablet contains :
 Atorvastatin Calcium IP
 eq. to Atorvastatin 80mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Atorvastatin is used along with a proper diet to help lower "bad" cholesterol and fats (such as LDL, triglycerides) and raise "good" cholesterol (HDL) in the blood. It belongs to a group of drugs known as "statins." It works by reducing the amount of cholesterol made by the liver.



20x10 Tablets (Alu Alu)

M.R.P. 325/-Per 10 Tablets

Vasafe-10

Composition :

Each film coated tablet contains :
 Atorvastatin Calcium IP
 eq. to Atorvastatin 10mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Atorvastatin is used along with a proper diet to help lower "bad" cholesterol and fats (such as LDL, triglycerides) and raise "good" cholesterol (HDL) in the blood. It belongs to a group of drugs known as "statins." It works by reducing the amount of cholesterol made by the liver.



20x10 Tablets (Blister)

M.R.P. 49.05/-Per 10 Tablets

Vasafe-40

Composition :

Each film coated tablet contains :
 Atorvastatin Calcium IP
 eq. to Atorvastatin 40mg.
 Excipients q.s.
 Colour : Titanium Dioxide IP

Indication : Atorvastatin is used along with a proper diet to help lower "bad" cholesterol and fats (such as LDL, triglycerides) and raise "good" cholesterol (HDL) in the blood. It belongs to a group of drugs known as "statins." It works by reducing the amount of cholesterol made by the liver.



20x10 Tablets (Blister)

M.R.P. 216.04/-Per 10 Tablets

Vasafe-E

Composition :

Each film coated tablet contains :
 Atorvastatin Calcium IP
 Eq. to Atorvastatin 10mg.
 Ezetimibe IP 10mg.
 Excipients q.s.
 Colour: Titanium Dioxide IP

Indication : Ezetimibe and atorvastatin combination is used together with a proper diet to treat high cholesterol and triglyceride (fats) levels in the blood. This medicine may help prevent medical problems (eg, heart attack, stroke) caused by clogged blood vessels.



10x10 Tablets (Alu Alu)

M.R.P. 130/-Per 10 Tablets

Vasafe-F

Composition :

Each film coated tablet contains :
 Atorvastatin Calcium IP
 Eq. to Atorvastatin 10mg.
 Fenofibrate IP 160mg
 Excipients q.s.
 Colour : Titanium Dioxide IP



Indication : Atorvastatin+fenofibrate Taken With Dietary Measures Helps In The Treatment Of High Cholesterol And Prevention Of Life-threatening Conditions To The Heart, Especially For Heart Patients When Initial Dietary Measures Fail To Lower Cholesterol.

10x10 Tablets (Alu Alu)
 M.R.P. 145/-Per 10 Tablets

Ventricose-500

Composition

Each film coated tablet contains :
 Diosmin BP 450mg.
 Hesperidin 50mg.
 Excipients q.s.
 Colour : Ferric Oxide Red, Ferric Oxide Yellow & Titanium Dioxide IP



Indication : MPFF is indicated as a first-line treatment of oedema and the symptoms of CVI in patients in any stage of the disease. In more advanced disease stages, MPFF may be used in conjunction with sclerotherapy, surgery and/or compression therapy, or as an alternative treatment when surgery is not indicated or is unfeasible.

10x10 Tablets (Blister)
 M.R.P. 299/-Per 10 Tablets

Vertifast[®]-8

Composition

Each uncoated tablet contains :
 Betahistine Hydrochloride IP 8mg.
 Excipients q.s.



Indication : Betahistine is a medicine used to treat the symptoms of Ménière's disease. Symptoms include: dizziness (vertigo) ringing in the ears (tinnitus)

10x10 Tablets (Alu Alu)
 M.R.P. 35/-Per 10 Tablets

Vertifast[®]-16

Composition :

Each uncoated tablet contains :
 Betahistine Hydrochloride IP 16mg.
 Excipients q.s.



Indication : Betahistine is a medicine used to treat the symptoms of Ménière's disease. Symptoms include: dizziness (vertigo) ringing in the ears (tinnitus)

10x15 Tablets (Alu Alu)
 M.R.P. 148/-Per 15 Tablets

Velfax-37.5

Composition

Each uncoated prolonged -release tablet contains :
 Venlafaxine Hydrochloride BP 37.5mg
 Excipients q.s.



Indication : It works by increasing the amounts of serotonin and norepinephrine, natural substances in the brain that help maintain mental balance.

10x10 Tablets (Blister)
 M.R.P. 65/-Per 10 Tablets

Velfax-75

Composition

Each uncoated prolonged -release tablet contains :
 Venlafaxine Hydrochloride BP 75mg
 Excipients q.s.



Indication : It works by increasing the amounts of serotonin and norepinephrine, natural substances in the brain that help maintain mental balance.

10x10 Tablets (Blister)
 M.R.P. 122/-Per 10 Tablets



TABLETS

Vildaily-DZ 5/100 SR

Composition :

Each film coated Bi-Layered tablet contains:
 eq. to Dapagliflozin 5mg.
 Vildagliptin 100mg.
 (As Sustained Release)
 Excipients q.s.
 Colour : Sunset Yellow



Indication : Dapagliflozin+Vildagliptin is used in the treatment of Type 2 diabetes mellitus.

10x10 Tablets (Alu Alu)

M.R.P. 425/-Per 10 Tablets

Vildaily-DZ 10/100 SR

Composition :

Each film coated Bi-Layered tablet contains:
 eq. to Dapagliflozin 10mg.
 Vildagliptin 100mg.
 (As Sustained Release)
 Excipients q.s.
 Colour : Sunset Yellow



Indication : Dapagliflozin+Vildagliptin is used in the treatment of Type 2 diabetes mellitus.


10x10 Tablets (Alu Alu)

M.R.P. 495/-Per 10 Tablets

Vildaily-50

Composition :

Each uncoated tablet contains :
 Vildagliptin 50mg.
 Excipients q.s.
 Colour: Titanium Dioxide IP



Indication : It helps to control the levels of sugar (glucose) in your blood. If your body does not make enough insulin, or if it does not use the insulin it makes effectively, this results in the condition called

10x15 Tablets (Alu Alu)

M.R.P. 255/-Per 15 Tablets

Vildaily-M

Composition :

Each film coated tablet contains :
 Vildagliptin 50mg.
 Metformin Hydrochloride IP 500mg.
 Excipients q.s.
 Colour : Iron Oxide red, Iron oxide Yellow & Titanium Dioxide IP.



Indication : Metformin and vildagliptin are both anti-diabetic agents that are used in combination to treat Diabetes Mellitus Type 2. In particular, metformin increases the body's sensitivity to insulin.

10x15 Tablets (Alu Alu)

M.R.P. 205.96/-Per 15 Tablets

Vildaily-M Forte

Composition :

Each film coated tablet contains :
 Vildagliptin 50mg.
 Metformin Hydrochloride IP 1000mg.
 Excipients q.s.
 Colour : Iron Oxide red, Iron oxide Yellow & Titanium Dioxide IP.



Composition : Metformin and vildagliptin are both anti-diabetic agents that are used in combination to treat Diabetes Mellitus Type 2. In particular, metformin increases the body's sensitivity to insulin.

10x15 Tablets (Alu Alu)

M.R.P. 405/-Per 15 Tablets

Vorhappy-10

Composition

Each Film coated tablet contains :
 Vortioxetine Hydrobromide
 Eq. to Vortioxetine 10mg
 Excipients q.s.
 Colour : Yellow oxide of Iron & TitaniumDioxide IP



Indication : Vortioxetine is used to treat depression in adults. Vortioxetine is in a class of medications called serotonin modulators. It works mainly by increasing the amount of serotonin, a natural substance in the brain that helps maintain mental balance.

10x10 Tablets (Alu Alu)

M.R.P. 188/-Per 10 Tablets



TABLETS

Voglicare Rep 0.2/1

Composition :

Each uncoated tablet contains :

Voglibose	IP	0.2mg.
Repaglinide	IP	1mg.
Excipients		q.s.



Indication : Repaglinide+voglibose is a combination drug that belongs to the anti-diabetic class. It is primarily used to treat type 2 diabetes. Type 2 diabetes is a common condition that causes the level of glucose (sugar) in the blood to become too high. It's a lifelong condition that can affect your everyday life.

10x10 Tablets (Alu Alu)
M.R.P. 147/-Per 10 Tablets

Voglicare Rep 0.2/0.5

Composition :

Each uncoated tablet contains :

Voglibose	IP	0.2mg.
Repaglinide	IP	0.5mg.
Excipients		q.s.



Indication : Repaglinide+voglibose is a combination drug that belongs to the anti-diabetic class. It is primarily used to treat type 2 diabetes. Type 2 diabetes is a common condition that causes the level of glucose (sugar) in the blood to become too high. It's a lifelong condition that can affect your everyday life.

10x10 Tablets (Alu Alu)
M.R.P. 121/-Per 10 Tablets

Voglicare[®]-0.2

Composition :

Each uncoated tablet contains :

Voglibose IP	0.2 mg.
Excipients	q.s.



Indication : Voglibose is an alpha-glucosidase inhibitor used for lowering post-prandial blood glucose levels in people with diabetes mellitus. It is made in India by Ranbaxy Labs and sold under the trade name Volix.

4x5x10 Tablets (Blister)
M.R.P. 70/-Per 10 Tablets

Voglicare[®]-0.3

Composition :

Each uncoated tablet contains :

Voglibose IP	0.3 mg.
Excipients	q.s.



Indication : Voglibose is an alpha-glucosidase inhibitor used for lowering post-prandial blood glucose levels in people with diabetes mellitus. It is made in India by Ranbaxy Labs and sold under the trade name Volix.

4x5x10 Tablets (Blister)
M.R.P. 88/-Per 10 Tablets

Voglicare[®]-M 0.2

Composition :

Each film coated tablet contains :

Voglibose	IP	0.2mg.
Metformin HCl	IP	500mg.
(As Sustained Release)		
Excipients		q.s.
Colour : Sunset Yellow FCF		



Indication : Metformin+Voglibose is used in the treatment of type 2 diabetes mellitus. Metformin + Voglibose is a combination of two antidiabetic medicines: Metformin and voglibose. Metformin lowers the production and absorption of sugar in your body and allows better use of existing insulin.

10x10 Tablets (Blister)
M.R.P. 78/-Per 10 Tablets

Voglicare[®]-M 0.3

Composition :

Each uncoated bilayered tablet contains :

Voglibose	IP	0.3mg.
Metformin Hydrochloride	IP	500mg.
(in extended release form)		
Excipients		q.s.
Colour : Ponceau 4R		



Indication : Metformin+Voglibose is used in the treatment of type 2 diabetes mellitus. Metformin + Voglibose is a combination of two antidiabetic medicines: Metformin and voglibose. Metformin lowers the production and absorption of sugar in your body and allows better use of existing insulin.

10x15 Tablets (Blister)
M.R.P. 62/-Per 15 Tablets

Voglicare[®]-MG 3

Composition :

Each uncoated bilayered tablet contains :

Glimepiride	IP	2mg
Metformin		
Hydrochloride	IP	
(in Sustained release form)		500mg
Voglibose	IP	0.3mg
Excipients		q.s.

Colour : Ferric Oxide USP-NF Yellow and Ferric Oxide USP-NF Red

Indication : is used to lower the blood sugar levels in diabetes type-2 patients. Insulin helps in uptake of glucose by muscles.

10x15 Tablets (Blister)
M.R.P. 165/-Per 15 Tablets



Voglicare[®]-MG 2

Composition :

Each uncoated bilayered tablet contains :

Glimepiride	IP	2mg
Metformin		
Hydrochloride	IP	
(in Sustained release form)		500mg
Voglibose	IP	0.2mg
Excipients		q.s.

Colour : Ferric Oxide USP-NF Yellow and Ferric Oxide USP-NF Red

Indication : is used to lower the blood sugar levels in diabetes type-2 patients. Insulin helps in uptake of glucose by muscles.

10x10 Tablets (Blister)
M.R.P. 93.50/-Per 10 Tablets



Warfarin-2

Composition :

Each uncoated tablet contains:
 Warfarin Sodium Clathrate IP
 Eq. to (Anhydrous) Warfarin Sodium 2mg.
 Excipients q.s.

Indication : Warfarin is an anticoagulant (blood thinner). Warfarin reduces the formation of blood clots. Warfarin is used to treat or prevent blood clots in veins or arteries, which can reduce the risk of stroke, heart attack, or other serious

20X10 Tablets (Blister)
M.R.P. 30.46/-Per 10 Tablets



Warfarin-5

Composition :

Each uncoated tablet contains:
 Warfarin Sodium Clathrate IP
 Eq. to (Anhydrous) Warfarin Sodium 5mg.
 Excipients q.s.

Indication : Warfarin is an anticoagulant (blood thinner). Warfarin reduces the formation of blood clots. Warfarin is used to treat or prevent blood clots in veins or arteries, which can reduce the risk of stroke, heart attack, or other serious conditions.

20x10 Tablets (Blister)
M.R.P. 26.30/-Per 10 Tablets





Valstep

Sacubitril & Valsartan Tablets

Valstep-S 100

Composition

Each film coated tablet contains :
Sacubitril Sodium eq. to Sacubitril 49mg.
Valsartan IP 51mg.
Excipients q.s.
Colour : Red Oxide of Iron, Black Oxide of Iron & Titanium Dioxide IP

Indication : Sacubitril Valsartan is a combination of a neprilysin inhibitor and a an angiotensin II receptor blocker. It treats heart failure. This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.

10x14 Tablets (Alu Alu)
M.R.P. 630/-Per 14 Tablets



Valstep-S 200

Composition

Each film coated tablet contains :
Sacubitril Sodium eq. to Sacubitril 97mg.
Valsartan IP 103mg.
Excipients q.s.
Colour : Red Oxide of Iron, Black Oxide of Iron & Titanium Dioxide IP

Indication : Sacubitril Valsartan is a combination of a neprilysin inhibitor and a an angiotensin II receptor blocker. It treats heart failure. This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.

10x7 Tablets (Alu Alu)
M.R.P. 615/-Per 7 Tablets



Valstep-S 50

Composition

Each film coated tablet contains :
Sacubitril Sodium eq. to Sacubitril 24mg.
Valsartan IP 26mg.
Excipients q.s.
Colour : Red Oxide of Iron, Black Oxide of Iron & Titanium Dioxide IP

Indication : Sacubitril Valsartan is a combination of a neprilysin inhibitor and a an angiotensin II receptor blocker. It treats heart failure. This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.

10x14 Tablets (Alu Alu)
M.R.P. 490/-Per 14 Tablets



Keeping hearts resilient, strong and restored.

Hardgelatin Capsules



Acethama[®]-100

Composition :
Each Hard gelatin Capsules :
Acebrophylline 100mg.
Excipients q.s.

Indication : Acebrophylline works by blocking the effects of leukotriene (substance produced by the body in response to certain triggers) that cause narrowing and swelling of airways in the lungs. Acebrophylline is used in the prevention and treatment of asthma and chronic obstructive pulmonary disease.



10x10 Capsules (Blister)

M.R.P. 95/-Per 10 Caps.

Dbgron-110

Composition :
Each hard gelatin capsule contains :
Dabigatran Etexilate Mesilate
equivalent to Dabigatran

Etexilate 110 mg
Colours : Brilliant blue & Titanium
Dioxide IP

Indication : Dabigatran is used to treat deep vein thrombosis (DVT; a blood clot, usually in the leg) and pulmonary embolism (PE; a blood clot in the lung) in adults and children 3 months of age and older who have been treated with an injectable anticoagulant ('blood thinner').



10x1x10 Capsules (Alu Alu)

M.R.P. 390/-Per 10 Caps.

Duloxit-30

Composition :
Each hard gelatin capsules
contains :
Duloxetine Hydrochloride IP
Eq. to Duloxetine 30mg.
(As enteric coated pellets)
Excipients q.s.

Indication : Duloxetine is an antidepressant medicine. It's used to treat depression and anxiety. It's also used to treat nerve pain, such as fibromyalgia, and can be used to treat stress in women.



10x10 Capsules (Blister)

M.R.P. 120/-Per 10 Caps.

Caresil-300

Composition :
Each hard gelatin capsule contains
Clindamycin HCl IP 300mg.
Eq. to Clindamycin 300mg.
Excipients q.s.

Indication : Clindamycin is used to treat certain types of bacterial infections, including infections of the lungs, skin, blood, female reproductive organs, and internal organs. Clindamycin is in a class of medications called lincosamin antibiotics. It works by slowing or stopping the growth of bacteria.



10x10 Capsules (Blister)

M.R.P. 242/-Per 10 Caps.

Dbgron-150

Composition :
Each hard gelatin capsule contains :
Dabigatran
Etexilate Mesylate 172.95 mg.
Eq. to Dabigatran Etexilate 150 mg.
Excipients q.s.

Indication : Dabigatran is used to treat deep vein thrombosis (DVT; a blood clot, usually in the leg) and pulmonary embolism (PE; a blood clot in the lung) in adults and children 3 months of age and older who have been treated with an injectable anticoagulant ('blood thinner').



10x1x10 Capsules (Alu Alu)

M.R.P. 710/-Per 10 Caps.

Fluoxetine

Composition :
Each hard gelatin capsule contains
Fluoxetine HCl I.P
Eq. To Fluoxetine 20mg.
Excipients q.s.

Indication : Fluoxetine is a type of antidepressant known as an SSRI (selective serotonin reuptake inhibitor). It is often used to treat depression, and also sometimes obsessive compulsive disorder and bulimia. Fluoxetine helps many people recover from depression, and it has fewer unwanted effects than older antidepressants.



20x10 Capsules (Blister)

M.R.P. 39.80/-Per 10 Caps.

Flupry-100

Composition :

Each hard gelatin capsule contains :
 Flupirtine Maleate 100mg.
 Excipients q.s.

Indication : Flupirtine is a non-opioid painkiller that has been used to treat pain, such as pain associated with muscle tension, cancer pain, menstrual pain and pain following orthopaedic surgery or injury



10x10 Capsules (Blister)

M.R.P. 124/-Per 10 Caps.

Hydoxip-500

Composition :

Each Hard Gelatin Capsule contains :
 Hydroxyurea USP 500mg.
 Excipients q.s.

Indication : Hydroxyurea is used to prevent painful episodes and reduce the need for blood transfusions in patients with sickle cell anemia. It works by making the red blood cells more flexible. This medicine is available only with your doctor's prescription.



10x10 Capsules (Blister)

M.R.P. 113.50/-Per 10 Caps.

Intocare®-SB 50

Composition :

Each Hard Gelatin Capsule contains :
 Itraconazole BP 50mg.
 Excipients q.s.

Indication : Itraconazole capsules are used to treat fungal infections in the lungs that can spread throughout the body. Itraconazole capsules are also used to treat fungal infections of the fingernails. Itraconazole tablets and capsules are used to treat fungal infections of the toenails.



10x1x10 Capsules (Alu Alu)

M.R.P. 190/-Per 10 Caps.

Gebcare-300

Composition :

Each hard gelatin capsule contains :
 Gabapentin I.P. 300 mg.
 Excipients q.s.

Indication : Gabapentin is used to treat epilepsy. It's also taken for nerve pain. Nerve pain can be caused by different illnesses, including diabetes and shingles, or it can happen after an injury. Occasionally, gabapentin is used to treat migraine headaches.



10x10 Capsules (Blister)

M.R.P. 155/-Per 10 Caps.

Indolus®-75 ER

Composition :

Each hard gelatin capsule contains :
 Indomethacin IP 75mg.
 (In extended release form)

Indication : Indomethacin is used to relieve moderate to severe pain, tenderness, swelling, and stiffness caused by osteoarthritis (arthritis caused by a breakdown of the lining of the joints), rheumatoid arthritis (arthritis caused by swelling of the lining of the joints), and ankylosing spondylitis (arthritis that mainly affects).



10x10 Capsules (Blister)

M.R.P. 115/-Per 10 Caps.

Intocare®-100

Composition :

Each hard gelatin capsule contains :
 Itraconazole pellets eq. to 100mg.
 Excipients q.s.
 Colour : Approved colour used empty capsules shells.

Indication : Itraconazole capsules are used to treat fungal infections in the lungs that can spread throughout the body. Itraconazole capsules are also used to treat fungal infections of the fingernails. Itraconazole tablets and capsules are used to treat fungal infections of the toenails.



10x10 Capsules (Blister)

M.R.P. 170/-Per 10 Caps.

Intocare[®]-200

Composition :

Each hard gelatin capsule contains :
 Itraconazole pellets eq. to Itraconazole BP 200mg.
 Excipients q.s.
 Colour : Approved colour used empty capsules shells.

Indication : Itraconazole capsules are used to treat fungal infections in the lungs that can spread throughout the body. Itraconazole capsules are also used to treat fungal infections of the fingernails. Itraconazole tablets and capsules are used to treat fungal infections of the toenails.


10x10 Capsules (Blister)
M.R.P. 247/-Per 10 Caps.

Mebaus-200

Composition

Each hard gelatin capsule contains
 Mebeverine Hydrochloride IP
 Eq. to Mebeverine 200mg.
 (as prolonged release pellets)
 Excipients q.s.

Indication : Mebeverine (sometimes called mebeverine hydrochloride) is a type of medicine known as an antispasmodic. It helps with muscle spasms. It can be used to ease painful stomach cramps if you have irritable bowel syndrome (IBS) or other conditions. It works by relaxing certain muscles in your gut.


10x10 Capsule (Alu Alu)
M.R.P. 235/-Per 10 Caps.

Nekszole[®]-LS

Composition :

Each hard gelatin capsule contains :
 Esomeprazole
 Magnesium Trihydrate IP
 Eq. to Esomeprazole 40 mg.
 (As enteric coated pellets)
 Levosulpiride 75mg.
 (As Sustained release pellets)
 Excipients q.s.

Indication: Domperidone+Esomeprazole is used in the treatment of gastroesophageal reflux disease (acid reflux) and functional dyspepsia. Domperidone + Esomeprazole is a combination of two medicines: Domperidone and Esomeprazole.


10x1x10 Capsules (Alu Alu)
M.R.P. 241/-Per 10 Caps.

Lenocell-10

Composition

Each hard gelatin capsule contains
 Lenalidomide 10mg.
 Excipients q.s.

Indication : Lenalidomide is used to treat anemia (low red blood cells) in patients with a certain type of myelodysplastic syndrome (MDS). Patients with MDS may have very low red blood cell counts and require blood transfusions.



1x10 (Capsule)
M.R.P. 1202/-Per 10 Caps.

Nekszole[®]-DSR

Composition :

Each hard gelatin capsule contains :
 Esomeprazole Sodium IP 40mg.
 (as enteric coated pellets)
 Domperidone IP 30mg.
 (as sustained release pellets)
 Excipients q.s.

Indication: Domperidone+Esomeprazole is used in the treatment of gastroesophageal reflux disease (acid reflux) and functional dyspepsia. Domperidone + Esomeprazole is a combination of two medicines: Domperidone and Esomeprazole.


10x10 Capsules (Alu Alu)
M.R.P. 160/-Per 10 Caps.

Olpavir[®]-75

Composition :

Each hard gelatin capsule contains :
 Oseltamivir Phosphate IP
 Eq. to Oseltamivir 75mg.
 Excipients q.s.

Indication : Oseltamivir is used to treat symptoms caused by the flu virus (influenza). It helps make the symptoms (such as stuffy nose, cough, sore throat, fever/chills, aches, tiredness) less severe and shortens the recovery time by 1-2 days.


10x10 Capsules (Blister)
M.R.P. 620/-Per 10 Caps.

Orlifit-N-Slim-120

Composition :

Each hard gelatin capsule contains :
Orlistat USP 120mg.
Excipients q.s.

Indication : Orlistat (prescription and nonprescription) is used with an individualized low-calorie, low-fat diet and exercise program to help people lose weight. Prescription orlistat is used in overweight people who may also have high blood pressure, diabetes, high cholesterol, or heart disease.



10x10 Capsules (Alu Alu)
M.R.P.475/-Per 10 Caps.

Pantaus®-10000

Composition :

Each hard gelatin capsule contains :
Pancreatin (entetic coated pellets)
eq. to Pancreatin IP 150mg.
Excipients q.s.
Colour : Approved colour used in empty capsule shells.



Pancreatin Minimicropellets with declared enzyme activity :

Lipase	10000 USP units
Protease	37500 USP units
Amylase	33200 USP units

10x10 Capsules (Aluminium)
M.R.P.355/-Per 10 Caps.

Pgnerv®-75

Composition :

Each hard gelatin capsule contains:
Pregabalin IP 75 mg.
Excipients q.s.

Indication : This medication is used to treat pain caused by nerve damage due to diabetes or shingles (herpes zoster) infection.



10x10 Capsules (Blister)
M.R.P.145/-Per 10 Caps.

Ostokey®-DOB

Composition :

Each hard gelatin capsule contains :
Calcium Debesilate
Monohydrate IP 500mg.
Excipients q.s.

Indication : Calcium dobesilate (Doxium) is a veno-tonic drug, which is widely prescribed in more than 60 countries from Europe, Latin America, Asia and the Middle East for three main indications: chronic venous disease, diabetic retinopathy and the symptoms of haemorrhoidal attack.




10x10 Capsules (Blister)
M.R.P.129/-Per 10 Caps.

Pantaus®-25000

Composition :

Each hard gelatin capsule contains :
Pancreatin (entetic coated pellets)
eq. to Pancreatin IP 300mg.
Excipients q.s.
Colour : Approved colour used in empty capsule shells.



Pancreatin Minimicropellets with declared enzyme activity :

Lipase	25000 USP units
Protease	62500 USP units
Amylase	74700 USP units

10x10 Capsules (Aluminium)
M.R.P.649/-Per 10 Caps.

Pgnerv®-M

Composition :

Each hard gelatin capsule contains
Pregabalin IP 75 mg.
Methylcobalamin USP 750 mcg.
Excipients q.s.

Indication : This medication is used to treat pain caused by nerve damage due to diabetes, shingles (herpes zoster) infection, or spinal cord injury. This medication is also used to treat pain in people with fibromyalgia. It is also used with other medications to treat certain types of seizures (focal seizures).



10x10 Capsules (Alu Alu)
M.R.P. 146/-Per 10 Caps.



Pgnerv-M Forte

Composition :

Each hard gelatin capsule contains
Pregabalin IP 75 mg.
Methylcobalamin USP 1500mcg.
Excipients q.s.
Approved colour used in empty capsule shells

Indication : Pregabalin and Methylcobalamin. This medicine is useful in the treatment of nerve pain (neuropathic pain). It calms the damaged or overactive nerves by acting on the brain, thereby decreasing the pain sensation



10x1x10 Capsules (Blister)

M.R.P. 245/-Per 10 Caps.

Palbocell-125

Composition

Each hard gelatin capsule contains :
Palbociclib 125mg.
Excipients q.s.

Indication : Palbociclib is in a class of medications called kinase inhibitors. It works by blocking the action of the abnormal protein that signals cancer cells to multiply. This helps stop or slow the spread of cancer cells.



1x21 Capsules


M.R.P. 7600/-Per 21 Caps.

Rosudoz-AS 10

Composition :

Each hard gelatin capsule contains :
Rosuvastatin Calcium IP
eq. to Rosuvastatin 10mg.
(As granules)
Aspirin IP 75mg.
(As enteric coated tablet)
Colour : Titanium Dioxide IP

Indication : Rosuvastatin+Aspirin is used to prevent prevention of heart attack and stroke. Rosuvastatin + Aspirin is a combination of two medicines: Rosuvastatin and Aspirin, which prevents heart attack and stroke. Rosuvastatin is a lipid-lowering medication that blocks an enzyme required in the body to make cholesterol.



10x15 Capsules (Aluminium)

M.R.P. 120/-Per 15 Caps.

Rosudoz-AS 20

Composition :

Each hard gelatin capsule contains :
Rosuvastatin Calcium IP
eq. to Rosuvastatin 20mg.
(As granules)
Aspirin IP 75mg.
(As enteric coated tablet)
Colour : Titanium Dioxide IP
Excipients q.s.

Indication : Rosuvastatin+Aspirin is used to prevent prevention of heart attack and stroke. Rosuvastatin + Aspirin is a combination of two medicines: Rosuvastatin and Aspirin, which prevents heart attack and stroke. Rosuvastatin is a lipid-lowering medication that blocks an enzyme required in the body to make cholesterol.



10x15 Capsules (Aluminium)

M.R.P. 87/-Per 15 Caps.

Rosudoz-CV 10/75

Composition :

Each hard gelatin capsule contains :
Rosuvastatin Calcium IP
Eq. to Rosuvastatin 10mg.
(As Pellets)
Clopidogrel Bisulphate IP
Eq. to Clopidogrel 75mg.
(As Pellets)
Excipients q.s.

Indication : Rosuvastatin + Clopidogrel is a combination of two medicines: Rosuvastatin and Clopidogrel, which prevents heart attack and stroke. Rosuvastatin is a lipid lowering medication that blocks the body's ability to produce "bad" cholesterol (LDL) and improves the level of "good" cholesterol (HDL).



10x10 Capsules (Alu Alu)

M.R.P. 147/-Per 10 Caps.

Rosudoz-CV 20/75

Composition :

Each hard gelatin capsule contains :
Rosuvastatin Calcium IP
Eq. to Rosuvastatin 20mg.
(As Pellets)
Clopidogrel Bisulphate IP
Eq. to Clopidogrel 75mg.
(As Pellets)
Excipients q.s.

Indication : Rosuvastatin + Clopidogrel is a combination of two medicines: Rosuvastatin and Clopidogrel, which prevents heart attack and stroke. Rosuvastatin is a lipid lowering medication that blocks the body's ability to produce "bad" cholesterol (LDL) and improves the level of "good" cholesterol (HDL).



10x10 Capsules (Alu Alu)

M.R.P. 179/-Per 10 Caps.

Rosudoz-Gold 10

Composition :

Each hard gelatin capsule contains :
 Rosuvastatin Calcium IP
 Eq. to Rosuvastatin 10mg.
 (As Granules)
 Clopidogrel Bisulphate IP
 Eq. to Clopidogrel 75mg.
 (As Granules)
 Aspirin IP 75mg.
 (As Gastro Resistant Tablet)
 Colour : Titanium Dioxide IP
 Excipients q.s.

Indication : Rosuvastatin + Clopidogrel is a combination of two medicines: Rosuvastatin and Clopidogrel, which prevents heart attack and stroke. Rosuvastatin is a lipid lowering medication that blocks the body's ability to produce "bad" cholesterol (LDL) and improves the level of "good" cholesterol (HDL).



10x15 Capsules (Aluminium)
M.R.P. 210/-Per 15 Caps.

Rosudoz-Gold 20

Composition :

Each hard gelatin capsule contains :
 Rosuvastatin Calcium IP
 Eq. to Rosuvastatin 20mg.
 (As Granules)
 Clopidogrel Bisulphate IP
 Eq. to Clopidogrel 75mg.
 (As Granules)
 Aspirin IP 75mg.
 (As Gastro Resistant Tablet)
 Colour : Titanium Dioxide IP
 Excipients q.s.

Indication : Rosuvastatin + Clopidogrel is a combination of two medicines: Rosuvastatin and Clopidogrel, which prevents heart attack and stroke. Rosuvastatin is a lipid lowering medication that blocks the body's ability to produce "bad" cholesterol (LDL) and improves the level of "good" cholesterol (HDL).



10x15 Capsules (Aluminium)
M.R.P. 258/-Per 15 Caps.

Silodom-4

Composition :

Each hard gelatin capsule contains
 Silodosin 4mg.
 Excipients q.s.

Indication : Silodosin is used in men to treat the symptoms of an enlarged prostate (benign prostatic hyperplasia; BPH), which include difficulty urinating (hesitation, dribbling, weak stream, and incomplete bladder emptying), painful urination, and urinary frequency and urgency.



10x1x10 Capsules (Alu Alu)
M.R.P. 175/-Per 10 Caps.

Silodom-8

Composition :

Each hard gelatin capsule contains
 Silodosin 8mg.
 Excipients q.s.

Indication : Silodosin is used in men to treat the symptoms of an enlarged prostate (benign prostatic hyperplasia; BPH), which include difficulty urinating (hesitation, dribbling, weak stream, and incomplete bladder emptying), painful urination, and urinary frequency and urgency.



10x1x10 Capsules (Alu Alu)
M.R.P. 350/-Per 10 Caps.

Silodom-D

Composition :

Each hard gelatin capsule contains :
 Silodosin 8 mg
 Dutasteride 0.5 mg
 (As uncoated tablet)
 Excipients q.s.
 Approved colours used in empty capsule shells.

Indication : Silodosin is used in men to treat the symptoms of an enlarged prostate (benign prostatic hyperplasia; BPH), which include difficulty urinating (hesitation, dribbling, weak stream, and incomplete bladder emptying), painful urination, and urinary frequency and urgency.



10x1x10 Capsules (Aluminium)
M.R.P. 450/-Per 10 Caps.

Sporiphax-500

Composition

Each hard gelatin capsule contains :
 Cephalexin IP
 eqv. to Anhydrous Cephalexin 500mg.
 Excipients q.s.

Indication : Cephalexin is used to treat certain infections caused by bacteria such as pneumonia and other respiratory tract infections; and infections of the bone, skin, ears, , genital, and urinary tract. Cephalexin is in a class of medications called cephalosporin antibiotics.



10x10 Capsules (Blister)
M.R.P. 210/-Per 10 Caps.

Tacrofast-0.5

Composition :

Each hard gelatin capsule contains:
 Tacrolimus IP 0.5mg.
 Excipients q.s.

Indication : Tacrolimus is used with other medications to prevent rejection of a kidney, heart, or liver transplant. This medication belongs to a class of drugs known as immunosuppressants. It works by weakening your body's defense system (immune system) to help your body accept the new organ as if it were your own



10x1x10 Capsules (Alu Alu)
 M.R.P. 206.40/-Per 10 Caps.

Tacrofast-1

Composition :

Each hard gelatin capsule contains:
 Tacrolimus IP 1mg.
 Excipients q.s.

Indication : Tacrolimus is used with other medications to prevent rejection of a kidney, heart, or liver transplant. This medication belongs to a class of drugs known as immunosuppressants. It works by weakening your body's defense system (immune system) to help your body accept the new organ as if it were your own



10x1x10 Capsules (Alu Alu)
 M.R.P. 405/-Per 10 Caps.

Tacrofast-2

Composition :

Each hard gelatin capsule contains:
 Tacrolimus IP 2mg.
 Excipients q.s.

Indication : Tacrolimus is used with other medications to prevent rejection of a kidney, heart, or liver transplant. This medication belongs to a class of drugs known as immunosuppressants. It works by weakening your body's defense system (immune system) to help your body accept the new organ as if it were your own



10x1x10 Capsules (Alu Alu)
 M.R.P. 842.40/-Per 10 Caps.

Temida-100

Composition

Each hard gelatin capsule contains :
 Temozolomide IP 100mg.
 Excipients q.s.

Indication : Temozolomide is used to treat certain types of brain tumors. Temozolomide is in a class of medications called alkylating agents. It works by slowing or stopping the growth of cancer cells in your body.



1x5 Capsules
 M.R.P. 2765/-Per 5 Caps.

Temida-250

Composition

Each hard gelatin capsule contains :
 Temozolomide IP 250mg.
 Excipients q.s.

Indication : Temozolomide is used to treat certain types of brain tumors. Temozolomide is in a class of medications called alkylating agents. It works by slowing or stopping the growth of cancer cells in your body.



1x5 Capsules
 M.R.P. 5720.65/-Per 5 Caps.

Thioaus-4

Composition :

Each hard gelatin capsule contains :
 Thiocolchicoside IP 4mg.
 Excipients q.s.
 Approved colour used in empty capsule shells.

Indication : Thiocolchicoside is used for muscle relaxation. It relieves painful spasms of the skeletal muscles. Thiocolchicoside is a muscle relaxant. It works on the centres in the brain and spinal cord to relieve muscle stiffness or spasm without reduction in strength.



10x1x10 Capsules (Blister)
 M.R.P. 190/-Per 10 Caps.

Vasafe-AS 10

Composition :

Each hard gelatin capsule contains :
 Atorvastatin Calcium IP
 Eq. to Atorvastatin 10mg
 (As Granules)
 Aspirin IP 75mg
 (As Gastro Resistant Tablet)
 Colour : Titanium Dioxide IP
 Excipients q.s.

Indication : Atorvastatin is a lipid-lowering medication that blocks an enzyme required in the body to make cholesterol. It lowers the "bad" cholesterol (LDL), triglycerides and raises the "good" cholesterol (HDL). Aspirin is a non-steroidal anti-inflammatory drug (NSAID) with anti-platelet action.

10x10 Capsules (Alu Alu)
M.R.P. 61.45/-Per 10 Caps.



Vasafe-AS 20

Composition :

Each hard gelatin capsule contains :
 Atorvastatin Calcium IP
 Eq. to Atorvastatin 20mg
 (As Granules)
 Aspirin IP 75mg
 (As Gastro Resistant Tablet)
 Colour : Titanium Dioxide IP
 Excipients q.s.

Indication : Atorvastatin is a lipid-lowering medication that blocks an enzyme required in the body to make cholesterol. It lowers the "bad" cholesterol (LDL), triglycerides and raises the "good" cholesterol (HDL). Aspirin is a non-steroidal anti-inflammatory drug (NSAID) with anti-platelet action.

10x10 Capsules (Alu Alu)
M.R.P. 71/-Per 10 Caps.



Vasafe-Gold 10

Composition :

Each hard gelatin capsule contains :
 Atorvastatin Calcium IP
 Eq. to Atorvastatin 10mg.
 (As Granules)
 Clopidogrel Bisulphate IP
 Eq. to Clopidogrel 75mg.
 (As Granules)
 Aspirin IP 75mg.
 (As Gastro Resistant Tablet)
 Colour : Titanium Dioxide IP
 Excipients q.s.

Indication : Atorvastatin is a lipid-lowering medication that blocks an enzyme required in the body to make cholesterol. It lowers the "bad" cholesterol (LDL), triglycerides and raises the "good" cholesterol (HDL). Aspirin is a non-steroidal anti-inflammatory drug (NSAID) with anti-platelet action.

10x15 Capsules (Aluminium)
M.R.P. 112/-Per 15 Caps.



Vasafe-Gold 20

Composition :

Each hard gelatin capsule contains :
 Atorvastatin Calcium IP
 Eq. to Atorvastatin 20mg.
 (As Granules)
 Clopidogrel Bisulphate IP
 Eq. to Clopidogrel 75mg.
 (As Granules)
 Aspirin IP 75mg.
 (As Gastro Resistant Tablet)
 Colour : Titanium Dioxide IP
 Excipients q.s.

Indication : Atorvastatin is a lipid-lowering medication that blocks an enzyme required in the body to make cholesterol. It lowers the "bad" cholesterol (LDL), triglycerides and raises the "good" cholesterol (HDL). Aspirin is a non-steroidal anti-inflammatory drug (NSAID) with anti-platelet action.

10x15 Capsules (Aluminium)
M.R.P. 178/-Per 15 Caps.



Zokovit

NUTRITIONAL INFORMATION (APPROXIMATE VALUES)	PER CAPSULES	RDA %
Vitamin A (Retinyl Palmitate)	600 mcg	60.0 %
Vitamin E (DL-Alpha-tocopheryl Acetate)	5 mg	50 %
Vitamin B1 (Thiamine Mononitrate)	1.7 mg	73.91 %
Vitamin B2 (Riboflavin 5-Phosphate)	2.1 mg	65.63 %
Vitamin B6 (Pyridoxal 5-Phosphate)	2 mg	64.52 %
Vitamin B12 (Cyanocobalamin)	1 mcg	45.45 %
Vitamin D3 (Cholecalciferol)	200 IU	33.33 %
Elemental Calcium Phosphate	255 mg	12.5 %
Magnesium Oxide		--
Eq. to Elemental Magnesium	55 mg	12.5 %
L-Lysine	2 mg	--
Zinc (as Zinc Oxide)	10 mg	58.82 %
Sodium Molybdate Dihydrate		--
Eq. to Elemental Molybdenum	30 mcg	66.67 %

CONTAINS PERMITTED SYNTHETIC FOOD COLOUR USED E120, E124 & E130 IN SOFTGEL CAPSULE SHELLS.
 Health Supplement - Not Intended To Treat, Mitigate, Diagnose Or Prevent Any Disease Or Disorder In Human Being.
Other Ingredients: Gelatin (Animal Origin Capsule Shell), Glycerin, Sorbitol, Refined Soybean Oil, Hydrogenated Vegetable Oil, Bees Wax, Soy Lecithin, Polyethylene Glycol 4000 & Preservative (214).
Recommended Usage: One capsule daily with water or as directed by Health Professionals.
 Appropriate overages of vitamins are added to compensate the loss of potency on storage.
 Keep out of reach of children.

20x1x10 Capsules (Blister)
M.R.P. 150/-Per 10 Caps.



Ausflu-S 250

Composition :

Each Hard gelatin capsule contains :
 Salmeterol xinafoate IP
 eq. to Salmeterol 50mcg.
 Fluticasone Propionat IP 250mcg.
 Excipients q.s.



Indication : The combination of fluticasone and salmeterol (Advair Diskus, Advair HFA, AirDuo Respiclick) is used to treat difficulty breathing, wheezing, shortness of breath, coughing, and chest tightness caused by asthma.

1x30 Capsules (Rotta Cap)
M.R.P. 432/-Per 30 Caps.

Ausflu-S 500

Composition :

Each hard gelatin capsule contains :
 Salmeterol xinafoate IP
 eq. to Salmeterol 50mcg.
 Fluticasone Propionat IP 500mcg.
 Excipients q.s.



Indication : The combination of fluticasone and salmeterol (Advair Diskus, Advair HFA, AirDuo Respiclick) is used to treat difficulty breathing, wheezing, shortness of breath, coughing, and chest tightness caused by asthma.

1x30 Capsules (Rotta Cap)
M.R.P. 485/-Per 30 Caps.

Budsocare-F6/200

Composition :

Each Hard gelatin capsule contains :
 Formoterol Fumarate
 Dihydrate IP
 Eq. to Formoterol
 Fumarate 6mcg
 Budesnide IP 200mcg
 Excipients q.s.



Indication : Budesonide and formoterol is a combination of two medicines that are used to help control the symptoms of asthma and improve lung function.

1x30 Capsules (Rotta Cap)
M.R.P. 185/-Per 30 Caps.

Budsocare-F6/400

Composition :

Each Hard gelatin capsule contains :
 Formoterol Fumarate
 Dihydrate IP
 Eq. to Formoterol
 Fumarate 6mcg
 Budesnide IP 400mcg
 Excipients q.s.



Indication : Budesonide and formoterol is a combination of two medicines that are used to help control the symptoms of asthma and improve lung function.

1x30 Capsules (Rotta Cap)
M.R.P. 239.90/-Per 30 Caps.

Tioway

Composition :

Each capsule contains :
 Tiotropium Bromide
 Monohydrate IP
 eq. to Tiotropium 18mcg
 Excipients q.s.
 Colour : Approved colour used in empty capsule shells.



Indication : Tiotropium is used to prevent wheezing, shortness of breath, coughing, and chest tightness in patients with chronic obstructive pulmonary disease (COPD, a group of diseases that affect the lungs and airways) such as chronic bronchitis (swelling of the air passages that lead to the lungs) and emphysema (damage to air sacs in the lungs). Tiotropium is in a class of medications called bronchodilators. It works by relaxing and opening the air passages to the lungs to make breathing easier.

1x30 Capsules (Rotta Cap)
M.R.P. 346/-Per 30 Caps.



Softgelatin Capsules



Primocare-500

Composition :

Each soft Gelatin capsule contains :
 Evening Primrose Oil 500 mg
 Evening Primrose Oil is a rich natural sources of Gamma-linolenic Acid (GLA), an essential Omega-6 fatty acid.

Indication : Nerve damage caused by diabetes. Research shows that taking evening primrose oil daily for 6-12 months improves symptoms of nerve damage caused by diabetes.

Osteoporosis. Taking evening primrose oil with fish oil and calcium seems to decrease bone loss and increase bone density in elderly people with osteoporosis.



10x1x15 Capsules (Blister)
M.R.P. 290/-Per 15 Caps.

Primocare-1000

Composition :

Each soft Gelatin capsule contains :
 Evening Primrose Oil 1000 mg
 Evening Primrose Oil is a rich natural sources of Gamma-linolenic Acid (GLA), an essential Omega-6 fatty acid.

Indication : Nerve damage caused by diabetes. Research shows that taking evening primrose oil daily for 6-12 months improves symptoms of nerve damage caused by diabetes.

Osteoporosis. Taking evening primrose oil with fish oil and calcium seems to decrease bone loss and increase bone density in elderly people with osteoporosis.



10x1x15 Capsules (Blister)
M.R.P. 390/-Per 15 Caps.

Re Caution Plus

Composition

Each Soft Gelatin capsule contains :
 Methylcobalamin IP1500 mcg
 Alpha Lipoic Acid USP..... 200 mg
 Folic Acid IP 1.5 mg
 Chromium Picolinate USP200 mcg
 Seleno Methionine USP55 mcg
 Inositol USP100 mg
 Zinc Monomethionine BP 25 mg
 Excipients.....q.s.

Indication : Methylcobalamin + Alpha Lipoic Acid + Folic Acid is used to treat certain types of long-lasting pain caused by damage to nerves as well as to fulfill your nutritional requirements.



10x1x15 Capsules (Blister)
M.R.P. 255/-Per 15 Caps.

Tocopower

Nutritional Information (Approximate Value)
Value per Serving / Per Capsule

Energy	5.64Kcal
Carbohydrates	0.286 gm
Protein	0.278 gm
Fat	0.375 gm
Saturated Fatty Acid	0.246 gm
Polysaturated Fatty Acid	0.135 gm
Monounsaturated Fatty Acid	0.372 gm
Trans Fatty Acid	0.00 gm
Cholesterol	0.00 gm
Each soft gelatin capsule contains:	RDA%*
Tocotrienol(50%)	200 mg **
Excipients	q.s.



10x1x10 Capsules (Blister)
M.R.P. 390/-Per 10 Caps.



INJECTION



Ausnate

Composition

Each vial contains :
Artesunate IP 60mg.

Indication : Artesunate injection is used to treat severe malaria. This medicine is also used together with other medicines (eg, 8-aminoquinoline medicine) to treat severe malaria caused by Plasmodium ovale or Plasmodium vivax. Artesunate belongs to a group of medicines known as antimalarials.



60mg

M.R.P. 230.05/-Per Vial

Ausnate-120

Composition

(a) Each vial contains IP 20mg. artesunate (sterile)
(b) this also contains :
1ml. ampoule of sodium bicarbonate injection IP 5% w/v
5ml. Ampoule of sodium Chloride injection IP 0.9% w/v

Indication : Artesunate injection is used to treat severe malaria. This medicine is also used together with other medicines (eg, 8-aminoquinoline medicine) to treat severe malaria caused by Plasmodium ovale or Plasmodium vivax. Artesunate belongs to a group of medicines known as antimalarials.



120mg.

M.R.P. 453.91/-Per Vial

Amekaday-500

Composition:

Each 2ml contains :
Amikacin Sulphate IP 500mg
equivalent to Amikacin
Methyl Paraben IP 0.018%w/v
(As Preservative)
Propyl Paraben IP 0.02%w/v
(As Preservative)
Water for Injection IP q.s.

Amikacin injection is used to treat certain serious infections that are caused by bacteria such as meningitis (infection of the membranes that surround the brain and spinal cord) and infections of the blood, abdomen (stomach area), lungs, skin, bones, joints, and urinary tract.



500mg/2ml.

M.R.P. 108.20/-Per Vial

Caresil-300

Composition :

Each ml contains :
Clindamycin Phosphate IP
Eq. to Clindamycin 150mg.
Benzyl Alcohol IP 0.945% v/v
(as Preservative)
Water for Injection IP q.s.

Clindamycin injection is used to treat certain types of bacterial infections, including infections of the lungs, skin, blood, bones, joints, female reproductive organs, and internal organs. Clindamycin is in a class of medications called lincomycin antibiotics. It works by slowing or stopping the growth of bacteria.



10x1x2ml.

M.R.P. 127/-Per Ampoule

Caresil-600

Composition :

Each ml contains :
Clindamycin Phosphate IP
Eq. to Clindamycin 150mg.
Benzyl Alcohol IP 0.945% v/v
(as Preservative)
Water for Injection IP q.s.

Clindamycin injection is used to treat certain types of bacterial infections, including infections of the lungs, skin, blood, bones, joints, female reproductive organs, and internal organs. Clindamycin is in a class of medications called lincomycin antibiotics. It works by slowing or stopping the growth of bacteria.



10x1x4ml.

M.R.P. 295/-Per Ampoule

Citirise

Composition :

Each ml Contain :
Citicoline Sodium
Eq. to Citicolin 250 mg.
Water Injection IP q.s.

Citicoline is taken by mouth or given as an injection to help memory loss due to aging, improve vision in people with glaucoma, and help with recovery in stroke patients. It is also used for Alzheimer disease, Parkinson disease, bipolar disorder, lazy eye, and other conditions of the brain.




10x5x2ml.

M.R.P. 145/-Per Ampoule

Impod-C

Composition :

Each ml Contain :
Sterile Imipenem IP 500mg.
sterile cilastatin sodium IP 500mg.
eq. to cilastatin 500mg.



Citicoline is taken by mouth or given as an injection to help memory loss due to aging, improve vision in people with glaucoma, and help with recovery in stroke patients. It is also used for Alzheimer disease, Parkinson disease, bipolar disorder, lazy eye, and other conditions of the brain.

1gm
M.R.P. 2400/-Per Vial

Livogrid-500

Composition :

Each 5ml. contains :
Levetiracetam I.P. 500mg.
Water for Injection I.P. q.s.



Levetiracetam is used in combination with other medications to treat certain types of seizures in adults and children with epilepsy. Levetiracetam is in a class of medications called anticonvulsants. It works by decreasing abnormal excitement in the brain.

25x1x5ml.
M.R.P. 135.60/-Per Ampoule

Methodol[®]

Composition :

Each 1ml. contains :
Methylprednisolone Acetate IP 40mg.
Water for injection q.s.



Levetiracetam is used in combination with other medications to treat certain types of seizures in adults and children with epilepsy. Levetiracetam is in a class of medications called anticonvulsants. It works by decreasing abnormal excitement in the brain.

25x1ml.
M.R.P. 49.87/-Per Vial

Methodol[®]

Composition :

Each 2ml. contains :
Methylprednisolone Acetate IP 80mg.
Water for injection q.s.




Levetiracetam is used in combination with other medications to treat certain types of seizures in adults and children with epilepsy. Levetiracetam is in a class of medications called anticonvulsants. It works by decreasing abnormal excitement in the brain.

25x2ml.
M.R.P. 134.20/-Per Vial

Methyq[®]-2500

Composition :

Each combination pack contains :
One glass ampoule of 1ml. contains
Methylcobalamin I.P. 2500 mcg.
Benzyle Alcohol I.P. 2% w/v.
(As Preservative)
water for injection I.P. q.s.
and one 2ml. Presterilized Disposable Syringe with a Needle.



Indication : Methylcobalamin is used to treat vitamin B12 deficiency. Vitamin B12 is important for the brain and nerves, and for the production of red blood cells. Methylcobalamin is sometimes used in people with pernicious anemia, diabetes, and other conditions.

1x10x1ml.
M.R.P. 122/-Per Ampoule

Nekszole[®]-40

Composition :

Each ml. contains :
Esomeprazole Sodium (Sterile)
eq. to Esomeprazole 40mg.
(As Lyophilized Powder)



Indication : Esomeprazole injection is used to treat gastroesophageal reflux disease (GERD; a condition in which backward flow of acid from the stomach causes heartburn and possible injury of the esophagus [the tube between the throat and stomach]) in adults and children 1 month of age or older who have had damage to their esophagus and are unable to take esomeprazole by mouth. Esomeprazole injection is also used in adults to reduce the risk of further ulcer bleeding after an endoscopy (examination of the inside of the esophagus, stomach, and intestines). Esomeprazole is in a class of medications called proton pump inhibitors. It works by decreasing the amount of acid made in the stomach.

40mg
M.R.P. 148/-Per Vial

Piracetam

Composition :

Each ml. contains :
 Piracetam IP 200mg.
 water for injection IP q.s.

Indication : Piracetam is most commonly used for breath-holding attacks, seizure disorder (epilepsy), dizziness (vertigo), a learning disorder marked by difficulty reading (dyslexia), and a movement disorder often caused by antipsychotic drugs (tardive dyskinesia).



60ml.
 M.R.P. 450/-Per Vial

Teiplan-400

Composition :

Each vial contains :
 Teicoplanin IP 400mg.
 As Sterile freeze dried powder for Reconstitution with sterile water for injection I.P. 3ml.

Teicoplanin is an antibiotic used in the prophylaxis and treatment of serious infections caused by Gram-positive bacteria, including methicillin-resistant Staphylococcus aureus and Enterococcus faecalis. It is a semisynthetic glycopeptide antibiotic with a spectrum of activity similar to vancomycin.



400mg.
 M.R.P. 2400/-Per Vial

Tigamine

Composition :

Each vial contains :
 Tigecycline (Sterile) 50mg.
 (As Lyophilized Powder)

Indication : Tigecycline injection used to treat certain serious infections including community acquired pneumonia (a lung infection that developed in a person who was not in the hospital), skin infections, and infections of the abdomen (area between the chest and the waist).



50mg.
 M.R.P. 2800/-Per Vial

Traxpain®

For Intravenous Use Injection

Composition :

Each 5ml. contains :
 Tranexamic Acid BP 500 gm.
 Water for Injection IP q.s.

Indication : Tranexamic acid injection is used to control or prevent excessive or heavy bleeding during dental procedures in patients with hemophilia.



5x5ml
 M.R.P. 77/-Per Ampoule

Vancofast-500

Composition :

Each vial contains :
 Vancomycin HCl IP
 eq. to Vancomycin 500mg.
 (As sterile freeze dried powder)

Indication : Vancomycin is used to treat an infection of the intestines caused by Clostridium difficile, which can cause watery or bloody diarrhea. This medicine is also used to treat staph infections that can cause inflammation of the colon and small



500mg
 M.R.P. 333.40/-Per Vial

Vancofast-1000

Composition :

Each vial contains :
 Vancomycin HCl IP
 eq. to Vancomycin 250mg.
 (As sterile freeze dried powder)

Indication : Vancomycin is used to treat an infection of the intestines caused by Clostridium difficile, which can cause watery or bloody diarrhea. This medicine is also used to treat staph infections that can cause inflammation of the colon and small



1gm
 M.R.P. 440.37/-Per Vial



FASTEST ACTION against Hyperacid Attack

Raybepoll-20

Composition :

Each vial contains :
Rabeprazole Sodium IP 20mg.
(As Sterile Freeze dried Powder)

Indication : Rabeprazole is used to treat the symptoms of gastroesophageal reflux disease (GERD), a condition in which backward flow of acid from the stomach causes heartburn and possible injury of the esophagus (the tube that connects the throat and stomach) in adults and children 12 year of age and older.

20mg

M.R.P. 210/-Per Vial



Omicer-40

Composition :

Each vial contains :
Sterile Omeprazole IP 40mg.
(In the form of Omeprazole Sodium Suitably buffered)

Indication : Omeprazole Injection 40 mg is indicated primarily for the treatment of Zollinger-Ellison syndrome, and may also be used for the treatment of gastric ulcer, duodenal ulcer and reflux oesophagitis.

40mg.

M.R.P. 120/-Per Vial





Ointment

Desonice

**Composition :**

Desonide USP 0.05 % w/w
 Preservative
 Potassium Sorbate & Sorbic Acid
 Combination eq. to
 Sorbic Acid 0.175% w/w
 In a cream base q.s.

Indication : Desonide is used to treat the redness, swelling, itching, and discomfort of various skin conditions, including psoriasis (a skin disease in which red, scaly patches form on some areas of the body and eczema (a skin disease that causes the skin to be dry and itchy and to sometimes develop red, scaly rashes).

24x1x15gm

M.R.P. 122/-Per Tube

Mupricare

**Composition :**

Mupirocin IP
 in water soluble base

Indication : Mupirocin topical cream is used to treat secondarily infected q.s. traumatic skin lesions due to specific bacteria. Mupirocin topical ointment is used to treat impetigo. This medicine works by killing bacteria or preventing their growth. This medicine is available only with your doctor's prescription.

20x1x5gm

M.R.P. 113.62/-Per Tube

Tacrofast

**Composition :**


Tacrolimus IP 1% w/w
 in an ointment base q.s.

Indication : Tacrolimus ointment is used to treat the symptoms of eczema (atopic dermatitis; a skin disease that causes the skin to be dry and itchy and to sometimes develop red, scaly rashes) in patients who cannot use other medications for their condition or whose eczema has not responded to another medication.

24x1x10gm

M.R.P. 345/-Per Tube

Thromboclid

**Composition :**

Each g contains :
 Heparin Sodium IP 50IU
 Benzyl Nicotinate 2mg
 Sorbic Acid IP 1.97mg
 (Preservative)
 Excipients q.s.

Indication : Benzyl Nicotinate+Heparin is used to treat thrombophlebitis, it is a condition in which presence of blood clot causes inflammation or swelling of a vein. When applied to the affected area, Benzyl Nicotinate+Heparin penetrates the skin to help reduce inflammation and pain, and aid faster healing.

20x1x20gm

M.R.P. 165/-Per Tube

Syrup

Bisalto-M

Composition :

Each 5ml. suspension contains :
 Bilastine 10mg.
 Montelukast Sodium IP 4mg.
 Eqv. to Montelukast 4mg.
 Flavoured syrupy base q.s.
 Colour : Sunset Yellow

Indication : Bilastine+montelukast belongs to a class of medication called 'anti-allergic medication, primarily used to treat allergic conditions like sneezing, runny nose, congestion, stuffy nose or watery eyes.



60ml.
 M.R.P. 185/-Per Bottle

Livogrid®

Composition:

Each ml contains:
 Levetiracetam IP 100mg
 Preservatives
 Methylparaben IP 1.35mg
 Propylparaben IP 0.15mg
 In a flavoured syrupy base

Indication :Levetiracetam is used alone or together with other medicines to help control certain types of seizures (eg, partial seizures, myoclonic seizures, or tonic-clonic seizures) in the treatment of epilepsy. This medicine cannot cure epilepsy and will only work to control seizures for as long as you continue to use it.intestines.



100ml.
 M.R.P. 453.60/-Per Bottle

Raniday®

Composition :

Each 5ml. contains :
 Ranitidine Hydrochloride IP 75mg.
 equivalent to Ranitidine
 Colour : Quinoline Yellow WS
 In a flavoured syrupy base

Indication :Ranitidine is a type of antihistamine that blocks the release of stomach acid. It is used to treat stomach or intestinal ulcers. It can relieve ulcer pain and discomfort, and the heartburn from acid reflux.intestines.



100ml.
 M.R.P. 134.40/-Per Bottle

Piracetam

Composition:

Each ml contains:
 Piracetam IP 500mg.
 Colour : Erythrosine
 In a flavoured syrupy base

Indication :Piracetam is most commonly used for breath-holding attacks, seizure disorder (epilepsy), dizziness (vertigo), a learning disorder marked by difficulty reading (dyslexia), and a movement disorder often caused by antipsychotic drugs (tardive dyskinesia).



100ml.
 M.R.P. 355/-Per Bottle



Protect your Little star with...



Linezolid

SUSPENSION

Linezolid for Oral Suspension

- Act primarily against gram-positive bacteria by inhibiting Protein synthesis
- Linezolid has proved to be successful in the treatment of A wide range of gram-positive infections



Composition:

Each 5ml after reconstitution suspension contains:

Linezolid	I.P.	100mg
Excipients		q.s.

Colour: Sunset Yellow FCF

18gm/30ml.

M.R.P. 118.00/-Per Bottle

SPRAY/SOLUTION

Sensofast

Composition :

Chlorhexidine Gluconate Solution IP Diluted to Chlorhexidine Gluconate 0.2% w/v
 Colour : Brilliant Blue FCF
 In pleasantly flavoured aqueous base q.s.

Indication : Chlorhexidine is an antiseptic and disinfectant. It helps reduce the number of germs (bacteria) in your mouth or on your skin. It can help with: mouth infections, mouth ulcers and gum disease.



100ml.
 M.R.P. 95/-Per Bottle

Sensofast

Composition :

Chlorhexidine Gluconate Solution IP Diluted to Chlorhexidine Gluconate 0.2% w/v
 Colour : Brilliant Blue FCF
 In pleasantly flavoured aqueous base q.s.

Indication : Chlorhexidine is an antiseptic and disinfectant. It helps reduce the number of germs (bacteria) in your mouth or on your skin. It can help with: mouth infections, mouth ulcers and gum disease.



450ml.
 M.R.P. 300/-Per Bottle

Povirub[®]-S Scrub

Composition :

Povidone Iodine IP 7.5% w/v
 (Available Iodine 0.75% w/v)
 Surfactive agent q.s.

Indication : Povidone Iodine is used on the skin to decrease risk of infection. This medicine is also used as a surgical hand scrub and to wash the skin and surface of the eye before surgery to help prevent infections.



500ml.
 M.R.P. 996.80/-Per Bottle

Climazole

Composition:

Clotrimazole IP 1%
 In glycerine IP & Propylene Glycol IP base q.s.

Indication : Climazole Mouth Paint contains clotrimazole, an antifungal drug used to treat or prevent oropharyngeal candidiasis. Candid Mouth Paint 25 ml works by killing the fungus that causes the infection. Candid Mouth Paint 25 ml works by damaging the fungal cell membrane



15ml.
 M.R.P. 99/-Per Bottle



The convenient way to ensure hair Regeneration

Promotes hair growth

- ➔ Prevent hair folicle shrinkage.
- ➔ Enhance hair growth by reduction calcium influx in the cells.
- ➔ Reduce hair shaft shading.
- ➔ Increases thickness of hair folicle.
- ➔ work to relieve hypertension by acting a vasodilator.

MITOXESS-F

60ml.
M.R.P. 699/-Per Bottle

MITOXESS-5 MITOXESS-10

60ml.
M.R.P. 625/-Per Bottle

60ml.
M.R.P. 778/-Per Bottle



Oral Spray/ Nasal Spray & Drop

Care Melatonin

Supplement Facts: 1 Spray = 0.1 ml		
Serving Size: 1 sprays (approx. 0.1 ml)	Amount per Serving	% DV**
Melatonin	0.5mg	**

* % values are based on 2,000 calorie diet
** Daily value not established

Nutritional Information (Approximate Value)	
Serving Size: 1 sprays (approx. 0.1ml)	
Energy	0.092 Kcal
Carbohydrate	0.023 g
Protein	0 g
Fat	0



10x15ml.
M.R.P. 490/-Per Bottle

Momecare

Composition :
Each spray contains :
Mometasone Furoate Monohydrate
eq. to Mometasone Furoate IP 50mcg
Composition :
Mometasone Furoate
Monohydrate eq. to Mometasone
Furoate IP 0.05% w/v
Benzalkonium Chloride IP 0.02% w/v
(as preservative)
Phenyl Ethyl Alcohol USP 0.25%w/v
Excipients q.s.

Indication : Mometasone nasal (nose) spray is used to treat cold-like symptoms caused by allergic rhinitis. This is inflammation of the inside of the nose that can be brought on by hay fever. Mometasone is also used to treat other conditions, such as nasal polyps in adults.



10x10ml.
M.R.P. 475/-Per Bottle

Care-D3



Composition:
Each spray (0.1 ml) delivers:
Vitamin D3 (Cholecalciferol) 400 IU
Excipients q.s.
Serving Size: 1 Spray

Supplement Facts: 1 Spray = 0.1 ml		Nutritional Information (Approx value) / 0.1 ml		
Serving Size 1 spray (Approx. 0.1 ml)	Amount	%RDA*	Amount	Units
Vitamin D3 (Cholecalciferol Solution)	400 IU	100	Energy	0.092
			Carbohydrates	0
			Protein	0
			Fat	0

Indication : Vitamin D3 Oral Spray: Our D3 vitamin spray helps in the absorption of daily supplements, aids immune support, and promotes healthy bones without the hassle of hard to swallow capsules and tablets.

10x15ml.
M.R.P. 470/-Per Bottle

Ligocare



Composition :
Each ml contains :
Lidocaine USP 100mg.
Excipients q.s.

Indication : Lidocaine spray is an anesthetic. It prevents and treats pain caused by some procedures. It can also treat minor burns, scrapes and insect bites. When you use this spray you'll lose feeling in your skin or other tissues.

10x20ml.
M.R.P. 499/-Per Bottle

Nasorub-Oxy

Composition :

Oxymetazoline Hydrochloride IP 0.05% w/v
Benzalkonium chloride Solution IP 0.02% w/v
In an aqueous isotonic solution using distilled water IP

Indication : Oxymetazoline is used for the temporary relief of nasal (of the nose) congestion or stuffiness caused by hay fever or other allergies, colds, or sinus trouble.



25x10ml.

M.R.P. 92/-Per Bottle

Ausflu

Composition :

Each actuation delivers :
Fluticasone Furoate 27.5mcg.
Composition :
Fluticasone Furoate 0.055% w/w
Benzalkonium chloride IP 0.015% w/w (as preservative)
Excipients q.s.

Indication : Fluticasone nasal spray (Flonase Allergy) is used to relieve symptoms of rhinitis such as sneezing and a runny, stuffy, or itchy nose and itchy, watery eyes caused by hay fever or other allergies (caused by an allergy to pollen, mold, dust, or pets).



10x6gm

M.R.P. 345/-Per Bottle

Ausflu-Plus

Composition :

Each Actuation delivers :
Fluticasone Propionate IP 50mcg.
Active :
Fluticasone Propionate IP 0.05% w/v
Preservatives
Benzalkonium Chloride IP 0.01% w/v
Phenyl Ethyl Alcohol IP 0.25% w/v (As Preservative)
Excipients q.s.

Indication : Fluticasone nasal spray is used to relieve symptoms of rhinitis such as sneezing and a runny, stuffy, or itchy nose and itchy, watery eyes caused by hay fever or other allergies (caused by an allergy to pollen, mold, dust, or pets).



10x10ml.

M.R.P. 390/-Per Bottle

Nasorub

Composition :

Xylometazoline Hydrochloride IP 0.01% w/v
Sterile aqueous base q.s.

Indication : This medication is used for temporary relief of congestion in the nose caused by various conditions including the common cold, sinusitis, hay fever, and allergies. It works by narrowing the blood vessels in the nose area, reducing swelling and congestion.



20x10ml.

M.R.P. 57.35/-Per Bottle

Ausflu-AZ

Composition:

Azelastine Hydrochloride IP 140mcg
Fluticasone Propionate IP 50mcg.
Composition :
Azelaastine Hydrochloride IP 0.14% w/v
Fluticasone Propionate IP 0.05% w/v
Benzalkonium chloride IP 0.01% w/v (As Preservative)
Alcohol IP 0.25% w/v (As preservative)
Excipients q.s.

Indication : Azelastine and fluticasone combination nasal spray is used to treat an itchy or runny nose, sneezing, or other symptoms caused by seasonal hay fever (allergic rhinitis).



10x7ml.

M.R.P. 395/-Per Bottle



EYE & EAR DROP

Alcatay

Composition :

Each ml contains :	
Alcaftadine	2.5mg.
Benzalkonium Chloride Solution IP (As preservative)	0.1mg.
Water for injection IP	q.s.

Indication : Alcaftadine eye drops are used to relieve itching of the eyes caused by pollen, ragweed, grass, animal hair, and dander. This medicine is available as an over-the-counter (OTC) product. This product is available in the following dosage forms.



25x1x5ml.

M.R.P. 255/-Per Bottle

Bepotosin

Composition :

Each ml. contains :	
Bepotastine Besilate	15mg.
eq. to Bepotastine	10.70mg.
Benzalkonium chloride solution IP	
Chloride	0.005% w/v
(as preservative)	
Sterile Aqueous base	q.s.

Indication : Bepotastine besilate eye drops is used to treat itching of the eye caused by a condition known as allergic conjunctivitis. It works by preventing the effects of certain inflammatory substances, which are produced by cells in your eyes and sometimes cause allergic reactions.



25x1x5ml.

M.R.P. 240/-Per Bottle

Bimto

Composition :

Bimatoprost	0.03% w/v
Benzalkonium Chloride Solution IP (as preservative)	0.01%v/v
Sterile aqueous base	q.s.

Indication : Bimatoprost is a medicine that treats high pressure in the eye (ocular hypertension) and helps with glaucoma. High pressure can happen when too much fluid builds up inside your eye. Bimatoprost helps to drain the fluid. If not treated, high pressure can damage your optic nerve and cause vision loss or blindness.



25x3ml.

M.R.P. 375/-Per Bottle

Bimto-T

Composition :

Bimatoprost	0.03%w/v
Timolol Maleate IP	
Eq. to Timolol	0.5%w/v
Benzalkonium Chloride Solution IP (as preservative)	0.02% w/v
Sterile aqueous base	q.s.

Indication : Bimatoprost/Timolol works by reducing the production of liquid and also increasing the amount of liquid that is drained. This reduces the pressure inside the eye. Bimatoprost/Timolol eye drops are used to treat high pressure in the eye in adults, including the elderly. This high pressure can lead to glaucoma.



25x1x3ml.

M.R.P. 440/-Per Bottle

Brimcare

Composition :

Brimonidine Tartrate IP	0.2%w/v
Benzalkonium chloride Solution IP	0.005%v/v
Sterile Aqueous base	q.s.

Indication : This combination medication is used to treat high pressure inside the eye due to glaucoma (open-angle type) or other eye diseases (such as ocular hypertension). Lowering high pressure inside the eye helps to prevent blindness.


25x1x5ml.
M.R.P. 210/-Per Bottle

Brimcare-T

Composition :

Brimonidine Tartrate IP	0.2%w/v
Timolol Maleate IP eq. to Timolol	0.5%w/v
Benzalkonium chloride Solution IP (as preservative)	0.005% v/v
Sterile Aqueous base	q.s.

Indication : This combination medication is used to treat high pressure inside the eye due to glaucoma (open-angle type) or other eye diseases (such as ocular hypertension). Lowering high pressure inside the eye helps to prevent blindness.


25x1x5ml.
M.R.P. 387/-Per Bottle

Cycfrain

Composition

Cyclopentolate HCl IP	1% w/v
Chlorbutal IP (as Preservative)	0.5% w/v
Aqueous Base	q.s

Indication : Cyclopentolate is used to dilate (enlarge) the pupil. It is used before eye examinations (such as cycloplegic refraction or ophthalmoscopy). This medicine is available only with your doctor's prescription.


20x1x5ml.
M.R.P. 85.50/-Per Bottle

Brotrace

Composition :

Bromfenac Sodium eq. to Bromfenac	0.09% w/v
Benzalkonium Chloride Solution IP (As preservative)	0.005%v/v
Sterile aqueous base	q.s.

Indication : Bromfenac ophthalmic (eye) solution is used to treat pain or swelling of the eye following cataract surgery. This medicine is a topical nonsteroidal anti-inflammatory drug (NSAID). This medicine is available only with your doctor's prescription.


25x1x5ml.
M.R.P. 167/-Per Bottle

Catadoze

Composition :

Potassium Iodide IP	3.3%w/v
Sodium Chloride IP	0.83%w/v
Calcium Chloride IP	1% w/v
Chlorbutol IP (as preservative)	0.5%w/v
Sterile Aqueous base	q.s.

Indication : Potassium Iodide + Sodium Chloride + Calcium Chloride is a combination of three mineral salts. Altogether, Potassium Iodide, Sodium Chloride, and Calcium Chloride work to improve the eyesight. It is also believed to prevent lenticular opacity (cataract or opacity) and other eye diseases.


25x1x10ml.
M.R.P. 107/-Per Bottle

Doreye

Composition :

Dorzolamide Hydrochloride IP	2%w/v
Benzalkonium chloride IP (as preservative)	0.02%v/v
Sterile aqueous base	q.s.

Indication : Ophthalmic dorzolamide is used to treat glaucoma, a condition in which increased pressure in the eye can lead to gradual loss of vision. Dorzolamide is in a class of medications called carbonic anhydrase inhibitors. It works by decreasing the pressure in the eye.


25x5ml.
M.R.P. 285/-Per Bottle

Doreye-T

Composition :
 Dorzolamide HCl IP 2%w/v
 eq. to Dorzolamide
 Timolol Maleate IP 0.5%w/v
 eq. to Timolol
 Benzalkonium Chloride Solution IP 0.02% v/v
 (as preservative)
 Sterile Aqueous base q.s.

Indication : The combination of dorzolamide and timolol is used to treat eye conditions, including glaucoma and ocular hypertension, in which increased pressure can lead to a gradual loss of vision. Dorzolamide and timolol is used for patients whose eye condition has not responded to another medication



25x1x5ml.
 M.R.P. 490/-Per Bottle

Fluppy

Composition :
 Flurbiprofen Sodium IP 0.03%w/v
 Benzalkonium Chloride Solution IP 0.001%v/v
 (as preservative)
 Sterile Aqueous base q.s.

Indication : Flurbiprofen ophthalmic is used to prevent or reduce changes in the eye that may occur during eye surgery. Flurbiprofen ophthalmic is in a class of medications called nonsteroidal anti-inflammatory drugs (NSAIDs). It works by stopping the release of certain natural substances that cause pain and swelling.



25x1x5ml.
 M.R.P. 74/-Per Bottle

Gatifaux-F

Composition :
 Gatifloxacin eq. to Anhydrous 0.3%w/v
 Gatifloxacin 0.3%w/v
 Flubiprofen Solution IP 0.03%w/v
 Solution IP 0.02%v/v
 (as preservative)
 Sterile Aqueous base q.s.

Indication : Gatifloxacin eye drops are used to treat infections of the eye, such as bacterial conjunctivitis. Gatifloxacin belongs to a group of medicines called fluoroquinolone antibiotics. It works by killing bacteria that cause conjunctivitis. This medicine is available only with your doctor's prescription.



25x1x5ml.
 M.R.P. 105/-Per Bottle

Flumron

Composition :
 Fluorometholone IP 0.1% w/v
 Benzalkonium Chloride Solution IP 0.004%v/v
 (as preservative)
 Sterile Aqueous base q.s.

Indication : Fluorometholone is used to treat eye conditions caused by inflammation (swelling). This medicine is a corticosteroid (steroid medicine) that helps reduce the swelling in the eye. This medicine is available only with your doctor's prescription.



25x1x5ml.
 M.R.P. 152.93/-Per Bottle

Gatifaux

Composition :
 Gatifloxacin eq. to Anhydrous 0.3%w/v
 Gatifloxacin 0.3%w/v
 Benzalkonium Chloride Solution IP 0.01%v/v
 (as preservative)
 Sterile Aqueous base q.s.

Indication : Gatifloxacin eye drops are used to treat infections of the eye, such as bacterial conjunctivitis. Gatifloxacin belongs to a group of medicines called fluoroquinolone antibiotics. It works by killing bacteria that cause conjunctivitis. This medicine is available only with your doctor's prescription.



25x1x5ml.
 M.R.P. 77/-Per Bottle

Hyalcare

Composition :
 Sodium Hyaluronate BP 0.1%w/v
 Sodium Perborate BP 0.028%w/v
 (as preservative)
 Sterile Aqueous base q.s.

Indication : Sodium hyaluronate eye drops are also known as 'artificial tears'. They are used to relieve eye dryness and soreness. They moisten, soothe and lubricate the surface of your eye, making it feel more comfortable.



25x1x10ml.
 M.R.P. 420/-Per Bottle



Ketozox

Composition :

Ketorolac	
Tromethamine IP	0.5% w/v
Benzalkonium Chloride Solution IP (as preservative)	0.02%v/v
Sterile aqueous base	q.s.

Indication : Ketorolac eye drops is used to treat itching caused by seasonal allergic conjunctivitis (an allergy that occurs at only certain times of the year). It is also used to treat pain, burning, and inflammation of the eye following cataract surgery or corneal refractive surgery.

**25x1x5ml.****M.R.P. 60/-Per Bottle**

Ketozox-OF

Composition :

Ketorolac Tromethamine IP	0.5%w/v
Ofloxacin IP	0.3%w/v
Benzalkonium Chloride Solution IP (as preservative)	0.04% v/v
Sterile Aqueous base	q.s.

Indication : Ofloxacin+Ketorolac is used in eye infection. Ofloxacin + Ketorolac is a combination of two medicines: Ofloxacin and Ketorolac which treat eye infections. Ofloxacin is an antibiotic. It kills bacteria by preventing the bacterial cells from dividing and repairing.

**25x1x10ml.****M.R.P. 265/-Per Bottle**

Latofix-T

Composition

Latanoprost IP	0.005% w/v
Timolol Maleate IP eq. to Timolol	0.5%w/v
Benzalkonium chloride Solution IP (as preservative)	0.2% v/v
Sterile aqueous base	q.s.

Indication : Timolol/latanoprost is used is indicated in adults (including the elderly) for the reduction of intraocular pressure (IOP) in patients with open angle glaucoma and ocular hypertension who are insufficiently responsive to topical beta-blockers or prostaglandin analogues.

**2.5ml.****M.R.P. 422/-Per Bottle**

Ketozox-CP

Composition :

Ketorolac Tromethamine IP	0.5%w/v
Phenylephrine HCl. IP	0.0125%w/v
Chlorpheniramine Maleate IP	0.02%w/v
Cellulose IP	0.25%w/v
Phenylmercuric Nitrate IP	0.001% w/v
Sterile aqueous base	q.s.

Indication : Ketorolac and phenylephrine ophthalmic solution is combined with an eye irrigation solution used during cataract surgery or intraocular lens replacement. This is to maintain pupil size by preventing intraoperative miosis (pupil constriction) and to reduce pain after eye surgery.

**25x1x5ml.****M.R.P. 90/-Per Bottle**

Latofix

Composition :

Latanoprost IP	0.005%w/v
Benzalkonium Chloride Solution IP (as preservative)	0.02%v/v
Sterile aqueous base	q.s.

Indication : Latanoprost ophthalmic is used to treat glaucoma (a condition in which increased pressure in the eye can lead to gradual loss of vision) and ocular hypertension (a condition which causes increased pressure in the eye). Latanoprost is in a class of medications called prostaglandin analogs.

**2.5ml.****M.R.P. 399/-Per Bottle**

Loteway

Composition :

Loteprednol Etabonate	0.5% w/v
Benzalkonium chloride Solution IP (as preservative)	0.01%v/v
Sterile aqueous base	q.s.

Indication : Loteprednol is used to treat swelling and pain after cataract surgery (procedure to treat clouding of the lens in the eye). Loteprednol (Alex) is used to reduce eye redness, itching, and swelling caused by seasonal allergies.

**25x1x5ml.****M.R.P. 180/-Per Bottle**

Moiscare

Composition :

Each ml. contains :
 Sodium Hyaluronate BP 1.0mg.
 Eq. to Hyaluronate Acid 50mg.
 D-Panthenol IP
 Stabilized oxychloro
 Complex 0.01%w/v
 (as preservative)
 Sterile Aqueous vehicle q.s.

Indication : Sodium Hyaluronate+D-Panthenol is used in the treatment of dry eyes. Sodium Hyaluronate + D-Panthenol is a combination of two lubricants: Sodium Hyaluronate and D-Panthenol which treat dry eyes. They moisten the eyes and provide temporary relief from burning and discomfort.



25x1x10ml.

M.R.P. 425/-Per Bottle

Moxiyst

Composition :

Moxifloxacin Hydrochloride IP 0.5%w/v
 eq. to Moxifloxacin 0.5%w/v
 Sterile Aqueous Buffered Vehicle q.s.

Indication : Moxifloxacin eye solution is used to treat infections of the eye, including bacterial conjunctivitis. This medicine belongs to a group of medicines called fluoroquinolone antibiotics. It works by killing bacteria that cause conjunctivitis. This medicine is available only with your doctor's prescription.



20x10ml.

M.R.P. 99/-Per Bottle

Moxiyst-D

Composition :

Moxifloxacin HCL IP 0.5%w/v
 eq. to Moxifloxacin 0.5%w/v
 Dexamethasone Sodium
 Phospaite IP eq. to Dexmehtasone
 Phospae 0.1%w/v
 Benzalkonium Chloride
 Solution (As preservative)IP0.02%v/v
 Sterile Aqueous base q.s.

Indication : Moxifloxacin belongs to Dexamethasone+ the class of ocular medications called 'antibiotics' used to treat inflammation associated with bacterial eye infections. It treats eye infections like conjunctivitis (infected conjunctiva) and other inflammatory conditions after eye surgery.



25x1x5ml.

M.R.P. 183/-Per Bottle

Moxiyst-P

Composition :

Moxifloxacin HCl IP 0.5%w/v
 eq. to Moxifloxacin 0.5%w/v
 Prednisolone Acetate IP 1%w/v
 Benzalkonium chloride Solution
 (as preservative) IP 0.02% v/v
 Sterile Aqueous base q.s.

Indication : Moxifloxacin+Prednisolone is used in the treatment of bacterial infections. Moxifloxacin + Prednisolone is a combination of two medicines: Moxifloxacin and Prednisolone. Moxifloxacin is an antibiotic which prevents bacterial cells from dividing and repairing, thereby killing the bacteria. This treats your infection.



25x1x5ml.

M.R.P. 70/-Per Bottle

Moxiyst-KT

Composition :

Moxifloxacin HCl IP 0.5%w/v
 eq. to Moxifloxacin 0.5%w/v
 Ketorolac 0.5%w/v
 Tromethamine IP 0.5%w/v
 Sterile Aqueous base q.s.

Indication : Ketorolac+Moxifloxacin is an ophthalmic (for eye use) medication, primarily used to treat and prevent bacterial eye infections and itchy eyes caused by seasonal allergies. Ketorolac+moxifloxacin also relieves swelling, pain, and burning or stinging sensation after cataract surgery or corneal refractive surgery.



25x1x5ml.

M.R.P. 125/-Per Bottle

Moxiyst-DF

Composition :

Moxifloxacin HCl IP 0.5w/v
 eq. to Moxifloxacin 0.5w/v
 Difluprednate 0.05%w/v
 Sorbic Acid IP 0.1% w/v
 as preservative
 Sterile Aqueous base q.s.

Indication : Moxifloxacin+ Difluprednate is used in the treatment of Bacterial eye infections. Moxifloxacin and Difluprednate is an antibiotic which prevents bacterial cells from dividing and repairing, thereby killing the bacteria.



25x1x5ml.

M.R.P. 210/-Per Bottle




Moxiyst-LP

Composition :

Moxifloxacin HCl IP	0.5%w/v
eq. to Moxifloxacin	
Loteprednol Etabonate	0.5%w/v
Benzalkonium Chloride Solution	
(As preservative)IP	0.01%v/v
Sterile Aqueous base	q.s.

Indication : Loteprednol etabonate + Moxifloxacin is used for the treatment of bacterial infection and inflammation of the eye. Do not skip any doses and finish the full course of treatment even if you feel better. Stopping it early may make the infection come back and harder to treat.



25x1x5ml.

M.R.P. 175/-Per Bottle

Natadoz

Composition

Natamycin IP	5% w/v
Benzalkonium chloride Soution IP	0.01% v/v
(as preservative)	
Sterile aqueous base	q.s.

Indication : TNatamycin belongs to the group of medicines called antifungals. It is used to treat some types of fungus infections of the eye. Natamycin is available only with your doctor's prescription.



25x1x5ml.


M.R.P. 130.87/-Per Bottle

Prednitore

Composition :

Prednisolone Acetate IP	1% w/v
Benzalkonium chloride Solution IP	0.02%v/v
(as preservative)	
Aqueous base	q.s.

Indication : Ophthalmic prednisolone reduces the irritation, redness, burning, and swelling of eye inflammation caused by chemicals, heat, radiation, infection, allergy, or foreign bodies in the eye. It sometimes is used after eye surgery. Prednisolone is in a class of medications called steroids.



25x1x10ml.

M.R.P. 65.51/-Per Bottle

Nepalook

Composition :

Nepafenac	0.1% w/v
Benzalkonium chloride solution IP	0.005% v/v
(as preservative)	
Aqueous Vehicle	q.s.

Indication : Ophthalmic nepafenac is used to treat eye pain, redness, and swelling in patients who are recovering from cataract surgery (procedure to treat clouding of the lens in the eye). Nepafenac is in a class of medications called nonsteroidal anti-inflammatory drugs (NSAIDs)



25x1x5ml.

M.R.P. 270/-Per Bottle

Oloease

Composition :

Olopatadine Hydrochloride IP	
Eq. to Olopatadine	0.1%w/v
Benzalkonium chloride Solution IP	0.02% v/v
(as preservative)	
Sterile Aqueous base	q.s.

Indication : Olopatadine Hydrochloride ophthalmic (eye) drops is used to treat itching of the eye caused by a condition known as allergic conjunctivitis (pink eye). This medicine is also used to treat eye itching or redness caused by pollen, ragweed, grass, animal hair, or dander.



25x1x5ml.


M.R.P. 145/-Per Bottle

Polyemoise

Composition :

Polyethylene Glyco 400	0.4%w/v
Prppylene Glycol IP	0.3%w/v
Sterile Aqueous base	q.s.

Indication : This medication is used to relieve dry, irritated eyes. Common causes for dry eyes include wind, sun, heating/air conditioning, computer use/reading, and certain medications.



20x1x10ml.

M.R.P.289/-Per Bottle

Re-Tearoute

Composition :

Hydroxypropyl Methyl Cellulose IP 0.3% w/v
Sterile aqueous base q.s.

Indication : It is used to relieve dryness and irritation caused by reduced tear flow. It helps prevent damage to the eye in certain eye diseases. Hydroxypropyl methylcellulose may also be used to moisten hard contact lenses and artificial eyes. In addition, it may be used in certain eye examinations



25x1x5ml.

M.R.P. 250/-Per Bottle

Tavapost-T

Composition :

Travoprost IP 0.004%w/v
Timolol Maleate IP eq. to Timolol 0.5% w/v
Benzalkonium chloride solution IP 0.02% v/v (as preservative)
Sterile Aqueous base q.s.

Indication : Travoprost/Timolol eye drops are used to treat high pressure in the eye in adults, including the elderly. This pressure can lead to an illness called glaucoma.



25x1x2.5ml.

M.R.P. 394/-Per Bottle

Timoday-D

Composition :

Timolol Maleate IP eq. to Timolol 0.5% w/v
Dexamethasone Phosphate 0.1% w/v
Benzalkonium Chloride Solution IP 0.02% v/v (as preservative)
Sterile Aqueous base q.s.

Indication : This medication is used to temporarily relieve itching eyes caused by seasonal allergies. It is also used to prevent and treat eye swelling due to a certain type of eye surgery (cataract removal).



25x1x5ml.

M.R.P. /-Per Bottle

Travoprost

Composition :

Composition :
Travoprost HydrochloridelP 0.004%w/v
Benzalkonium ChloridelP 0.02%v/v (as preservative)
Sterile Aqueous base q.s.

Indication : Travoprost eye drops are used to treat increased pressure in the eye caused by open-angle glaucoma or hypertension of the eye. Both conditions are caused by high pressure in the eye which can lead to pain or vision changes. This medicine is a prostaglandin.



35x2.5ml.

M.R.P. 289/-Per Bottle

Timoday

Composition :

Timolol Maleate IP eq. to Timolol 0.5%w/v
Benzalkonium chloride Solution IP 0.02%v/v (as preservative)
Sterile Aqueous base q.s.

Indication : Timolol is used to treat glaucoma, a condition in which increased pressure in the eye can lead to gradual loss of vision. Timolol is in a class of medications called beta-blockers. It works by decreasing the pressure in the eye.



25x1x5ml.

M.R.P. 66.90/-Per Bottle

Tropacyl-P

Composition

Tropicamide IP 0.8% w/v
Phenylephrine HCl 5% w/v
Chlorbutol IP 0.5%w/v (as preservative)
Sterile aqueous base q.s.

Indication : Tropicamide is used to dilate (enlarge) the pupil so that the doctor can see into the back of your eye. It is used before eye examinations, such as cycloplegic refraction and examination of the fundus of the eye. Tropicamide may also be used before and after eye surgery.



25x1x5ml.

M.R.P. 125/-Per Bottle



Tobraclide-F

Composition :

Fluorometholone IP	0.1%w/v
Tobramycin Sulphate USP	eq. to Tobramycin
Benzalkonium chloride Solution IP	0.3%w/v
Sterile aqueous base	0.004%v/v
	q.s.

Indication : Tobramycin + Fluorometholone is used for the treatment of bacterial skin infections. It should be applied to the affected areas as a thin film, two or three times daily, or as advised by your doctor. Do not use it more often or for longer than advised by your doctor.



Tobraclide

Composition :

Tobramycin Sulphate U.S.P.	
Eq. to Tobramycin	0.3% w/v
Benzalkonium chloride Solution I.P.	0.02% v/v
(As preservative)	
Sterile Aqueous base	q.s.

Indication : Tobramycin Ophthalmic is used in the eye to treat bacterial infections of the eye. Tobramycin works by killing bacteria. Ophthalmic tobramycin may be used alone or with other medicines for eye infections. Either the drops or the ointment form of this medicine may be used alone during the day.

25x1x5ml.

M.R.P. 181/-Per Bottle

25x1x5ml.

M.R.P. 98/-Per Bottle

EYE/EAR DROP

RESPULES/ INFUSION

Budsocare-0.5

Composition :

Each 2ml. Respules contains :
Budesonide IP 0.5mg



Indication : Budesonide suspension (liquid) for oral inhalation (Pulmicort Respules) is used in children 12 months to 8 years of age. Budesonide belongs to a class of medications called corticosteroids. It works by decreasing swelling and irritation in the airways to allow for easier breathing.

4x5x2ml.

M.R.P. 118.75/-

Budsocare-1

Composition :

Each 2ml. Respules contains :
Budesonide IP 1mg



Indication : Budesonide suspension (liquid) for oral inhalation (Pulmicort Respules) is used in children 12 months to 8 years of age. Budesonide belongs to a class of medications called corticosteroids. It works by decreasing swelling and irritation in the airways to allow for easier breathing.

4x5x2ml.

M.R.P. 147.15/-

L-Carebex

Composition :

Each 2.5ml. contains :
Levosaltamol Sulphate IP
eq. to Levosaltamol 1.25mg
Ipratropium Bromide IP
eq. to Ipratropium Bromide
(Anhydrous) 500mg.
(An Isotonic Solution) q.s.



Indication : levosaltamol+ ipratropium Bromide Is A Combination Medication Primarily Used To Treat Or Prevent Symptoms Of Chronic Obstructive Pulmonary Disease (copd) Likes Shortness Of Breath, Coughing Or Wheezing (whistling Sound While Breathing).

4x5x5ml.

M.R.P. 123/-

Parest®

Composition :

Each 100ml. contains :
Paracetamol IP 1000mg.
Water for injection IP q.s.



Indication : Paracetamol is recommended as a first-line treatment for pain and fever in paediatric patients. Intravenous (IV) infusions are recommended to be administered as a 15-min infusion to minimize local tissue trauma and related pain.





100ml.

M.R.P. 425/-Per Bottle

UPcoming Products

1	Clonazepam-5
2	Clonazepam-10
3	Everlimus-0.5
4	Rasgocin-1
5	Rasgocin-2
6	Lercanidipine-10mg.
7	Melphalan-2
8	Cypcare Syrup (mango, orange, kiwi, & Pineapple Flavour)
9	Caldicare joint
10	Imeglimin Hydrochloride
11	Bupropion Tablets 150mg
12	Danazol 200mg Capsules
13	Tadalafil Tablets 10mg.
14	Etodolac Extended Release Tablets
15	Clarithromycin IP 1% gel
16	Clarithromycin Tablets IP 500mg.
17	Levodopa And Carbidopa Tablets IP
18	Thiamine Hydrochloride Tablet 100mg
19	Erythropoietin Injection
20	Saxagliptin Tablets
21	Naltrexone 50mg tablets
22	Apremilast Tablets 30mg
23	Verapamil 40mg Tablets
24	Levocloperastine Hydrochloride + Fendizoate Syup
25	levodropropizine and chlorpheniramine maleate syrup
26	Levocetirizine & Ambroxol Hydrochloride Tablets
27	Brinzolamide & Brimonidine Eye Drops
28	Parecoxib Tablet


New Product Launch





Care Formulation Labs

Care Formulation Labs Pvt. Ltd
(An ISO 9001 : 2015 Certified Company)
F-14/37 2nd Floor Model Town-II, Delhi-110009
Customer Care No.: 011-47878729
E-Mail: cflabs@gmail.com
Website : www.careformulationlabs.com



August-2023